



A NOTE ON A GRAPH RELATED TO THE COMAXIMAL IDEAL GRAPH OF A COMMUTATIVE RING

SUBRAMANIAN VISWESWARAN* AND JAYDEEP PAREJIYA

Communicated by S. Alikhani

ABSTRACT. The rings considered in this article are commutative with identity which admit at least two maximal ideals. This article is inspired by the work done on the comaximal ideal graph of a commutative ring. Let R be a ring. We associate an undirected graph to R denoted by $\mathcal{G}(R)$, whose vertex set is the set of all proper ideals I of R such that $I \not\subseteq J(R)$, where $J(R)$ is the Jacobson radical of R and distinct vertices I_1, I_2 are adjacent in $\mathcal{G}(R)$ if and only if $I_1 \cap I_2 = I_1 I_2$. The aim of this article is to study the interplay between the graph-theoretic properties of $\mathcal{G}(R)$ and the ring-theoretic properties of R .

1. INTRODUCTION

The rings considered in this article are commutative with identity which admit at least two maximal ideals. Let R be a ring. We denote the set of all maximal ideals of R by $Max(R)$. This article is inspired by the interesting theorems proved by M. Ye and T. Wu in [15]. Motivated

<http://dx.doi.org/10.29252/asta.4.1.57>

MSC(2010): Primary 13A15; Secondary 05C25.

Keywords: Comaximal ideal graph of a commutative ring, complete graph, von Neumann regular ring, bipartite graph, clique number.

Received: 11 November 2017, Accepted: 19 April 2018

* Corresponding author

by the research work done on the comaximal graph of a ring in [9, 10, 11, 13, 14] and on the annihilating-ideal graph of a ring in [5, 6], M. Ye and T. Wu in [15] introduced and studied a graph structure on a ring R whose vertex set is the set of all proper ideals I of R such that $I \not\subseteq J(R)$, where $J(R)$ is the Jacobson radical of R and distinct vertices I_1 and I_2 are adjacent if and only if $I_1 + I_2 = R$. M. Ye and T. Wu called the graph introduced and investigated by them in [15] as the *comaximal ideal graph* of R and denoted it by $\mathcal{C}(R)$ and they investigated the influence of certain graph parameters of $\mathcal{C}(R)$ on the ring structure of R .

We denote the cardinality of a set A by $|A|$. Let R be a ring with $|Max(R)| \geq 2$. In this article, we introduce a graph structure on R , denoted by $\mathcal{G}(R)$, is an undirected graph whose vertex set is the set of all proper ideals I of R such that $I \not\subseteq J(R)$ and distinct vertices I_1, I_2 are adjacent in $\mathcal{G}(R)$ if and only if $I_1 \cap I_2 = I_1 I_2$. The graphs considered in this article are undirected and simple. We denote the set of all vertices of a graph G by $V(G)$ and the set of all edges of G by $E(G)$. A subgraph H of a graph G is said to be a *spanning subgraph* of G if $V(G) = V(H)$. Let R be a ring. If I_1, I_2 are ideals of R such that $I_1 + I_2 = R$, then we know from [2, Proposition 1.10(i)] that $I_1 \cap I_2 = I_1 I_2$. Let I_1, I_2 be proper ideals of a ring such that $I_i \not\subseteq J(R)$ for each $i \in \{1, 2\}$. If I_1 and I_2 are adjacent in $\mathcal{C}(R)$, then they are adjacent in $\mathcal{G}(R)$. This shows that $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$. Hence, it is natural to compare the graph-theoretic properties of $\mathcal{G}(R)$ with that of the graph-theoretic properties of $\mathcal{C}(R)$. The aim of this article is to study the interplay between the graph-theoretic properties of $\mathcal{G}(R)$ and the ring-theoretic properties of R .

First, it is useful to recall certain definitions and results from commutative ring theory that are used in this article. Let R be a ring. The nil radical of R is denoted by $nil(R)$. A ring R is said to be *reduced* if $nil(R) = (0)$. Recall from [7, Exercise 16, page 111] that a ring R is said to be *von Neumann regular* if for each $a \in R$, there exists $b \in R$ such that $a = a^2 b$. A principal ideal ring is said to be a *special principal ideal ring* (SPIR) if R has a unique prime ideal. If \mathfrak{m} is the only prime ideal of a special principal ideal ring R , then we denote it by mentioning that (R, \mathfrak{m}) is a SPIR. If (R, \mathfrak{m}) is a SPIR, then \mathfrak{m} is nilpotent. Let (R, \mathfrak{m}) be a SPIR which is not a field. Let $n \geq 2$ be least with the property that $\mathfrak{m}^n = (0)$. Then it follows from the proof of (iii) \Rightarrow (i) of [2, Proposition 8.8] that $\{\mathfrak{m}^i | i \in \{1, \dots, n-1\}\}$ is the set of all nonzero proper ideals of R . A ring with a unique maximal ideal is referred to as a *quasilocal* ring. A ring which admits only a finite number of maximal ideals is referred to as a *semiquasilocal* ring. A Noetherian quasilocal (respectively, semiquasilocal) ring is referred to as a *local* (respectively, *semilocal*) ring. For a ring R , we denote the set of all units of R by $U(R)$ and the set of all nonunits of R by $NU(R)$. The Krull dimension of a ring R is simply denoted by $dim R$. We use $Spec(R)$ to denote the set of all prime ideals of a ring R . We use \subset to denote proper inclusion. For any $n \geq 2$, we denote the ring of integers modulo n by \mathbb{Z}_n .

Next, it is useful to recall the following results from graph theory before we give an account of the results that are proved in this article. Let $G = (V, E)$ be a graph. Let $a, b \in V, a \neq b$. Recall that the *distance* between a and b , denoted by $d(a, b)$ is defined as the length of a shortest path in G between a and b if there exists such a path in G ; otherwise, we define $d(a, b) = \infty$. We define $d(a, a) = 0$. The *diameter* of G , denoted by $diam(G)$, is defined as $diam(G) = \sup\{d(a, b) | a, b \in V\}$ [3]. A graph $G = (V, E)$ is said to be *connected* if for any distinct $a, b \in V$, there exists a path in G between a and b [3]. Let $G = (V, E)$ be a connected graph. Let $a \in V$. Then the *eccentricity* of a , denoted by $e(a)$, is defined as $e(a) = \sup\{d(a, b) | b \in V\}$. The *radius* of G , denoted by $r(G)$, is defined as $r(G) = \min\{e(a) | a \in V\}$. A simple graph $G = (V, E)$ is said to be *complete* if every pair of distinct vertices of G are adjacent in G [3, Definition 1.1.11]. Let $n \in \mathbb{N}$. A complete graph on n vertices is denoted by K_n . A graph $G = (V, E)$ is said to be *bipartite* if V can be partitioned into two nonempty subsets V_1 and V_2 such that each edge of G has one end in V_1 and the other in V_2 . A bipartite graph with vertex partition V_1 and V_2 is said to be *complete* if each element of V_1 is adjacent to every element of V_2 .

Let $G = (V, E)$ be a graph. Recall from [3, Definition 1.2.2] that a *clique* of G is a complete subgraph of G . The *clique number* of G , denoted by $\omega(G)$, is defined as the largest integer $n \geq 1$ such that G contains a clique on n vertices [3, page 185]. We set $\omega(G) = \infty$ if G contains a clique on n vertices for all $n \geq 1$. Recall from [3, page 129] that a *vertex coloring* of G is a map $f : V \rightarrow S$, where S is a set of distinct colors. A vertex coloring $f : V \rightarrow S$ is said to be *proper*, if adjacent vertices of G receive different colors of S ; that is, if a and b are adjacent vertices of G , then $f(a) \neq f(b)$. The *chromatic number* of G , denoted by $\chi(G)$, is the minimum number of colors needed for a proper vertex coloring of G [3, Definition 7.1.2]. It is well-known that for any graph G , $\omega(G) \leq \chi(G)$.

Let R be a ring with $|Max(R)| \geq 2$. It is shown in Section 2 of this article that $\mathcal{G}(R)$ is connected and $diam(\mathcal{G}(R)) \leq 3$. With the hypothesis that $J(R) = (0)$, it is shown that $\mathcal{G}(R)$ is complete if and only if R is von Neumann regular. Some classes of rings R are provided such that $diam(\mathcal{G}(R))$ is either 2 or 3. Moreover, some examples are given to illustrate the results proved in this section.

Let R be a ring with $|Max(R)| \geq 2$. It is proved in Section 3 of this article that $\mathcal{G}(R)$ is a finite bipartite graph if and only if $R \cong F_1 \times F_2$ as rings, where F_i is a field for each $i \in \{1, 2\}$.

Let R be a ring such that $|Max(R)| \geq 2$. With the assumption that $J(R) = (0)$, it is proved in Section 4 of this article that $\omega(\mathcal{G}(R)) < \infty$ if and only if there exist $n \geq 2$ and fields F_1, F_2, \dots, F_n such that $R \cong F_1 \times F_2 \times \dots \times F_n$ as rings. Moreover, in such a case, it is verified that $\omega(\mathcal{G}(R)) = \chi(\mathcal{G}(R)) = 2^n - 2$. Moreover, an example of a ring R with $|Max(R)| = 2$ is provided such that $\omega(\mathcal{G}(R)) = 5 < \chi(\mathcal{G}(R)) = 6$.

2. Some basic properties of $\mathcal{G}(R)$

Let R be a ring such that $|Max(R)| \geq 2$. The aim of this section is to study some basic properties of $\mathcal{G}(R)$.

Proposition 2.1. *Let R be a ring such that $|Max(R)| \geq 2$. Then $\mathcal{G}(R)$ is connected with $diam(\mathcal{G}(R)) \leq 3$.*

Proof. It is already noted in the introduction that $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$. We know from [15, Theorem 2.4] that $\mathcal{C}(R)$ is connected with $diam(\mathcal{C}(R)) \leq 3$. Therefore, $\mathcal{G}(R)$ is connected with $diam(\mathcal{G}(R)) \leq 3$. \square

Proposition 2.2. *Let R be a ring such that $|Max(R)| \geq 2$. If either $J(R) = (0)$ or $J(R)$ is a prime ideal of R , then $diam(\mathcal{G}(R)) \leq 2$.*

Proof. Let $I_1, I_2 \in V(\mathcal{G}(R))$ be such that $I_1 \neq I_2$. We claim that $d(I_1, I_2) \leq 2$ in $\mathcal{G}(R)$. This is clear if I_1, I_2 are adjacent in $\mathcal{G}(R)$. Hence, we can assume that I_1 and I_2 are not adjacent in $\mathcal{G}(R)$. We claim that $I_1 \cap I_2 \not\subseteq J(R)$. We are assuming that either $J(R) = (0)$ or $J(R)$ is a prime ideal of R . We consider the following cases.

Case(i) $J(R) = (0)$

Observe that $I_1 \cap I_2 \neq (0)$. For if $I_1 \cap I_2 = (0)$, then $I_1 \cap I_2 = I_1 I_2 = (0)$ and this implies that I_1 and I_2 are adjacent in $\mathcal{G}(R)$. This is in contradiction to the assumption that I_1 and I_2 are not adjacent in $\mathcal{G}(R)$. Therefore, $I_1 \cap I_2 \neq (0)$. As $J(R) = (0)$, we get that $I_1 \cap I_2 \not\subseteq J(R)$.

Case(ii) $J(R)$ is a prime ideal of R

Let $i \in \{1, 2\}$. As $I_i \in V(\mathcal{G}(R))$, we obtain that $I_i \not\subseteq J(R)$. Since $J(R) \in Spec(R)$, we obtain from [2, Proposition 1.11(ii)] that $I_1 \cap I_2 \not\subseteq J(R)$.

Thus $I_1 \cap I_2 \not\subseteq J(R)$. Hence, there exists $\mathfrak{m} \in Max(R)$ such that $I_1 \cap I_2 \not\subseteq \mathfrak{m}$. This implies that $I_i \not\subseteq \mathfrak{m}$ for each $i \in \{1, 2\}$. Therefore, $I_i + \mathfrak{m} = R$ for each $i \in \{1, 2\}$. Hence, $I_1 - \mathfrak{m} - I_2$ is a path of length two in $\mathcal{C}(R)$ and this is also a path in $\mathcal{G}(R)$ since $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$. This proves that $d(I_1, I_2) \leq 2$ in $\mathcal{G}(R)$ for any $I_1, I_2 \in V(\mathcal{G}(R))$. Therefore, $diam(\mathcal{G}(R)) \leq 2$. \square

Remark 2.3. (i) Let R be a ring such that $|Max(R)| \geq 2$. If $J(R) \in Spec(R)$, then $diam(\mathcal{C}(R)) \leq 2$.

(ii) Let R be a ring such that $|Max(R)| \geq 3$. Suppose that there exist $I_1, I_2 \in V(\mathcal{C}(R))$ such that I_1 and I_2 are not adjacent in $\mathcal{C}(R)$ and $I_1 I_2 \subseteq J(R)$. Then $diam(\mathcal{C}(R)) = 3$.

Proof. (i) Let $I_1, I_2 \in V(\mathcal{C}(R))$ be such that $I_1 \neq I_2$ and I_1, I_2 are not adjacent in $\mathcal{C}(R)$. Using the hypothesis that $J(R) \in Spec(R)$, it is noted in the proof of Case(ii) of Proposition

2.2 that there exists $\mathfrak{m} \in \text{Max}(R)$ such that $I_1 - \mathfrak{m} - I_2$ is a path of length two between I_1 and I_2 in $\mathcal{C}(R)$. Hence, $\text{diam}(\mathcal{C}(R)) \leq 2$.

(ii) Now, by assumption $I_1, I_2 \in V(\mathcal{C}(R))$ are such that I_1, I_2 are not adjacent in $\mathcal{C}(R)$ and $I_1 I_2 \subseteq J(R)$. Since $I_1 \not\subseteq J(R)$, $I_1 I_2 \subseteq J(R)$, and $J(R)$ is a radical ideal of R , it follows that $I_1 \neq I_2$. We claim that $d(I_1, I_2) = 3$ in $\mathcal{C}(R)$. We know from [15, Theorem 2.4] that $\mathcal{C}(R)$ is connected and $\text{diam}(\mathcal{C}(R)) \leq 3$. We verify that there exists no path of length two between I_1 and $I_2 \in \mathcal{C}(R)$. Suppose that there exists a path of length two between I_1 and I_2 in $\mathcal{C}(R)$. Thus there exists $I \in V(\mathcal{C}(R))$ such that $I_1 - I - I_2$ is a path of length two in $\mathcal{C}(R)$. Hence, $I_i + I = R$ for each $i \in \{1, 2\}$. Let $\mathfrak{m} \in \text{Max}(R)$ be such that $I \subseteq \mathfrak{m}$. Note that $I_i + \mathfrak{m} = R$ for each $i \in \{1, 2\}$. As $I_1 I_2 \subseteq J(R) \subset \mathfrak{m}$, we get that either $I_1 \subseteq \mathfrak{m}$ or $I_2 \subseteq \mathfrak{m}$. Therefore, $I_i + \mathfrak{m} \neq R$ for at least one $i \in \{1, 2\}$. This is a contradiction. Therefore, $d(I_1, I_2) \geq 3$ in $\mathcal{C}(R)$ and so, we obtain that $\text{diam}(\mathcal{C}(R)) = 3$. \square

Let R be a ring such that $|\text{Max}(R)| \geq 2$. Suppose that $J(R) = (0)$. In Theorem 2.4, we characterize rings R such that $\mathcal{G}(R)$ is complete.

Theorem 2.4. *Let R be a ring with $|\text{Max}(R)| \geq 2$. Suppose that $J(R) = (0)$. Then the following statements are equivalent:*

- (i) $\mathcal{G}(R)$ is complete.
- (ii) R is von Neumann regular.

Proof. (i) \Rightarrow (ii) Assume that $\mathcal{G}(R)$ is complete. By hypothesis, $J(R) = (0)$. Since $\text{nil}(R) \subseteq J(R)$, we obtain that $\text{nil}(R) = (0)$. Hence, R is reduced. Therefore, to prove R is von Neumann regular, it follows from (d) \Rightarrow (a) of [7, Exercise 16, page 111] that it is enough to show that $\dim R = 0$. Let $\mathfrak{p} \in \text{Spec}(R)$. Let $a \in R \setminus \mathfrak{p}$. We claim that $\mathfrak{p} + Ra = R$. If $Ra = Ra^2$, then $a = ra^2$ for some $r \in R$. Hence, $a(1 - ra) = 0 \in \mathfrak{p}$. As $a \notin \mathfrak{p}$, we obtain that $1 - ra \in \mathfrak{p}$. This implies that $\mathfrak{p} + Ra = R$. Suppose that $Ra \neq Ra^2$. Then $Ra, Ra^2 \in V(\mathcal{G}(R))$. Since $\mathcal{G}(R)$ is complete, we obtain that Ra and Ra^2 are adjacent in $\mathcal{G}(R)$. Therefore, $Ra \cap Ra^2 = Ra^3$. Thus $Ra^2 = Ra^3$ and this implies that $a^2 = sa^3$ for some $s \in R$. Hence, $a^2(1 - sa) = 0$. From $a^2 \notin \mathfrak{p}$, we get that $1 - sa \in \mathfrak{p}$. Therefore, $\mathfrak{p} + Ra = R$. This proves that \mathfrak{p} is a maximal ideal of R for any $\mathfrak{p} \in \text{Spec}(R)$. Therefore, $\dim R = 0$ and so, R is von Neumann regular.

(ii) \Rightarrow (i) We are assuming that R is von Neumann regular. Let $a \in R$. We know from (1) \Rightarrow (3) of [7, Exercise 29, page 113] that there exist a unit $u \in R$ and an idempotent element e of R such that $a = ue$. Using this fact, it can be shown that any ideal of R is a radical ideal of R . Let $I_1, I_2 \in V(\mathcal{G}(R))$ with $I_1 \neq I_2$. We know from [2, Exercise 1.13(iii), page 9] that $\sqrt{I_1 I_2} = \sqrt{I_1 \cap I_2}$. Since for any ideal I of R , $I = \sqrt{I}$, we obtain that $I_1 \cap I_2 = I_1 I_2$. Hence, I_1 and I_2 are adjacent in $\mathcal{G}(R)$. This proves that $\mathcal{G}(R)$ is complete. \square

Example 2.5. Let F_n be a field for each $n \in \mathbb{N}$. Let $R = \prod_{n=1}^{\infty} F_n$. Then $\mathcal{G}(R)$ is complete but $\text{diam}(\mathcal{C}(R)) = 3$.

Proof. Note that R is von Neumann regular. We know from (a) \Rightarrow (d) of [7, Exercise 16, page 111] that R is reduced and $\text{dim} R = 0$. Hence, $J(R) = \text{nil}(R) = (0)$. It now follows from (ii) \Rightarrow (i) of Theorem 2.4 that $\mathcal{G}(R)$ is complete. We next verify that $\text{diam}(\mathcal{C}(R)) = 3$. We know from [15, Theorem 2.4] that $\mathcal{C}(R)$ is connected and $\text{diam}(\mathcal{C}(R)) \leq 3$. Note that $\text{Max}(R)$ is infinite and $J(R) = (0, 0, 0, \dots)$. Let $I_1 = \{(\alpha_1, \alpha_2, \alpha_3, \dots) \in R \mid \alpha_i = 0 \text{ for all } i \in \mathbb{N} \text{ with } i \text{ odd}\}$ and let $I_2 = \{(0, \alpha_2, \alpha_3, \alpha_4, \dots) \in R \mid \alpha_i = 0 \text{ for all } i \in \mathbb{N} \text{ with } i \text{ even}\}$. Observe that $I_1, I_2 \in V(\mathcal{C}(R))$ with $I_1 \neq I_2$ and $I_1 + I_2 \neq R$. Therefore, I_1 and I_2 are not adjacent in $\mathcal{C}(R)$. As $I_1 I_2 = (0, 0, 0, \dots) \subseteq J(R)$, we obtain from the proof of Remark 2.3(ii) that $d(I_1, I_2) \geq 3$ in $\mathcal{C}(R)$ and so, it follows that $\text{diam}(\mathcal{C}(R)) = 3$. \square

Corollary 2.6. Let R be a ring such that $|\text{Max}(R)| \geq 2$. Suppose that $J(R) = (0)$. If R is not von Neumann regular, then $\text{diam}(\mathcal{G}(R)) = 2$.

Proof. Let $\mathfrak{m} \in \text{Max}(R)$. As $|\text{Max}(R)| \geq 2$, it follows that $\mathfrak{m} \in V(\mathcal{G}(R))$. Let $\mathfrak{m}, \mathfrak{m}' \in \text{Max}(R)$ be such that $\mathfrak{m} \neq \mathfrak{m}'$. As $\mathfrak{m} + \mathfrak{m}' = R$, we obtain that \mathfrak{m} and \mathfrak{m}' are adjacent in $\mathcal{C}(R)$ and so, they are adjacent in $\mathcal{G}(R)$. Hence, $\mathcal{G}(R)$ admits at least one edge. We know from Proposition 2.1 that $\mathcal{G}(R)$ is connected. We are assuming that $J(R) = (0)$ and R is not von Neumann regular. Therefore, we obtain from (i) \Rightarrow (ii) of Theorem 2.4 that $\text{diam}(\mathcal{G}(R)) \geq 2$. Since $J(R) = (0)$, we know from Proposition 2.2 that $\text{diam}(\mathcal{G}(R)) \leq 2$. Therefore, we get that $\text{diam}(\mathcal{G}(R)) = 2$. \square

Note that $J(\mathbb{Z}) = (0)$ and \mathbb{Z} is not von Neumann regular. Hence, we obtain from Corollary 2.6 that $\text{diam}(\mathcal{G}(\mathbb{Z})) = 2$. Let R be a principal ideal domain such that $|\text{Max}(R)| \geq 2$. We verify in Remark 2.7 that $\mathcal{C}(R) = \mathcal{G}(R)$.

Remark 2.7. Let R be a principal ideal domain such that $|\text{Max}(R)| \geq 2$. Then $\mathcal{C}(R) = \mathcal{G}(R)$.

Proof. Let T be any ring with $|\text{Max}(T)| \geq 2$. It is already noted in the introduction that $\mathcal{C}(T)$ is a spanning subgraph of $\mathcal{G}(T)$. Hence, it follows that $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$. Let $I_1, I_2 \in V(\mathcal{G}(R))$ be such that $I_1 \neq I_2$ and they are adjacent in $\mathcal{G}(R)$. Hence, $I_1 \cap I_2 = I_1 I_2$. Since R is a principal domain, there exist nonzero nonunits $a, b \in R$ such that $I_1 = Ra$ and $I_2 = Rb$. As R is a principal ideal domain, it follows that $Ra + Rb = Rd$ and $Ra \cap Rb = R(\frac{ab}{d})$, where d is the greatest common divisor of a, b in R . From $I_1 \cap I_2 = I_1 I_2$, it follows that $R(\frac{ab}{d}) = Rab$. This implies that $\frac{ab}{d} = rab$ for some $r \in R$ and so, $dr = 1$.

Therefore, $d \in U(R)$. Hence, $Ra + Rb = Rd = R$ and so, I_1 and I_2 are adjacent in $\mathcal{C}(R)$. This shows that $\mathcal{G}(R)$ is a spanning subgraph of $\mathcal{C}(R)$. Therefore, we obtain that $\mathcal{C}(R) = \mathcal{G}(R)$. \square

Let R be a ring such that $|Max(R)| \geq 2$. Suppose that $J(R) \neq (0)$ and $J(R) \in Spec(R)$. We prove in Corollary 2.10 that $diam(\mathcal{G}(R)) = 2$.

Lemma 2.8. *Let R be a ring such that $|Max(R)| \geq 2$. If $\mathcal{G}(R)$ is complete, then $\mathcal{G}(\frac{R}{J(R)})$ is complete.*

Proof. It is convenient to denote $\frac{R}{J(R)}$ by T . Note that $|Max(T)| = |Max(R)| \geq 2$ and $J(T)$ is the zero ideal of T . Observe that $V(\mathcal{G}(T))$ is the set of all nonzero proper ideals of T . Let $A, B \in V(\mathcal{G}(T))$ with $A \neq B$. Note that $A = \frac{I_1}{J(R)}$ and $B = \frac{I_2}{J(R)}$ for some proper ideals I_1, I_2 of R with $I_i \not\subseteq J(R)$ for each $i \in \{1, 2\}$. Now, $I_1, I_2 \in V(\mathcal{G}(R))$ with $I_1 \neq I_2$. By hypothesis, $\mathcal{G}(R)$ is complete. Hence, I_1 and I_2 are adjacent in $\mathcal{G}(R)$. Therefore, $I_1 \cap I_2 = I_1 I_2$. This implies that $\frac{I_1 \cap I_2}{J(R)} = \frac{I_1 I_2}{J(R)}$ and so, $A \cap B = AB$. This shows that A and B are adjacent in $\mathcal{G}(T)$. Therefore, we get that $\mathcal{G}(\frac{R}{J(R)})$ is complete. \square

Corollary 2.9. *Let R be a ring such that $|Max(R)| \geq 2$. If $\mathcal{G}(R)$ is complete, then $\frac{R}{J(R)}$ is von Neumann regular.*

Proof. Assume that $\mathcal{G}(R)$ is complete. Let us denote the ring $\frac{R}{J(R)}$ by T . Note that $|Max(T)| = |Max(R)| \geq 2$. We know from Lemma 2.8 that $\mathcal{G}(T)$ is complete. Since $J(T)$ is the zero ideal of T , we obtain from (i) \Rightarrow (ii) of Theorem 2.4 that $T = \frac{R}{J(R)}$ is von Neumann regular. \square

Corollary 2.10. *Let R be a ring such that $|Max(R)| \geq 2$. If $J(R) \in Spec(R)$, then $diam(\mathcal{G}(R)) = 2$.*

Proof. It is already noted in the proof of Corollary 2.6 that $\mathcal{G}(R)$ has at least one edge. Let us denote $\frac{R}{J(R)}$ by T . Since $J(R) \in Spec(R)$ by assumption, it follows that T is an integral domain and moreover, we know from Proposition 2.2 that $diam(\mathcal{G}(R)) \leq 2$. As $|Max(R)| \geq 2$, it follows that $J(R)$ is not a maximal ideal of R . Hence, T is not a field. It is well-known that an integral domain is von Neumann regular if and only if it is a field. Therefore, T is not von Neumann regular and so, it follows from Corollary 2.9 that $diam(\mathcal{G}(R)) \geq 2$. Hence, we obtain that $diam(\mathcal{G}(R)) = 2$. \square

Recall from [7, page 373] that a ring R is said to be a *Hilbert ring* if each prime ideal of R is an intersection of maximal ideals of R .

Example 2.11. Let $T = K[X, Y]$ be the polynomial ring in two variables X, Y over a field K . Let $I = X^2T$ and let $R = \frac{T}{I}$. Then $\text{diam}(\mathcal{G}(R)) = 2$.

Proof. We know from (1) \Rightarrow (7) of [7, Theorem 31.8] that T is a Hilbert ring and so, we obtain from (1) \Rightarrow (3) of [7, Theorem 31.8] that $R = \frac{T}{I}$ is a Hilbert ring. Therefore, $\text{nil}(R) = J(R)$ and as $\text{nil}(R) = \frac{TX}{I}$, we obtain that $J(R) = \frac{TX}{I} \in \text{Spec}(R)$. It is clear that $|\text{Max}(R)| \geq 2$. Indeed, $\text{Max}(R)$ is infinite. Now, it follows from Corollary 2.10 that $\text{diam}(\mathcal{G}(R)) = 2$. \square

Remark 2.12. Let T, I, R be as in Example 2.11. Since $J(R) \in \text{Spec}(R)$, it follows from Remark 2.3(i) that $\text{diam}(\mathcal{C}(R)) \leq 2$. Let $I_1 = \frac{TX}{I}$ and let $I_2 = \frac{TX^2+TY}{I}$. Note that $I_1 + I_2 \neq R$ and so, I_1 and I_2 are not adjacent in $\mathcal{C}(R)$. Hence, we obtain that $\text{diam}(\mathcal{C}(R)) = 2$. Observe that $I_1 \cap I_2 = R(XY + I) = I_1I_2$. Therefore, I_1 and I_2 are adjacent in $\mathcal{G}(R)$ and so, $\mathcal{C}(R) \neq \mathcal{G}(R)$.

We provide an example in Example 2.15 to illustrate that Corollary 2.10 can fail to hold if the hypothesis that $J(R) \in \text{Spec}(R)$ is omitted.

Lemma 2.13. *Let R be a ring such that $|\text{Max}(R)| \geq 2$. Then $e(\mathfrak{m}) \leq 2$ in $\mathcal{C}(R)$ for each $\mathfrak{m} \in \text{Max}(R)$.*

Proof. We know from [15, Theorem 2.4] that $\mathcal{C}(R)$ is connected and $\text{diam}(\mathcal{C}(R)) \leq 3$. Let $I \in V(\mathcal{C}(R))$ be such that $I \neq \mathfrak{m}$. If $I \not\subseteq \mathfrak{m}$, then $I + \mathfrak{m} = R$ and so, I and \mathfrak{m} are adjacent in $\mathcal{C}(R)$. Suppose that $I \subset \mathfrak{m}$. Since $I \in V(\mathcal{C}(R))$, there exists $\mathfrak{m}' \in \text{Max}(R)$ such that $I \not\subseteq \mathfrak{m}'$. It is clear that $I + \mathfrak{m}' = \mathfrak{m} + \mathfrak{m}' = R$. Hence, $\mathfrak{m} - \mathfrak{m}' - I$ is a path of length 2 between \mathfrak{m} and I in $\mathcal{C}(R)$. This proves that $d(\mathfrak{m}, I) \leq 2$ in $\mathcal{C}(R)$ for any $I \in V(\mathcal{C}(R))$. This shows that $e(\mathfrak{m}) \leq 2$ in $\mathcal{C}(R)$ for any $\mathfrak{m} \in \text{Max}(R)$. \square

Proposition 2.14. *Let R be a ring such that $|\text{Max}(R)| \geq 3$. Then $r(\mathcal{C}(R)) = 2$. If $\text{Max}(R)$ is finite, then $\text{diam}(\mathcal{C}(R)) = 3$.*

Proof. We know from [15, Theorem 2.4] that $\mathcal{C}(R)$ is connected and $\text{diam}(\mathcal{C}(R)) \leq 3$. Let $I \in V(\mathcal{C}(R))$. Then I is a proper ideal of R such that $I \not\subseteq J(R)$. Let \mathfrak{m} be a maximal ideal of R such that $I \subseteq \mathfrak{m}$. We consider the following cases.

Case(i) $I \neq \mathfrak{m}$

As $I + \mathfrak{m} = \mathfrak{m} \neq R$, it follows that I and \mathfrak{m} are not adjacent in $\mathcal{C}(R)$. Hence, $d(I, \mathfrak{m}) \geq 2$ in $\mathcal{C}(R)$. Therefore, $e(I) \geq 2$ in $\mathcal{C}(R)$.

Case(ii) $I = \mathfrak{m}$

Let $\mathfrak{m}' \in \text{Max}(R)$ be such that $\mathfrak{m}' \neq \mathfrak{m}$. Consider the ideal $\mathfrak{m} \cap \mathfrak{m}'$. Since $|\text{Max}(R)| \geq 3$, it is clear that $\mathfrak{m} \cap \mathfrak{m}' \not\subseteq J(R)$. As $\mathfrak{m} + (\mathfrak{m} \cap \mathfrak{m}') = \mathfrak{m} \neq R$, we get that \mathfrak{m} and $\mathfrak{m} \cap \mathfrak{m}'$ are not adjacent in $\mathcal{C}(R)$. Hence, $d(\mathfrak{m}, \mathfrak{m} \cap \mathfrak{m}') \geq 2$ in $\mathcal{C}(R)$. Therefore, $e(\mathfrak{m}) \geq 2$ in $\mathcal{C}(R)$.

This proves that $e(I) \geq 2$ in $\mathcal{C}(R)$ for any $I \in V(\mathcal{C}(R))$. Therefore, $r(\mathcal{C}(R)) \geq 2$.

We know from Lemma 2.13 that $e(\mathfrak{m}) \leq 2$ in $\mathcal{C}(R)$ for each $\mathfrak{m} \in \text{Max}(R)$. Therefore, $e(\mathfrak{m}) = 2$ in $\mathcal{C}(R)$ for each $\mathfrak{m} \in \text{Max}(R)$. This proves that $r(\mathcal{C}(R)) = 2$.

Suppose that $|\text{Max}(R)| \geq 3$ and $\text{Max}(R)$ is finite. Let $|\text{Max}(R)| = n$ and let $\{\mathfrak{m}_1, \mathfrak{m}_2, \mathfrak{m}_3, \dots, \mathfrak{m}_n\}$ denote the set of all maximal ideals of R . Let $I_1 = \mathfrak{m}_1 \cap \mathfrak{m}_2$ and let $I_2 = \mathfrak{m}_1 \cap \mathfrak{m}_3 \cap \dots \cap \mathfrak{m}_n$. Note that $I_1, I_2 \in V(\mathcal{C}(R))$ and $I_1 \neq I_2$. As $I_1 + I_2 \subseteq \mathfrak{m}_1$, it is clear that I_1 and I_2 are not adjacent in $\mathcal{C}(R)$. Observe that $I_1 I_2 \subseteq J(R)$. Now, it follows from Remark 2.3(ii) that $\text{diam}(\mathcal{C}(R)) = 3$. \square

Example 2.15. Let $n \geq 3$ and let $p_1, p_2, p_3, \dots, p_n$ be distinct prime numbers. Let $S = \mathbb{Z} \setminus (\cup_{i=1}^n p_i \mathbb{Z})$. Let $R = S^{-1}\mathbb{Z}$. Then $r(\mathcal{G}(R)) = 2$ and $\text{diam}(\mathcal{G}(R)) = 3$.

Proof. Note that R is a principal ideal domain and $\text{Max}(R) = \{p_1 R, p_2 R, p_3 R, \dots, p_n R\}$. We know from Remark 2.7 that $\mathcal{C}(R) = \mathcal{G}(R)$. Since $|\text{Max}(R)| = n \geq 3$, it follows from Proposition 2.14 that $r(\mathcal{C}(R)) = 2$ and $\text{diam}(\mathcal{C}(R)) = 3$. Therefore, we get that $r(\mathcal{G}(R)) = 2$ and $\text{diam}(\mathcal{G}(R)) = 3$. \square

3. When is $\mathcal{G}(R)$ a finite complete bipartite graph?

Let R be a ring such that $|\text{Max}(R)| \geq 2$. The aim of this section is to classify rings R such that $\mathcal{G}(R)$ is a finite complete bipartite graph.

Lemma 3.1. *Let R be a ring such that $|\text{Max}(R)| \geq 2$. If $\mathcal{G}(R)$ is bipartite, then $|\text{Max}(R)| = 2$.*

Proof. It is already noted in the introduction that $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$. Thus if $\mathcal{G}(R)$ is bipartite, then $\mathcal{C}(R)$ is also a bipartite graph. Hence, we obtain from (2) \Rightarrow (3) of [15, Theorem 4.5] that $|\text{Max}(R)| = 2$. \square

Lemma 3.2. *Let H be a spanning subgraph of a graph $G = (V, E)$. Suppose that H is a complete bipartite graph. If G is a bipartite graph, then $H = G$.*

Proof. Let H be a complete bipartite graph with vertex partition V_1 and V_2 . Let G be a bipartite graph with vertex partition W_1 and W_2 . Note that $V = V_1 \cup V_2 = W_1 \cup W_2$ and $V_1 \cap V_2 = W_1 \cap W_2 = \emptyset$. Let $x \in W_1$. Then either $x \in V_1$ or $x \in V_2$. Without loss of generality, we can assume that $x \in V_1$. Let $x' \in W_1$ be such that $x' \neq x$. If $x' \in V_2$, then x

and x' are adjacent in H and hence, they are adjacent in G . This is impossible since x and x' are not adjacent in G . Therefore, $x' \in V_1$ and this proves that $W_1 \subseteq V_1$. Let $y \in V_2$. Now, $V_2 \subset V = W_1 \cup W_2$. As $W_1 \subseteq V_1$ and $V_1 \cap V_2 = \emptyset$, we obtain that $y \in W_2$. This shows that $V_2 \subseteq W_2$. Let $z \in W_2 \subset V = V_1 \cup V_2$. We claim that $z \in V_2$. Suppose that $z \in V_1$. Let $y \in V_2$. Then z and y are adjacent in H and so, they are adjacent in G . As both z and y are in W_2 , they are not adjacent in G . This is a contradiction and therefore, $z \in V_2$. This shows that $W_2 \subseteq V_2$ and so, $V_2 = W_2$. Hence, we get that $V_1 \subseteq W_1$ and so, $V_1 = W_1$. If $a - b$ is any edge of G , then one of a, b must be in V_1 and the other must be in V_2 . Since H is a complete bipartite graph with vertex partition V_1 and V_2 , we obtain that $a - b$ is an edge of H . This proves that $H = G$. \square

Corollary 3.3. *Let R be a ring such that $|Max(R)| \geq 2$. If $\mathcal{G}(R)$ is a bipartite graph, then $|Max(R)| = 2$, $\mathcal{G}(R)$ is a complete bipartite graph and $\mathcal{C}(R) = \mathcal{G}(R)$.*

Proof. Assume that $\mathcal{G}(R)$ is a bipartite graph. We know from Lemma 3.1 that $|Max(R)| = 2$. Note that $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$. Thus if $\mathcal{G}(R)$ is a bipartite graph, then $\mathcal{C}(R)$ is also a bipartite graph. In such a case, we get from (2) \Rightarrow (1) of [15, Theorem 4.5] that $\mathcal{C}(R)$ is a complete bipartite graph. Therefore, we obtain from Lemma 3.2 that $\mathcal{C}(R) = \mathcal{G}(R)$ and so, $\mathcal{G}(R)$ is a complete bipartite graph. \square

Lemma 3.4. *Let (R_i, \mathfrak{m}_i) be a quasilocal ring for each $i \in \{1, 2\}$ and let $R = R_1 \times R_2$. If $\mathcal{G}(R)$ is a bipartite graph, then R_i is a field for each $i \in \{1, 2\}$.*

Proof. Note that $\{\mathfrak{M}_1 = \mathfrak{m}_1 \times R_2, \mathfrak{M}_2 = R_1 \times \mathfrak{m}_2\}$ is the set of all maximal ideals of R . Assume that $\mathcal{G}(R)$ is a bipartite graph. First, we verify that R_1 is a field. Suppose that R_1 is not a field. Then $\mathfrak{m}_1 \neq (0)$. Note that $\mathfrak{m}_1 \times R_2 - (0) \times R_2 - R_1 \times (0) - \mathfrak{m}_1 \times R_2$ is a cycle of length three in $\mathcal{G}(R)$. This is in contradiction to the assumption that $\mathcal{G}(R)$ is a bipartite graph. Therefore, R_1 is a field. Similarly, it can be shown that R_2 is a field. This shows that R_i is a field for each $i \in \{1, 2\}$. \square

Let R be a ring. Recall from [8] that R is said to *satisfy descending chain condition on principal powers* if for any $a \in R$, the descending sequence of ideals $Ra \supseteq Ra^2 \supseteq Ra^3 \supseteq \dots$ stops after a finite stage.

Let R be a ring such that $|Max(R)| \geq 2$. In Theorem 3.5, we classify such rings R in order that $\mathcal{G}(R)$ to be a finite bipartite graph. Theorem 3.5 of this article is motivated by [15, Proposition 4.7].

Theorem 3.5. *Let R be a ring such that $|Max(R)| \geq 2$. The following statements are equivalent:*

- (i) $\mathcal{G}(R)$ is a finite bipartite graph.
- (ii) $\mathcal{G}(R)$ is a bipartite graph and R satisfies d.c.c. on principal powers on elements $a \in R$ such that $Ra \in V(\mathcal{G}(R))$.
- (iii) $R \cong F_1 \times F_2$ as rings, where F_i is a field for each $i \in \{1, 2\}$.

Proof. (i) \Rightarrow (ii) As $\mathcal{G}(R)$ is a finite bipartite graph, it follows that $V(\mathcal{G}(R))$ is finite. Let $a \in R$ be such that $Ra \in V(\mathcal{G}(R))$. Then $a \in NU(R) \setminus J(R)$. Then $a^n \in NU(R) \setminus J(R)$ for all $n \geq 1$ and so, $Ra^n \in V(\mathcal{G}(R))$ for all $n \geq 1$. Since $V(\mathcal{G}(R))$ is finite, we obtain that there exists $n \geq 1$ such that $Ra^n = Ra^j$ for all $j \geq n$.

(ii) \Rightarrow (iii) Now, by assumption, $\mathcal{G}(R)$ is a bipartite graph. We know from Corollary 3.3 that $|Max(R)| = 2$ and $\mathcal{C}(R) = \mathcal{G}(R)$ is a complete bipartite graph. Let $\{\mathfrak{M}_1, \mathfrak{M}_2\}$ denote the set of all maximal ideals of R . Let $a \in \mathfrak{M}_1 \setminus \mathfrak{M}_2$. Observe that $Ra \in V(\mathcal{G}(R))$. As we are assuming that R satisfies d.c.c. on principal powers on elements $x \in R$ such that $Rx \in V(\mathcal{G}(R))$, we get that there exists $n \geq 1$ such that $Ra^n = Ra^j$ for all $j \geq n$. This implies that $a^n = ra^{2n}$ for some $r \in R$. Therefore, $e = ra^n$ is a nontrivial idempotent element of R . Hence, the mapping $f : R \rightarrow Re \times R(1 - e)$ defined by $f(x) = (xe, x(1 - e))$ is an isomorphism of rings. Let us denote the ring Re by R_1 and the ring $R(1 - e)$ by R_2 . Now, $R \cong R_1 \times R_2$ as rings. Since $|Max(R)| = 2$, it follows that R_i is a quasilocal ring for each $i \in \{1, 2\}$. As $\mathcal{G}(R_1 \times R_2)$ is a bipartite graph, we obtain from Lemma 3.4 that R_i is a field for each $i \in \{1, 2\}$. Let $i \in \{1, 2\}$. With $F_i = R_i$, we obtain that F_i is a field and $R \cong F_1 \times F_2$ as rings.

(iii) \Rightarrow (i) Let us denote the ring $F_1 \times F_2$ by T , where F_i is a field for each $i \in \{1, 2\}$. Note that $V(\mathcal{G}(T)) = \{(0) \times F_2, F_1 \times (0)\}$. From $((0) \times F_2) + (F_1 \times (0)) = T$, we get that $\mathcal{G}(T)$ is a complete graph on two vertices. Since $R \cong T$ as rings, we obtain that $\mathcal{G}(R)$ is a complete graph on two vertices and hence, it is a finite bipartite graph. \square

In Example 3.6, we mention an example of a ring R which illustrates that (ii) \Rightarrow (iii) of Theorem 3.5 can fail to hold if the hypothesis that R satisfies d.c.c. on principal powers on elements $a \in R$ with $a \in V(\mathcal{G}(R))$ is omitted. The example 3.6 mentioned here is [15, Example 4.10].

Example 3.6. Let p, q be distinct prime numbers. Let $S = \mathbb{Z} \setminus (p\mathbb{Z} \cup q\mathbb{Z})$. Let $R = S^{-1}\mathbb{Z}$. Then $\mathcal{G}(R)$ is a complete bipartite graph and R has no nontrivial idempotent element.

Proof. Note that R is a semilocal principal ideal domain with $Max(R) = \{pR, qR\}$. We know from Remark 2.7 that $\mathcal{C}(R) = \mathcal{G}(R)$. Since $|Max(R)| = 2$, we know from [15, Lemma 4.1]

that $\mathcal{C}(R)$ is a complete bipartite graph. This shows that $\mathcal{G}(R)$ is a complete bipartite graph. As R is an integral domain, we obtain that 0 and 1 are the only idempotent elements of R . Hence, $R \not\cong R_1 \times R_2$ as rings for any quasilocal rings R_1 and R_2 . \square

Let I be an ideal of a ring R . As in [12], we denote $\{\mathfrak{m} \in \text{Max}(R) \mid \mathfrak{m} \supseteq I\}$ by $M(I)$.

Remark 3.7. Let R be a ring with $|\text{Max}(R)| = 2$. Let $\{\mathfrak{m}_1, \mathfrak{m}_2\}$ denote the set of all maximal ideals of R . We know from [15, Lemma 4.1] that $\mathcal{C}(R)$ is a complete bipartite graph with vertex partition V_1 and V_2 , where V_1 is the set of all ideals I of R such that $M(I) = \{\mathfrak{m}_1\}$ and V_2 is the set of all ideals J of R such that $M(J) = \{\mathfrak{m}_2\}$. It is noted in [15, Corollary 4.2] that $1 \leq \text{diam}(\mathcal{C}(R)) \leq 2$. It is already observed in the introduction that $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$. Therefore, we obtain that $1 \leq \text{diam}(\mathcal{G}(R)) \leq 2$. In Corollary 3.9, we classify rings R such that $\text{diam}(\mathcal{G}(R)) = 1$, that is, we classify rings R such that $\mathcal{G}(R)$ is complete.

Theorem 3.8. *Let (R_i, \mathfrak{m}_i) be a quasilocal ring for each $i \in \{1, 2\}$ and let $R = R_1 \times R_2$. The following statements are equivalent:*

- (i) $\mathcal{G}(R)$ is complete.
- (ii) For each $i \in \{1, 2\}$, \mathfrak{m}_i is principal and $\mathfrak{m}_i^2 = (0)$.

Proof. (i) \Rightarrow (ii) Assume that $\mathcal{G}(R)$ is complete. First, we verify that \mathfrak{m}_1 is principal and $\mathfrak{m}_1^2 = (0)$. This is clear if $\mathfrak{m}_1 = (0)$. Suppose that $\mathfrak{m}_1 \neq (0)$. Let $a \in \mathfrak{m}_1 \setminus \{0\}$. We assert that $\mathfrak{m}_1 = R_1 a$. If $\mathfrak{m}_1 \neq R_1 a$, then the ideals $I_1 = R_1 a \times R_2$ and $I_2 = \mathfrak{m}_1 \times R_2$ are distinct members of $V(\mathcal{G}(R))$. Since we are assuming that $\mathcal{G}(R)$ is complete, we get that $I_1 \cap I_2 = I_1 I_2$. This implies that $R_1 a = \mathfrak{m}_1 a$ and so, $a = xa$ for some $x \in \mathfrak{m}_1$. Hence, $a(1 - x) = 0$. Since $1 - x \in U(R_1)$, we get that $a = 0$. This is in contradiction to the fact that $a \neq 0$. This proves that $\mathfrak{m}_1 = R_1 a$ for any $a \in \mathfrak{m}_1 \setminus \{0\}$. Let $a \in \mathfrak{m}_1 \setminus \{0\}$. If $a^2 \neq 0$, then $R_1 a = R_1 a^2$. This implies that $a = xa^2$ for some $x \in R_1$ and so, $a(1 - xa) = 0$. Since $1 - xa \in U(R_1)$, we obtain that $a = 0$. This is a contradiction. Therefore, \mathfrak{m}_1 is principal and $\mathfrak{m}_1^2 = (0)$. Similarly, it can be shown that \mathfrak{m}_2 is principal and $\mathfrak{m}_2^2 = (0)$.

(ii) \Rightarrow (i) Assume that \mathfrak{m}_i is principal and $\mathfrak{m}_i^2 = (0)$ for each $i \in \{1, 2\}$. Note that $|\text{Max}(R)| = 2$ and $\{\mathfrak{M}_1 = \mathfrak{m}_1 \times R_2, \mathfrak{M}_2 = R_1 \times \mathfrak{m}_2\}$ is the set of all maximal ideals of R . We consider the following cases.

Case(1) $\mathfrak{m}_i = (0)$ for each $i \in \{1, 2\}$

In this case, both R_1 and R_2 are fields. Note that $V(\mathcal{G}(R)) = \{(0) \times R_2, R_1 \times (0)\}$ and $((0) \times R_2) + (R_1 \times (0)) = R$. Hence, $(0) \times R_2$ and $R_1 \times (0)$ are adjacent in $\mathcal{C}(R)$ and so, they are adjacent in $\mathcal{G}(R)$. Therefore, $\mathcal{G}(R)$ is a complete graph on two vertices.

Case(2) $\mathfrak{m}_1 \neq (0)$ but $\mathfrak{m}_2 = (0)$

Note that R_2 is a field. As \mathfrak{m}_1 is a nonzero principal ideal of R_1 with $\mathfrak{m}_1^2 = (0)$, it follows that \mathfrak{m}_1 is the only nonzero proper ideal of R_1 . Observe that $V(\mathcal{G}(R)) = \{(0) \times R_2, \mathfrak{m}_1 \times R_2, R_1 \times (0)\}$. Observe that $\mathcal{C}(R)$ is a complete bipartite graph with vertex partition $V_1 = \{\mathfrak{M}_1^2 = (0) \times R_2, \mathfrak{M}_1 = \mathfrak{m}_1 \times R_2\}$ and $V_2 = \{\mathfrak{M}_2 = R_1 \times (0)\}$. As $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$, it follows that each member of V_1 is adjacent to each member of V_2 in $\mathcal{G}(R)$. Observe that $\mathfrak{M}_1 \cap \mathfrak{M}_1^2 = \mathfrak{M}_1^2 = \mathfrak{M}_1^3$. Therefore, there is an edge of $\mathcal{G}(R)$ joining \mathfrak{M}_1 and \mathfrak{M}_1^2 . This proves that $\mathcal{G}(R)$ is the cycle Γ of length three given by $\Gamma : \mathfrak{M}_1 - \mathfrak{M}_2 - \mathfrak{M}_1^2 - \mathfrak{M}_1$ and so, $\mathcal{G}(R)$ is a complete graph on three vertices.

Case(3) $\mathfrak{m}_1 = (0)$ but $\mathfrak{m}_2 \neq (0)$

Since \mathfrak{m}_2 is a nonzero principal ideal of R_2 with $\mathfrak{m}_2^2 = (0)$, it follows as in Case(2) that $\mathcal{G}(R)$ is a complete graph on three vertices.

Case(4) $\mathfrak{m}_i \neq (0)$ for each $i \in \{1, 2\}$

Let $i \in \{1, 2\}$. Since \mathfrak{m}_i is a nonzero principal ideal of R_i with $\mathfrak{m}_i^2 = (0)$, it follows that \mathfrak{m}_i is the only nonzero proper ideal of R_i . Note that $V(\mathcal{G}(R)) = \{\mathfrak{M}_1 = \mathfrak{m}_1 \times R_2, \mathfrak{M}_1^2 = (0) \times R_2, \mathfrak{M}_2 = R_1 \times \mathfrak{m}_2, \mathfrak{M}_2^2 = R_1 \times (0)\}$. Observe that $\mathcal{C}(R)$ is a complete bipartite graph with vertex partition $V_1 = \{\mathfrak{M}_1, \mathfrak{M}_1^2\}$ and $V_2 = \{\mathfrak{M}_2, \mathfrak{M}_2^2\}$. Since $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$, it follows that each member of V_1 is adjacent to each member of V_2 in $\mathcal{G}(R)$. Moreover, it follows as in Case(2) that $\mathfrak{M}_i - \mathfrak{M}_i^2$ is an edge of $\mathcal{G}(R)$ for each $i \in \{1, 2\}$. Therefore, we get that $\mathcal{G}(R)$ is a complete graph on four vertices.

This proves that $\mathcal{G}(R)$ is complete. \square

Corollary 3.9. *Let R be a ring such that $|Max(R)| = 2$. The following statements are equivalent:*

- (i) $\mathcal{G}(R)$ is complete.
- (ii) R is isomorphic to one of the following rings:
 - (a) $F_1 \times F_2$, where F_i is a field for each $i \in \{1, 2\}$.
 - (b) $F_1 \times R_2$, where F_1 is a field and (R_2, \mathfrak{m}_2) is a SPIR with $\mathfrak{m}_2 \neq (0)$ but $\mathfrak{m}_2^2 = (0)$.
 - (c) $R_1 \times R_2$, where (R_i, \mathfrak{m}_i) is a SPIR with $\mathfrak{m}_i \neq (0)$ but $\mathfrak{m}_i^2 = (0)$ for each $i \in \{1, 2\}$.

Proof. (i) \Rightarrow (ii) Let $\{\mathfrak{M}_1, \mathfrak{M}_2\}$ denote the set of all maximal ideals of R . We assert that R admits at least one nontrivial idempotent. Let $a \in \mathfrak{M}_1 \setminus \mathfrak{M}_2$. Then for all $n \geq 2$, $a^n \in \mathfrak{M}_1 \setminus \mathfrak{M}_2$. Hence, $Ra^n \in V(\mathcal{G}(R))$ for all $n \geq 1$. If $Ra = Ra^2$, then $a = ra^2$ for some $r \in R$. In such a case, ra is a nontrivial idempotent element of R . Suppose that $Ra \neq Ra^2$. Since $\mathcal{G}(R)$ is complete, the vertices Ra and Ra^2 are adjacent in $\mathcal{G}(R)$. Hence, $Ra \cap Ra^2 = Ra^3$ and so, $Ra^2 = Ra^3$. Therefore, $Ra^2 = Ra^4$. This implies that $a^2 = sa^4$ for some $s \in R$ and so, sa^2 is a nontrivial idempotent element of R . This shows that there exists a nontrivial idempotent

element e of R . Note that the mapping $f : R \rightarrow Re \times R(1 - e)$ defined by $f(r) = (re, r(1 - e))$ is an isomorphism of rings. Let us denote the ring Re by R_1 and the ring $R(1 - e)$ by R_2 . Let us denote the ring $R_1 \times R_2$ by T . Since $R \cong T$ as rings, we obtain that $\mathcal{G}(T)$ is complete. As $|Max(T)| = 2$, it follows that R_i admits a unique maximal ideal for each $i \in \{1, 2\}$. Let \mathfrak{m}_i denote the unique maximal ideal of R_i for each $i \in \{1, 2\}$. Now, we know from (i) \Rightarrow (ii) of Theorem 3.8 that \mathfrak{m}_i is principal and $\mathfrak{m}_i^2 = (0)$ for each $i \in \{1, 2\}$. If $\mathfrak{m}_i = (0)$ for each $i \in \{1, 2\}$, then R_i is a field for each $i \in \{1, 2\}$. With $F_i = R_i$ for each $i \in \{1, 2\}$, we get that R is isomorphic to the ring mentioned in (ii)(a). Suppose that exactly one between \mathfrak{m}_1 and \mathfrak{m}_2 is the zero ideal. Without loss of generality, we can assume that $\mathfrak{m}_1 = (0)$. Then R_1 is a field and (R_2, \mathfrak{m}_2) is a SPIR with $\mathfrak{m}_2 \neq (0)$ but $\mathfrak{m}_2^2 = (0)$. In this case, with $F_1 = R_1$, we obtain that R is isomorphic to the ring mentioned in (ii)(b). If $\mathfrak{m}_i \neq (0)$ for each $i \in \{1, 2\}$, then (R_i, \mathfrak{m}_i) is a SPIR with $\mathfrak{m}_i \neq (0)$ but $\mathfrak{m}_i^2 = (0)$ for each $i \in \{1, 2\}$. and R is isomorphic to the ring mentioned in (ii)(c).

(ii) \Rightarrow (i) Assume that R is isomorphic to one of the rings mentioned in (ii)(a), (b) or (c). Let F_1, F_2 be fields. Let us denote $F_1 \times F_2$ by T_1 . We know from the proof of (ii) \Rightarrow (i) Case(1) of Theorem 3.8 that $\mathcal{G}(T_1)$ is a complete graph on two vertices. Let F_1 be a field and (R_2, \mathfrak{m}_2) be a SPIR with $\mathfrak{m}_2 \neq (0)$ but $\mathfrak{m}_2^2 = (0)$. Let us denote the ring $F_1 \times R_2$ by T_2 . We know from the proof of (ii) \Rightarrow (i) Case(3) of Theorem 3.8 that $\mathcal{G}(T_2)$ is a complete graph on three vertices. Suppose that (R_i, \mathfrak{m}_i) be a SPIR with $\mathfrak{m}_i \neq (0)$ but $\mathfrak{m}_i^2 = (0)$ for each $i \in \{1, 2\}$. Let us denote the ring $R_1 \times R_2$ by T_3 . We know from the proof of (ii) \Rightarrow (i) Case(4) of Theorem 3.8 that $\mathcal{G}(T_3)$ is a complete graph on four vertices. This proves that $\mathcal{G}(R)$ is complete. \square

4. On the clique number and the chromatic number of $\mathcal{G}(R)$

Let R be a ring with $|Max(R)| \geq 2$. The aim of this section is to discuss some results regarding $\omega(\mathcal{G}(R))$ and $\chi(\mathcal{G}(R))$.

Remark 4.1. Let R be a ring with $|Max(R)| \geq 2$. Suppose that $\omega(\mathcal{G}(R)) < \infty$. It is already noted in the introduction that $\mathcal{C}(R)$ is a spanning subgraph of $\mathcal{G}(R)$. Hence, $\omega(\mathcal{C}(R))$ is also finite. If $\omega(\mathcal{C}(R)) = n$, then we know from [15, Theorem 3.1] that $|Max(R)| = n = \chi(\mathcal{C}(R))$. Thus if $\omega(\mathcal{G}(R)) < \infty$, then R is semiquasilocal. In Example 4.2, we provide an example of a semiquasilocal ring R such that $\mathcal{G}(R)$ admits an infinite clique.

Example 4.2. Let V be an infinite dimensional vector space over a field K . Let $T = K \oplus V$ be the ring obtained on using Nagata's principle of idealization. Let $R = T \times T$. Then $|Max(R)| = 2$ and $\mathcal{G}(R)$ admits an infinite clique.

Proof. Note that T is quasilocal with $\mathfrak{m} = (0) \oplus V$ as its unique maximal ideal and $\mathfrak{m}^2 = (0) \oplus (0)$. Hence, $R = T \times T$ has $\{\mathfrak{M}_1 = \mathfrak{m} \times T, \mathfrak{M}_2 = T \times \mathfrak{m}\}$ as its set of all maximal ideals. This shows that $|Max(R)| = 2$. Since V is an infinite dimensional vector space over K , it is possible to find $v_i \in V$ for each $i \in \mathbb{N}$ such that $\{v_i | i \in \mathbb{N}\}$ is linearly independent over K . For each $i \in \mathbb{N}$, let us denote the ideal $(0) \oplus Kv_i$ of T by I_i . Let $i, j \in \mathbb{N}, i \neq j$. Since v_i, v_j are linearly independent over K , it follows that $Kv_i \cap Kv_j = (0)$ and so, $I_i \cap I_j = (0) \oplus (0)$. Hence, $I_i I_j = I_i \cap I_j = (0) \oplus (0)$. For each $i \in \mathbb{N}$, let us denote the ideal $I_i \times T$ of R by A_i . It is clear that A_i is a proper ideal of R and $A_i \not\subseteq J(R)$ for each $i \in \mathbb{N}$. Hence, $A_i \in V(\mathcal{G}(R))$ for each $i \in \mathbb{N}$. Let i, j be distinct elements of \mathbb{N} . From $I_i I_j = I_i \cap I_j$, it follows that $A_i A_j = I_i I_j \times T = (I_i \cap I_j) \times T = A_i \cap A_j$. Hence, the subgraph of $\mathcal{G}(R)$ induced on $\{A_i | i \in \mathbb{N}\}$ is an infinite clique. \square

In Proposition 4.3 we classify rings R with $|Max(R)| \geq 2$ and $J(R) = (0)$ such that $\omega(\mathcal{G}(R)) < \infty$.

Proposition 4.3. *Let R be a ring such that $|Max(R)| \geq 2$ and suppose that $J(R) = (0)$. The following statements are equivalent:*

- (i) $\omega(\mathcal{G}(R)) < \infty$.
- (ii) $\mathcal{G}(R)$ does not contain any infinite clique.
- (iii) There exist $n \in \mathbb{N}$ with $n \geq 2$ and fields F_1, F_2, \dots, F_n such that $R \cong F_1 \times F_2 \times \dots \times F_n$ as rings.

Moreover, if any one of the statements (i), (ii) or (iii) holds (and hence, all the three hold), then $\omega(\mathcal{G}(R)) = \chi(\mathcal{G}(R)) = 2^n - 2$.

Proof. (i) \Rightarrow (ii) This is clear.

(ii) \Rightarrow (iii) We claim that $Max(R)$ is finite. Suppose that $Max(R)$ is infinite. If \mathfrak{m} is any element of $Max(R)$, then $\mathfrak{m} \in V(\mathcal{G}(R))$. Now, for any distinct $\mathfrak{m}, \mathfrak{m}' \in Max(R)$, $\mathfrak{m} + \mathfrak{m}' = R$. Hence, \mathfrak{m} and \mathfrak{m}' are adjacent in $\mathcal{C}(R)$ and so, they are adjacent in $\mathcal{G}(R)$. Therefore, the subgraph of $\mathcal{G}(R)$ induced on $Max(R)$ is an infinite clique. This is in contradiction to the assumption that $\mathcal{G}(R)$ does not contain any infinite clique. Hence, $Max(R)$ is finite. Let $|Max(R)| = n$. It is clear that $n \geq 2$. Let $\{\mathfrak{m}_1, \mathfrak{m}_2, \dots, \mathfrak{m}_n\}$ denote the set of all maximal ideals of R . By hypothesis, $J(R) = (0)$. Hence, $\cap_{i=1}^n \mathfrak{m}_i = (0)$. Now, it follows from the Chinese remainder theorem [2, Proposition 1.10(ii) and (iii)] that the mapping $f : R \rightarrow \frac{R}{\mathfrak{m}_1} \times \frac{R}{\mathfrak{m}_2} \times \dots \times \frac{R}{\mathfrak{m}_n}$ given by $f(r) = (r + \mathfrak{m}_1, r + \mathfrak{m}_2, \dots, r + \mathfrak{m}_n)$ is an isomorphism of rings. For each $i \in \{1, 2, \dots, n\}$, let us denote the field $\frac{R}{\mathfrak{m}_i}$ by F_i . It is clear that $R \cong F_1 \times F_2 \times \dots \times F_n$ as rings.

(iii) \Rightarrow (i) Let us denote the ring $F_1 \times F_2 \times \cdots \times F_n$ by T . Note that T is a von Neumann regular ring. Hence, we obtain from (ii) \Rightarrow (i) of Theorem 2.4 that $\mathcal{G}(T)$ is complete. Since $V(\mathcal{G}(T))$ is the set of all nonzero proper ideals of T , it follows that $\mathcal{G}(T)$ is a complete graph on $2^n - 2$ vertices. From $R \cong T$ as rings, we get that $\mathcal{G}(R)$ is a complete graph on $2^n - 2$ vertices and so, $\omega(\mathcal{G}(R)) = \chi(\mathcal{G}(R)) = 2^n - 2$.

The moreover part of this proposition is already verified in (iii) \Rightarrow (i) of this proposition. \square

In Example 4.4, we provide an example of a ring R such that $|Max(R)| = 2$ with $\omega(\mathcal{G}(R)) = 5 < \chi(\mathcal{G}(R)) = 6$.

Example 4.4. Let $T = \mathbb{Z}_4[X, Y, Z]$ be the polynomial ring in three variables X, Y, Z over \mathbb{Z}_4 . Let I be the ideal of T generated by $\{X^2 - 2, Y^2 - 2, Z^2, XY, YZ - 2, XZ, 2X, 2Y, 2Z\}$. Let $S = \frac{T}{I}$ and let $R = S \times F$, where F is a field. Then $|Max(R)| = 2$ and $\omega(\mathcal{G}(R)) = 5 < \chi(\mathcal{G}(R)) = 6$.

Proof. The ring S mentioned in Example 4.4 is an interesting and inspiring example due to D.D. Anderson and M. Nasser [1] which answered a conjecture of I. Beck [4] in the negative. It was already noted in [1] that S is a finite local ring with $\mathfrak{m} = \frac{TX+TY+TZ}{I}$ as its unique maximal ideal, $\mathfrak{m}^2 = S(2 + I) = \{0 + I, 2 + I\}$, and $\mathfrak{m}^3 = (0 + I)$. Since $R = S \times F$, it is clear that $|Max(R)| = 2$ and $\{\mathfrak{M}_1 = \mathfrak{m} \times F, \mathfrak{M}_2 = S \times (0)\}$ is the set of all maximal ideals of R . It is convenient to denote $X + I$ by x , $Y + I$ by y , and $Z + I$ by z . It was already observed in the proof of [6, Proposition 2.1] that the set of all nonzero proper ideals of S equals $\{S(2 + I), Sx, Sy, Sz, S(x + y), S(y + z), S(x + z), S(x + y + z), Sx + Sy, Sy + Sz, Sx + Sz, Sx + S(y + z), Sy + S(z + x), Sz + S(x + y), Sx + Sy + Sz\}$. Observe that $\mathfrak{m}^2 = S(2 + I)$ is the unique minimal ideal of S . Note that $J(R) = \mathfrak{m} \times (0)$ and $V(\mathcal{G}(R))$ equals $\{\mathfrak{m} \times F, (0 + I) \times F, S \times (0), Sx \times F, Sy \times F, Sz \times F, S(x + y) \times F, S(y + z) \times F, S(x + z) \times F, S(x + y + z) \times F, (Sx + Sy) \times F, (Sy + Sz) \times F, (Sx + Sz) \times F, (Sx + S(y + z)) \times F, (Sy + S(x + z)) \times F, (Sz + S(x + y)) \times F, S(2 + I) \times F\}$.

We next proceed to verify that $\omega(\mathcal{G}(R)) \leq 5$. It is convenient to denote $\{Sx \times F, Sy \times F, Sz \times F\}$ by W_1 , $\{S(x + y) \times F, S(y + z) \times F, S(x + z) \times F, (Sx + Sy) \times F, (Sy + Sz) \times F, (Sx + Sz) \times F\}$ by W_2 , and $\{(Sx + S(y + z)) \times F, (Sy + S(x + z)) \times F, (Sz + S(x + y)) \times F\}$ by W_3 . Let $W \subseteq V(\mathcal{G}(R))$ be such that the subgraph of $\mathcal{G}(R)$ induced on W is a clique.

Suppose that $\mathfrak{m}^2 \times F = S(2 + I) \times F \in W$. As \mathfrak{m}^2 is the unique minimal ideal of S , it follows that for any nonzero proper ideal B of S , $\mathfrak{m}^2 \cap B = \mathfrak{m}^2$ but $\mathfrak{m}^2 B = (0 + I)$. Hence, $S(2 + I) \times F$ is not adjacent in $\mathcal{G}(R)$ to any vertex of the form $B \times F$, where B is a nonzero proper ideal of S . Hence, the only possible vertices that can belong to W are $\mathfrak{m}^2 \times F$, $(0 + I) \times F$, and $S \times (0)$. Thus if $S(2 + I) \times F \in W$, then $|W| \leq 3$.

Hereafter, we assume that $\mathfrak{m}^2 \times F \notin W$.

Suppose that $\mathfrak{m} \times F \in W$. Let B_1, B_2 be distinct nonzero proper ideals of S such that $B_1 \cap B_2 \not\subseteq \mathfrak{m}^2$. Note that $B_i \not\subseteq \mathfrak{m}^2$ for each $i \in \{1, 2\}$. Then $(B_1 \times F) \cap (B_2 \times F) = (B_1 \cap B_2) \times F \not\subseteq \mathfrak{m}^2 \times F$, whereas $(B_1 \times F)(B_2 \times F) = B_1 B_2 \times F \subseteq \mathfrak{m}^2 \times F$. Hence, $B_1 \times F$ and $B_2 \times F$ are not adjacent in $\mathcal{G}(R)$. Therefore, at most one between $B_1 \times F$ and $B_2 \times F$ can be in W . In particular, if C_1 and C_2 are nonzero proper ideals of S with $C_i \not\subseteq \mathfrak{m}^2$ for each $i \in \{1, 2\}$ and $C_1 \subset C_2$, then at most one between $C_1 \times F$ and $C_2 \times F$ can be in W . Thus if $\mathfrak{m} \times F \in W$, then the possible vertices that can be in W are $\mathfrak{m} \times F, S \times (0)$, and $(0) \times F$. Therefore, $|W| \leq 3$.

Hereafter, we assume that $\mathfrak{m} \times F \notin W$.

Since S is a local ring, it follows that if $m_1, m_2 \in \mathfrak{m} \setminus \mathfrak{m}^2$ are such that $Sm_1 \neq Sm_2$, then $Sm_1 \cap Sm_2 \subseteq \mathfrak{m}^2$. Hence, if $m_1 m_2 \neq 0 + I$, then $Sm_1 m_2 = Sm_1 \cap Sm_2$. Therefore, $Sm_1 \times F$ and $Sm_2 \times F$ are adjacent in $\mathcal{G}(R)$. If $m_1 m_2 = (0) + I$, then as $\mathfrak{m}^2 \times F \subseteq (Sm_1 \times F) \cap (Sm_2 \times F)$, it follows that $Sm_1 \times F$ and $Sm_2 \times F$ are not adjacent in $\mathcal{G}(R)$. Since $Sy \neq Sz$ and $yz = 2 + I \neq 0 + I$, we get that $Sy \times F$ and $Sz \times F$ are adjacent in $\mathcal{G}(R)$. Observe that $\mathfrak{m}^2 \times F \subseteq (Sx \times F) \cap (Sy \times F) \cap (Sz \times F)$, whereas $Sxy \times F = (0 + I) \times F = Sxz \times F$. Hence, we get that $(Sx \times F) \cap (Sy \times F) \neq Sxy \times F$ and $(Sx \times F) \cap (Sz \times F) \neq Sxz \times F$ and so, in $\mathcal{G}(R)$, $Sx \times F$ is not adjacent to any of the members from $\{Sy \times F, Sz \times F\}$. Thus if either $Sy \times F \in W$ or $Sz \times F \in W$, then $Sx \times F$ cannot be in W .

As $(x + y)(y + z) = 0 + I$, at most one between $S(x + y) \times F$ and $S(y + z) \times F$ can be in W . Hence, $|W \cap \{S(x + y) \times F, S(y + z) \times F, S(x + z) \times F\}| \leq 2$. It is clear that at most one among $(Sx + Sy) \times F, (Sy + Sz) \times F$, and $(Sx + Sz) \times F$ can belong to W . Note that $(x + y)(x + z) = 0 + I$ and so, $S(x + y) \times F$ and $S(x + z) \times F$ are not adjacent in $\mathcal{G}(R)$. It follows from $(y + z)(x + z) = 2 + I \neq 0 + I$ that $S(y + z) \times F$ and $S(x + z) \times F$ are adjacent in $\mathcal{G}(R)$. Suppose that both $S(y + z) \times F$ and $S(x + z) \times F$ are in W . Then no member from $\{(Sx + Sy) \times F, (Sy + Sz) \times F, (Sx + Sz) \times F\}$ can belong to W . This shows that $|W \cap W_2| \leq 2$.

Note that $x + y + z$ is in each member of W_3 . Hence, at most one element of W_3 can belong to W . Moreover, if $S(x + y + z) \times F \in W$, then $W \cap W_3 = \emptyset$.

Suppose that both $Sy \times F$ and $Sz \times F$ are in W . Note that $y(y + z) = y^2 + yz = (2 + I) + (2 + I) = 0 + I$. Observe that $0 + I = xy = xz = y(y + z) = y(x + y + z) = z(x + z)$. Also, as $Sy \times F \subset (Sx + Sy) \times F, Sy \times F \subset (Sy + Sz) \times F$, and $Sz \times F \subset (Sx + Sz) \times F$, it follows that the only possible member of W_2 that can belong to W is $S(x + y) \times F$. Moreover, it is clear that $Sx \times F$ and $S(x + y + z) \times F$ cannot be in W and furthermore, $W \cap W_3 = \emptyset$. From the above given arguments, it is clear that the only possible members of $V(\mathcal{G}(R))$ that can be in W are $Sy \times F, Sz \times F, S(x + y) \times F, (0) \times F$, and $S \times (0)$. Hence, $|W| \leq 5$.

Hereafter, we assume that at most one between $Sy \times F$ and $Sz \times F$ is in W .

Suppose that $Sy \times F \in W$ but $Sz \times F \notin W$.

It follows as in the previous paragraph that none of the members from $\{Sx \times F, S(y+z) \times F, S(x+y+z) \times F, (Sx+Sy) \times F, (Sy+Sz) \times F, (Sx+S(y+z)) \times F, (Sy+S(x+z)) \times F\}$ can be in W . Note that $y(x+y) = y(x+z) = 2+I \neq 0+I$ and $(x+y)(x+z) = 0+I$. It is clear that at most two members from $\{S(x+y) \times F, S(x+z) \times F, (Sx+Sz) \times F, (Sx+S(y+z)) \times F, (Sy+S(x+z)) \times F\}$ can be in W . From the above discussion, it follows that $W \subseteq \{Sy \times F\} \cup (W \cap \{S(x+y) \times F, S(x+z) \times F, (Sx+Sz) \times F, (Sx+S(y+z)) \times F, (Sy+S(x+z)) \times F\}) \cup \{(0) \times F, S \times (0)\}$. Therefore, we obtain that $|W| \leq 5$.

Suppose that $Sz \times F \in W$ but $Sy \times F \notin W$.

It follows from the reasons mentioned earlier in the verification of this example that none of the members from $\{Sx \times F, S(x+z) \times F, (Sy+Sz) \times F, (Sx+Sz) \times F, (Sx+S(y+z)) \times F\}$ can be in W . Observe that $z(x+y) = z(y+z) = 2+I \neq 0+I$. Hence, we obtain that the possible members from W_2 that can belong to W are $S(x+y) \times F, (Sx+Sy) \times F, S(y+z) \times F$ and so, we obtain that $|W \cap W_2| \leq 1$. Note that $z(x+y+z) = 2+I \neq 0+I$. Hence, $Sz \times F$ and $S(x+y+z) \times F$ are adjacent in $\mathcal{G}(R)$. If $S(x+y+z) \times F \in W$, then it is already observed that $W \cap W_3 = \emptyset$. Thus in this case, $W \subseteq \{Sz \times F, S(x+y+z) \times F\} \cup (W \cap W_2) \cup \{(0) \times F, S \times (0)\}$ and so, $|W| \leq 5$. Suppose that $S(x+y+z) \times F \notin W$. It is already noted that $|W \cap W_3| \leq 1$. In this case, $W \subseteq \{Sz \times F\} \cup (W \cap W_2) \cup (W \cap W_3) \cup \{(0) \times F, S \times (0)\}$. Therefore, we obtain that $|W| \leq 5$.

Hereafter, we assume that both $Sy \times F$ and $Sz \times F$ are not in W .

Suppose that $Sx \times F \in W$. Then none of the members from $\{S(y+z) \times F, (Sx+Sy) \times F, (Sx+Sz) \times F, (Sy+Sz) \times F, (Sx+S(y+z)) \times F\}$ can belong to W . It is easy to verify that the possible members from W_2 that can be in W are $S(x+y) \times F$ and $S(x+z) \times F$. Since $S(x+y) \times F$ and $S(x+z) \times F$ are not adjacent in $\mathcal{G}(R)$, we obtain that $|W \cap W_2| \leq 1$. From $x(x+y+z) \neq 0+I$, it follows that $Sx \times F$ and $S(x+y+z) \times F$ are adjacent in $\mathcal{G}(R)$. If $S(x+y+z) \times F \in W$, then $W \cap W_3 = \emptyset$. Therefore, in this case, $W \subseteq \{Sx \times F, S(x+y+z) \times F\} \cup (W \cap W_2) \cup \{(0) \times F, S \times (0)\}$ and so, $|W| \leq 5$. Suppose that $S(x+y+z) \times F \notin W$. It is already noted that $|W \cap W_3| \leq 1$. Hence, in this case, we get that $W \subseteq \{Sx \times F\} \cup (W \cap W_2) \cup (W \cap W_3) \cup \{(0) \times F, S \times (0)\}$ and so, $|W| \leq 5$.

Hereafter, we assume that $Sx \times F \notin W$.

Note that $|W \cap W_2| \leq 2$ and $|W \cap W_3| \leq 1$. If $S(x+y+z) \times F \in W$, then $W \cap W_3 = \emptyset$ and in this case, $W \subseteq \{S(x+y+z) \times F\} \cup (W \cap W_2) \cup \{(0) \times F, S \times (0)\}$ and so, $|W| \leq 5$. Suppose that $S(x+y+z) \times F \notin W$. Then as $W \subseteq (W \cap W_2) \cup (W \cap W_3) \cup \{(0) \times F, S \times (0)\}$, it follows that $|W| \leq 5$.

This proves that if W is any subset of $V(\mathcal{G}(R))$ such that the subgraph of $\mathcal{G}(R)$ induced on W is a clique, then $|W| \leq 5$.

Observe that the subgraph of $\mathcal{G}(R)$ induced on $\{Sx \times F, S(x+y) \times F, S(x+y+z) \times F, (0) \times F, S \times (0)\}$ is a clique on five vertices and so, $\omega(\mathcal{G}(R)) \geq 5$. Therefore, we obtain that $\omega(\mathcal{G}(R)) = 5$.

Now, $\chi(\mathcal{G}(R)) \geq \omega(\mathcal{G}(R)) = 5$. We claim that $\chi(\mathcal{G}(R)) > 5$. Suppose that $\chi(\mathcal{G}(R)) = 5$. Then the vertices of $\mathcal{G}(R)$ can be properly colored using a set of five distinct colors. Let $\{c_1, c_2, c_3, c_4, c_5\}$ be a set of five distinct colors that are used for a proper coloring of the vertices of $\mathcal{G}(R)$. Let $i \in \{1, 2, 3, 4, 5\}$ and let $V_i = \{I \in V(\mathcal{G}(R)) \mid I \text{ receives color } c_i\}$. Note that $V_i \neq \emptyset$ for each $i \in \{1, 2, 3, 4, 5\}$, $V_i \cap V_j = \emptyset$ for all distinct $i, j \in \{1, 2, 3, 4, 5\}$, and $V(\mathcal{G}(R)) = \cup_{i=1}^5 V_i$. Since the subgraph of $\mathcal{G}(R)$ induced on $W = \{(0) \times F, S \times (0), Sx \times F, S(x+y) \times F, S(x+y+z) \times F\}$ is a clique, no two members from W can be in the same V_i for any $i \in \{1, 2, 3, 4, 5\}$. Without loss of generality, we can assume that $(0) \times F \in V_1, S \times (0) \in V_2, Sx \times F \in V_3, S(x+y) \times F \in V_4$, and $S(x+y+z) \times F \in V_5$. Since $(0) \times F$ (respectively, $S \times (0)$) is adjacent in $\mathcal{G}(R)$ to all of its other vertices, we obtain that $V_1 = \{(0) \times F\}$ and $V_2 = \{S \times (0)\}$. Since $Sz \times F$ is adjacent to both $S(x+y) \times F$ and $S(x+y+z) \times F$ but it is not adjacent to $Sx \times F$ in $\mathcal{G}(R)$, we get that $Sz \times F$ must be in V_3 . Note that $S(x+z) \times F$ is adjacent to $Sx \times F$ and is not adjacent to any of the members from $\{S(x+y) \times F, S(x+y+z) \times F\}$ in $\mathcal{G}(R)$. Therefore, $S(x+z) \times F \in V_4 \cup V_5$. Suppose that $S(x+z) \times F \in V_4$. As $S(y+z) \times F$ is adjacent to each one of the members from $\{Sz \times F, S(x+z) \times F, S(x+y+z) \times F\}$ in $\mathcal{G}(R)$, $S(y+z) \times F \notin \cup_{i=1}^5 V_i$. This is a contradiction. Suppose that $S(x+z) \times F \in V_5$. Since $Sy \times F$ is adjacent to each one of the members from $\{Sz \times F, S(x+y) \times F, S(x+z) \times F\}$ in $\mathcal{G}(R)$, we get that $Sy \times F \notin \cup_{i=1}^5 V_i$. This is a contradiction. Therefore, $\chi(\mathcal{G}(R)) \geq 6$.

We next verify that $\chi(\mathcal{G}(R)) \leq 6$. Let $\{c_1, c_2, c_3, c_4, c_5, c_6\}$ be a set consisting of six distinct colors. We now show that the vertices of $\mathcal{G}(R)$ can be properly colored using $\{c_1, c_2, c_3, c_4, c_5, c_6\}$. Let us assign the color c_1 to $(0) \times F$, the color c_2 to $S \times (0)$, the color c_3 to $Sx \times F$, the color c_4 to $S(x+y) \times F$, the color c_5 to $S(x+y+z) \times F$, the color c_3 to each one of the members from $\{Sz \times F, (Sx + Sz) \times F, \mathbf{m} \times F, \mathbf{m}^2 \times F\}$, the color c_4 to each one of the members from $\{S(x+z) \times F, (Sz + S(x+y)) \times F\}$, the color c_5 to each one of the members from $\{Sy \times F, (Sy + S(x+z)) \times F\}$, and the color c_6 to each one of the members from $\{(Sx + Sy) \times F, S(y+z) \times F, (Sy + Sz) \times F, (Sx + S(y+z)) \times F\}$. Note that the above assignment of six colors to the vertices of $\mathcal{G}(R)$ is proper and so, $\chi(\mathcal{G}(R)) \leq 6$. Therefore, we obtain that $\chi(\mathcal{G}(R)) = 6$. \square

5. ACKNOWLEDGMENTS

We are very much thankful to the referee for many useful suggestions.

REFERENCES

- [1] D.D. Anderson and M. Naseer, *Beck's coloring of a commutative ring*, J. Algebra. 159 (2) (1993), 500–514.
- [2] M.F. Atiyah and I.G. Macdonald, *Introduction to Commutative Algebra*, Addison-Wesley, Reading, Massachusetts, (1969).
- [3] R. Balakrishnan and K. Ranganathan, *A Textbook of Graph Theory*, Universitext, Springer, (2000).
- [4] I. Beck, *Coloring of commutative rings*, J. Algebra. 116 (1988), 208–226.
- [5] M. Behboodi and Z. Rakeei, *The annihilating-ideal graph of commutative rings I* , J. Algebra Appl. 10 (4) (2011), 727–739.
- [6] M. Behboodi and Z. Rakeei, *The annihilating-ideal graph of commutative rings II* , J. Algebra Appl. 10 (4) (2011), 741–753.
- [7] R. Gilmer, *Multiplicative Ideal Theory*, Marcel-Dekker, New York, (1972).
- [8] C. Gottlieb, *Strongly prime ideals and strongly zero-dimensional rings*, J. Algebra Appl. 16 (10) (2017) Article ID. 1750191 (9 pages).
- [9] M.I. Jinnah and S.C. Mathew, *When is the comaximal graph split?*, Comm. Algebra. 40 (7) (2012), 2400–2404.
- [10] H.R. Maimani, M. Salimi, A. Sattari, and S. Yassemi, *Comaximal graph of commutative rings*, J. Algebra. 319 (2008), 1801-1808.
- [11] S.M. Moconja and Z.Z. Petrovic, *On the structure of comaximal graphs of commutative rings with identity*, Bull. Aust. Math. Soc. 83 (2011), 11–21.
- [12] K. Samei, *On the comaximal graph of a commutative ring*, Canad. Math. Bull. 57 (2) (2014), 413–423.
- [13] P.K. Sharma and S.M. Bhatwadekar, *A note on graphical representation of rings*, J. Algebra. 176 (1995), 124–127.
- [14] H.J. Wang, *Graphs associated to co-maximal ideals of commutative rings*, J. Algebra. 320 (7) (2008), 2917–2933.
- [15] M. Ye and T. Wu, *Co-maximal ideal graphs of commutative rings*, J. Algebra Appl. 11 (6) (2012), Article ID. 1250114 (14 pages).

Subramanian Visweswaran

Department of Mathematics,

Saurashtra University,

Rajkot, India.

s_visweswaran2006@yahoo.co.in

Jaydeep Parejiya

Department of Mathematics,

Saurashtra University,

Rajkot, India.

parejiyajay@gmail.com