



APPLICATIONS OF A GROUP IN GENERAL FUZZY AUTOMATA

M. HORRY

Communicated by S. Alikhani

ABSTRACT. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be a general fuzzy automaton and the set of its states be a group. The aim of this paper is the study of applications of a group in a general fuzzy automaton. For this purpose, we define the concepts of fuzzy normal kernel of a general fuzzy automaton, fuzzy kernel of a general fuzzy automaton, adjustable, multiplicative. Then we obtain the relationships between them.

1. INTRODUCTION

The concept of fuzzy automata was introduced by Wee in 1967 [11].

Let Σ be a set. A word in Σ is the product of a finite sequence of elements in Σ . Λ will denote the empty word and Σ^* the set of all words on Σ . The length $\ell(x)$ of the word $x \in \Sigma^*$ is the number of its letters, so $\ell(\Lambda) = 0$. For a nonempty set X , $\tilde{P}(X)$ will denote the set of all fuzzy sets on X and $P(X)$ will denote the set of all subsets on X .

A deterministic finite-state automaton is a five-tuple denoted as $A = (Q, \Sigma, f, T, s)$, where Q is a finite set of states, Σ is a finite set of input symbols, the total function f from $Q \times \Sigma$ into

DOI: <http://dx.doi.org/10.29252/asta.4.2.57>

MSC(2010): Primary:18B20

Keywords: (General) Fuzzy automata, group, normal subgroup, fuzzy subgroup, fuzzy normal subgroup.

Received: 10 November 2017, Accepted: 17 July 2018

*Corresponding author

Q is the state transition, T is a subset of Q of accepting states and $s \in Q$ is the initial state. A word $x = x_1x_2 \dots x_n \in \Sigma^*$ is said to be accepted by A if there exist states q_0, q_1, \dots, q_n satisfying

- (1) $q_0 = s$
- (2) $f(q_{i-1}, x_i) = q_i$ for $i = 1, 2, \dots, n$,
- (3) $q_n \in T$.

The empty word is accepted by A if and only if $s \in T$.

A nondeterministic finite-state automaton is a five-tuple denoted as $A = (Q, \Sigma, f, T, s)$, where Q is a finite set of states, Σ is a finite set of input symbols, the partial function f from $Q \times \Sigma$ into $P(Q)$ is the state transition, T is a subset of Q of accepting states and $s \in Q$ is the initial state.

A fuzzy finite-state automaton (FFA) is a six-tuple $\tilde{F} = (Q, \Sigma, R, Z, \delta, \omega)$, where Q is a finite set of states, Σ is a finite set of input symbols, R is the initial state of \tilde{F} , Z is a finite set of output symbols, $\delta : Q \times \Sigma \times Q \rightarrow [0, 1]$ is the fuzzy transition function which is used to map a state (current state) into another state (next state) upon an input symbol, attributing a value in the interval $[0, 1]$ and $\omega : Q \rightarrow Z$ is the output function. Associated with each fuzzy transition, there is a membership value in $[0, 1]$ called the weight of the transition. The transition from state q_i (current state) to state q_j (next state) upon input a_k is denoted by $\delta(q_i, a_k, q_j)$.

We use this notation to refer both to a transition and its weight. Whenever $\delta(q_i, a_k, q_j)$ is used as a value, it refers to the weight of the transition. Otherwise, it specifies the transition itself. The set of all transitions of \tilde{F} will be denoted by Δ . The above definition is generally accepted as a formal definition of a fuzzy finite-state automaton [4, 5, 6, 7, 8, 9].

In 2004, M. Doostfatemeleh and S.C. Kremer extended the notion of fuzzy automata and introduced the notion of general fuzzy automata [1].

In this paper, we define the concepts of fuzzy normal kernel of a general fuzzy automaton, fuzzy kernel of a general fuzzy automaton, adjustable, multiplicative and obtain the relationships between them.

Definition 1.1. [1] A general fuzzy automaton (GFA) is an eight-tuple machine $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \tilde{\delta}, \omega, F_1, F_2)$, where

- (i) Q is a finite set of states, $Q = \{q_1, q_2, \dots, q_n\}$,
- (ii) Σ is a finite set of input symbols, $\Sigma = \{a_1, a_2, \dots, a_m\}$,
- (iii) \tilde{R} is the set of fuzzy start states, $\tilde{R} \in \tilde{P}(Q)$,
- (iv) Z is a finite set of output symbols, $Z = \{b_1, b_2, \dots, b_k\}$,
- (v) $\omega : Q \rightarrow Z$ is the output function,
- (vi) $\tilde{\delta} : (Q \times [0, 1]) \times \Sigma \times Q \rightarrow [0, 1]$ is the augmented transition function,

- (vii) $F_1 : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is the membership assignment function,
- (viii) $F_2 : [0, 1]^* \rightarrow [0, 1]$ is called the multi-membership resolution function.

We note that the function $F_1(\mu, \delta)$ has two parameters μ and δ , where μ is the membership value of a predecessor and δ is the weight of a transition and $\delta : Q \times \Sigma \times Q \rightarrow [0, 1]$ is the fuzzy transition function which is used to map a state (current state) into another state (next state) upon an input symbol, attributing a value in the interval $[0, 1]$. In this definition, the process that takes place upon the transition from state q_i to q_j on input a_k is represented as:

$$\mu^{t+1}(q_j) = \tilde{\delta}((q_i, \mu^t(q_i)), a_k, q_j) = F_1(\mu^t(q_i), \delta(q_i, a_k, q_j)).$$

This means that $t \in \{0, 1, 2, \dots, n\}$ and the membership value (mv) of the state q_j at time $t + 1$ is computed by function F_1 using both the membership value of q_i at time t and the weight of the transition.

The usual options for the function $F_1(\mu, \delta)$ are $\max\{\mu, \delta\}$, $\min\{\mu, \delta\}$ and $(\mu + \delta)/2$.

The multi-membership resolution function resolves the multi-membership active states and assigns a single membership value to them.

Let $Q_{act}(t_i)$ be the fuzzy set of all active states at time t_i , $\forall i \geq 0$. We have $Q_{act}(t_0) = \tilde{R}$ and

$$Q_{act}(t_i) = \{(q, \mu^{t_i}(q)) : \exists q' \in Q_{act}(t_{i-1}), \exists a \in \Sigma, \delta(q', a, q) \in \Delta\}, \forall i \geq 1.$$

Since $Q_{act}(t_i)$ is a fuzzy set, in order to show that a state q belongs to $Q_{act}(t_i)$ and T is a subset of $Q_{act}(t_i)$, we should write: $q \in Domain(Q_{act}(t_i))$ and $T \subset Domain(Q_{act}(t_i))$.

Hereafter, we simply denote them as: $q \in Q_{act}(t_i)$ and $T \subset Q_{act}(t_i)$.

The combination of the operations of functions F_1 and F_2 on a multi-membership state q_j leads to the multi-membership resolution algorithm.

Algorithm 1. [1] (*Multi-membership resolution*) *If there are several simultaneous transitions to the active state q_j at time $t + 1$, the following algorithm will assign a unified membership value to it:*

(1) *Each transition weight $\delta(q_i, a_k, q_j)$ together with $\mu^t(q_i)$, will be processed by the membership assignment function F_1 , and will produce a membership value. Call this v_i .*

$$v_i = \tilde{\delta}((q_i, \mu^t(q_i)), a_k, q_j) = F_1(\mu^t(q_i), \delta(q_i, a_k, q_j)).$$

(2) *These membership values are not necessarily equal. Hence, they need to be processed by the multi-membership resolution function F_2 .*

(3) *The result produced by F_2 will be assigned as the instantaneous membership value of the active state q_j ,*

$$\mu^{t+1}(q_j) = F_2[v_i] = F_2[F_1(\mu^t(q_i), \delta(q_i, a_k, q_j))].$$

where

- n is the number of simultaneous transitions to the active state q_j at time $t + 1$.
- $\delta(q_i, a_k, q_j)$ is the weight of a transition from q_i to q_j upon input a_k .
- $\mu^t(q_i)$ is the membership value of q_i at time t .
- $\mu^{t+1}(q_j)$ is the final membership value of q_j at time $t + 1$.

Definition 1.2. [12] Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be a general fuzzy automaton. We define max-min general fuzzy automata as $\tilde{F}^* = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}^*, F_1, F_2)$ such that :

$$\tilde{\delta}^* : Q_{act} \times \Sigma^* \times Q \rightarrow [0, 1]$$

where $Q_{act} = \{Q_{act}(t_0), Q_{act}(t_1), Q_{act}(t_2), \dots\}$ and for all $i \geq 0$,

$$\tilde{\delta}^*((q, \mu^{t_i}(q)), \Lambda, p) = \begin{cases} 1, & q = p, \\ 0, & \text{otherwise} \end{cases}$$

Also, if the input at time t_i be u_i , where $u_i \in \Sigma, \forall 1 \leq i \leq n$, then

$$\begin{aligned} \tilde{\delta}^*((q, \mu^{t_{i-1}}(q)), u_i, p) &= \tilde{\delta}((q, \mu^{t_{i-1}}(q)), u_i, p), \\ \tilde{\delta}^*((q, \mu^{t_{i-1}}(q)), u_i u_{i+1}, p) &= \bigvee_{q' \in Q_{act}(t_i)} (\tilde{\delta}((q, \mu^{t_{i-1}}(q)), u_i, q') \wedge \tilde{\delta}((q', \mu^{t_i}(q')), u_{i+1}, p)), \end{aligned}$$

and recursively

$$\begin{aligned} \tilde{\delta}^*((q, \mu^{t_0}(q)), u_1 u_2 \dots u_n, p) &= \bigvee \{ \tilde{\delta}((q, \mu^{t_0}(q)), u_1, p_1) \wedge \tilde{\delta}((p_1, \mu^{t_1}(p_1)), u_2, p_2) \wedge \dots \\ &\wedge \tilde{\delta}((p_{n-1}, \mu^{t_{n-1}}(p_{n-1})), u_n, p) \mid p_1 \in Q_{act}(t_1), p_2 \in Q_{act}(t_2), \dots, p_{n-1} \in Q_{act}(t_{n-1}) \}. \end{aligned}$$

If $q \in Q_{act}(t_i)$, we should write q belongs to an element of Q_{act} . Hereafter, we simply denote it as: $q \in Q_{act}$.

Definition 1.3. [5] A fuzzy subset μ of X is a function of X into the closed interval $[0,1]$. The support of μ is defined to be the set,

$$Supp(\mu) = \{x \in X \mid \mu(x) > 0\}$$

Definition 1.4. [5] Let λ and μ be fuzzy subsets of G . The product $\lambda * \mu$ of λ and μ is defined by

$$(\lambda * \mu)(x) = \bigvee \{ \lambda(y) \wedge \mu(z) \mid y, z \in G, x = y * z \}$$

for all $x \in G$.

We let $FP(X)$ denote the fuzzy power set of X . So $FP(X)$ is the set of all fuzzy subsets of X .

$FP(X)$ is a semigroup with respect to the product of fuzzy subsets of X .

Definition 1.5. [5] Let $(G, *)$ be a group. A fuzzy subset λ of G is called a fuzzy subgroup of G if the following properties hold:

i) $\lambda(x * y) \geq \lambda(x) \wedge \lambda(y)$

ii) $\lambda(x) = \lambda(x^{-1})$

for all $x, y \in G$.

Definition 1.6. [5] A fuzzy subgroup λ of G is called a fuzzy normal subgroup of G if

$$\lambda(x * y * x^{-1}) \geq \lambda(y)$$

for all $x, y \in G$.

Definition 1.7. [5] Let λ and μ be fuzzy subsets of G and $\lambda \subseteq \mu$. Then λ is called a fuzzy normal subgroup of μ if

$$\lambda(x * y * x^{-1}) \geq \lambda(x) \wedge \lambda(y)$$

for all $x, y \in G$.

2. Applications of a group in general fuzzy automata

Definition 2.1. Let $\tilde{F}_1 = (Q_1, \Sigma_1, \tilde{R}_1, Z, \omega, \tilde{\delta}_1, F_1, F_2)$ and $\tilde{F}_2 = (Q_2, \Sigma_2, \tilde{R}_2, Z, \omega, \tilde{\delta}_2, F_1, F_2)$ be general fuzzy automata and $(Q_1, *)$ and $(Q_2, *)$ be groups. A pair of functions (f, g) , where $f : Q_1 \rightarrow Q_2, g : \Sigma_1 \rightarrow \Sigma_2$, is called a homomorphism from \tilde{F}_1 into \tilde{F}_2 if the following conditions hold:

i) f is a group homomorphism,

ii) $\tilde{\delta}_1((p, \mu^t(p)), x, q) \leq \tilde{\delta}_2((f(p), \mu^t(f(p))), g(x), f(q)), \forall p, q \in Q, x \in \Sigma_1$.

Definition 2.2. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be a general fuzzy automaton and let λ be a fuzzy subset of Q and $(Q, *)$ be a group. λ is called fuzzy normal kernel of \tilde{F} if the following conditions hold:

i) λ is a fuzzy normal subgroup of Q ,

ii) $\lambda(p * r^{-1}) \geq \tilde{\delta}((q * k, \mu^t(q * k)), x, p) \wedge \tilde{\delta}((q, \mu^t(q)), x, r) \wedge \lambda(k)$

for all $p, q, k, r \in Q, x \in \Sigma$.

Theorem 2.3. Let $\tilde{F}^* = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}^*, F_1, F_2)$ be a max-min general fuzzy automaton and let λ is a fuzzy normal subgroup of Q and $(Q, *)$ be a group. Then λ is a fuzzy normal kernel of \tilde{F}^* if and only if

$$\lambda(p * r^{-1}) \geq \tilde{\delta}^*((q * k, \mu^t(q * k)), x, p) \wedge \tilde{\delta}^*((q, \mu^t(q)), x, r) \wedge \lambda(k)$$

for all $p, q, k, r \in Q, x \in \Sigma^*$.

Proof. Let λ be a fuzzy normal kernel of \tilde{F}^* . We prove the theorem by induction on $|x| = n$. Let $n = 0$. Then $x = \Lambda$. If $p = q * k$, $r = q$, then since λ is a fuzzy normal subgroup, we have

$$\tilde{\delta}^*((q * k, \mu^t(q * k)), x, p) \wedge \tilde{\delta}^*((q, \mu^t(q)), x, r) \wedge \lambda(k) = \lambda(k) \leq \lambda(q * k * q^{-1}) = \lambda(p * r^{-1})$$

If $p \neq q * k$ or $r \neq q$, then

$$\tilde{\delta}^*((q * k, \mu^t(q * k)), x, p) \wedge \tilde{\delta}^*((q, \mu^t(q)), x, r) \wedge \lambda(k) = 0 \leq \lambda(p * r^{-1})$$

Thus the result holds for $n = 0$. Suppose that the result holds for all $x \in \Sigma^*$ such that $|x| = n - 1$. Let $x = ya$, $y \in \Sigma^*$, $|y| = n - 1$, $n > 0$. Then we have

$$\begin{aligned} & \tilde{\delta}^*((q * k, \mu^t(q * k)), x, p) \wedge \tilde{\delta}^*((q, \mu^t(q)), x, r) \wedge \lambda(k) \\ &= \tilde{\delta}^*((q * k, \mu^t(q * k)), ya, p) \wedge \tilde{\delta}^*((q, \mu^t(q)), ya, r) \wedge \lambda(k) \\ &= (\bigvee_{u \in Q} \tilde{\delta}^*((q * k, \mu^t(q * k)), y, u) \wedge \tilde{\delta}^*((u, \mu^t(u)), a, p)) \wedge \\ & \quad (\bigvee_{v \in Q} \tilde{\delta}^*((q, \mu^t(q)), y, v) \wedge \tilde{\delta}^*((v, \mu^t(v)), a, r)) \wedge \lambda(k) \\ &= \bigvee_{u \in Q} \bigvee_{v \in Q} \tilde{\delta}^*((q * k, \mu^t(q * k)), y, u) \wedge \tilde{\delta}^*((u, \mu^t(u)), a, p) \wedge \\ & \quad \tilde{\delta}^*((q, \mu^t(q)), y, v) \wedge \tilde{\delta}^*((v, \mu^t(v)), a, r) \wedge \lambda(k) \\ & \leq \bigvee_{u \in Q} \bigvee_{v \in Q} \lambda(u * v^{-1}) \wedge \tilde{\delta}^*((u, \mu^t(u)), a, p) \wedge \\ & \quad \tilde{\delta}^*((v, \mu^t(v)), a, r) \\ & \leq \bigvee_{u \in Q} \bigvee_{v \in Q} \lambda(v^{-1} * u) \wedge \tilde{\delta}^*((v * v^{-1} * u, \mu^t(v * v^{-1} * u)), a, p) \wedge \\ & \quad \tilde{\delta}^*((v, \mu^t(v)), a, r) \\ &= \bigvee_{u \in Q} \bigvee_{v \in Q} \lambda(k') \wedge \tilde{\delta}^*((v * k', \mu^t(v * k')), a, p) \wedge \tilde{\delta}^*((v, \mu^t(v)), a, r) \\ & \leq \lambda(p * r^{-1}) \end{aligned}$$

The converse is trivial. \square

Definition 2.4. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be a general fuzzy automaton and let λ be a fuzzy subset of Q and $(Q, *)$ be a group. λ is called fuzzy kernel of \tilde{F} if the following conditions hold:

- i) λ is a fuzzy subgroup of Q ,
- ii) $\lambda(p) \geq \tilde{\delta}^*((q, \mu^t(q)), x, p) \wedge \lambda(q)$ for all $p, q \in Q$, $x \in \Sigma$.

Theorem 2.5. Let $\tilde{F}^* = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}^*, F_1, F_2)$ be a max-min general fuzzy automaton and let λ is a fuzzy subgroup of Q and $(Q, *)$ be a group. Then λ is a fuzzy kernel of \tilde{F}^* if and only if $\lambda(p) \geq \tilde{\delta}^*((q, \mu^t(q)), x, p) \wedge \lambda(q)$ for all $p, q \in Q$, $x \in \Sigma^*$.

Proof. The proof is similar to that of Theorem 2.3. \square

Definition 2.6. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be a general fuzzy automaton and $(Q, *)$ be a group. \tilde{F} is called adjustable if

$$\tilde{\delta}((p * q, \mu^t(p * q)), x, p) \leq \tilde{\delta}((p, \mu^t(p)), x, k)$$

for all $p, q, r, k \in Q, x \in \Sigma$.

Theorem 2.7. Let $\tilde{F}^* = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}^*, F_1, F_2)$ be an adjustable general fuzzy automaton, $(Q, *)$ be a group and let λ be a fuzzy normal kernel of \tilde{F} and ν be a fuzzy kernel of \tilde{F} . Then $\lambda * \nu$ is a fuzzy kernel of \tilde{F}^* .

Proof. Since λ is a fuzzy normal subgroup of Q and ν is a fuzzy subgroup of Q , it follows that $\lambda * \nu$ is a fuzzy subgroup of Q and $\lambda * \nu = \nu * \lambda$. Since $p = (p * r^{-1}) * r$, then we have

$$\begin{aligned} (\lambda * \nu)(p) &\geq \lambda(p * r^{-1}) \wedge \nu(r) \\ &\geq (\tilde{\delta}((a * b, \mu^t(a * b)), x, p) \wedge \tilde{\delta}((a, \mu^t(a)), x, r) \wedge \lambda(b)) \wedge (\tilde{\delta}((a, \mu^t(a)), x, r) \wedge \nu(a)) \end{aligned}$$

for all $a, b, p \in Q, x \in \Sigma$.

Since \tilde{F} is adjustable, so we have

$$\tilde{\delta}((a * b, \mu^t(a * b)), x, p) \leq \tilde{\delta}((a, \mu^t(a)), x, r)$$

Then we have

$$\begin{aligned} &(\tilde{\delta}((a * b, \mu^t(a * b)), x, p) \wedge \tilde{\delta}((a, \mu^t(a)), x, r) \wedge \lambda(b)) \wedge (\tilde{\delta}((a, \mu^t(a)), x, r) \wedge \nu(a)) \\ &= \tilde{\delta}((a * b, \mu^t(a * b)), x, p) \wedge \lambda(b) \wedge \nu(a) \end{aligned}$$

Thus for all $p, q \in Q, x \in \Sigma$,

$$\begin{aligned} (\lambda * \nu)(p) &\geq \vee \{ \tilde{\delta}((a * b, \mu^t(a * b)), x, p) \wedge \lambda(b) \wedge \nu(a) \mid a, b \in Q, a * b = q \} \\ &= \tilde{\delta}((q, \mu^t(q)), x, p) \wedge (\vee \{ \lambda(b) \wedge \nu(a) \mid a, b \in Q, a * b = q \}) \\ &= \tilde{\delta}((q, \mu^t(q)), x, p) \wedge (\nu * \lambda)(q) \\ &= \tilde{\delta}((q, \mu^t(q)), x, p) \wedge (\lambda * \nu)(q) \end{aligned}$$

Hence $\lambda * \nu$ is a fuzzy kernel of \tilde{F} . \square

Theorem 2.8. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be an adjustable general fuzzy automaton and $(Q, *)$ be a group. If λ and ν are fuzzy normal kernels of \tilde{F} , then $\lambda * \nu$ is a fuzzy normal kernel of \tilde{F} .

Proof. Since λ and ν are fuzzy normal subgroups of Q , it follows that $\lambda * \nu$ is a fuzzy normal subgroup of Q and $\lambda * \nu = \nu * \lambda$. Since $p * r^{-1} = (p * q^{-1}) * (q * r^{-1})$, then we have

$$\begin{aligned} (\lambda * \nu)(p * r^{-1}) &\geq \lambda(p * q^{-1}) \wedge \nu(q * r^{-1}) \\ &\geq (\tilde{\delta}((a * b * c, \mu^t(a * b * c)), x, p) \wedge \tilde{\delta}((a * b, \mu^t(a * b)), x, q) \wedge \lambda(c)) \\ &\quad \wedge (\tilde{\delta}((a * b, \mu^t(a * b)), x, q) \wedge \tilde{\delta}((a, \mu^t(a)), x, r) \wedge \nu(b)) \end{aligned}$$

for all $a, b, c, p, r \in Q, x \in \Sigma$.

Since \tilde{F} is adjustable, so we have

$$\tilde{\delta}((a * b * c, \mu^t(a * b * c)), x, p) \leq \tilde{\delta}((a * b, \mu^t(a * b)), x, q)$$

Then we have

$$\begin{aligned} &\tilde{\delta}((a * b * c, \mu^t(a * b * c)), x, p) \wedge \tilde{\delta}((a * b, \mu^t(a * b)), x, q) \wedge \lambda(c) \\ &\quad \wedge \tilde{\delta}((a * b, \mu^t(a * b)), x, q) \wedge \tilde{\delta}((a, \mu^t(a)), x, r) \wedge \nu(b) \\ &= \tilde{\delta}((a * b * c, \mu^t(a * b * c)), x, p) \wedge \tilde{\delta}((a, \mu^t(a)), x, r) \wedge \lambda(c) \wedge \nu(b) \end{aligned}$$

Thus for all $p, q, r, k \in Q, x \in \Sigma$, we have

$$\begin{aligned} &(\lambda * \nu)(p * r^{-1}) \geq \\ &\vee \{ \tilde{\delta}((a * b * c, \mu^t(a * b * c)), x, p) \wedge \tilde{\delta}((a, \mu^t(a)), x, r) \wedge \lambda(c) \wedge \nu(b) \mid b, c \in Q, b * c = k \} \\ &= \tilde{\delta}((a * k, \mu^t(a * k)), x, p) \wedge \tilde{\delta}((a, \mu^t(a)), x, r) \wedge (\vee \{ \lambda(c) \wedge \nu(b) \mid b, c \in Q, b * c = k \}) \\ &= \tilde{\delta}((a * k, \mu^t(a * k)), x, p) \wedge \tilde{\delta}((a, \mu^t(a)), x, r) \wedge (\nu * \lambda)(k) \\ &= \tilde{\delta}((a * k, \mu^t(a * k)), x, p) \wedge \tilde{\delta}((a, \mu^t(a)), x, r) \wedge (\lambda * \nu)(k) \end{aligned}$$

Hence $\lambda * \nu$ is a fuzzy normal kernel of \tilde{F} . \square

Theorem 2.9. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be a general fuzzy automaton, $(Q, *)$ be a group and let λ be a fuzzy normal kernel of \tilde{F} with $\text{Supp}(\lambda) = Q$. Then there exists a general fuzzy automaton say \tilde{H} and there exists a homomorphism from \tilde{F} to \tilde{H} .

Proof. Now λ is a fuzzy normal subgroup of Q . Define $\lambda_p : Q \rightarrow [0, 1]$ by $\lambda_p(q) = \lambda(p * q^{-1})$ for all $q \in Q$. Let $E = \{\lambda_p | p \in Q\}$. Then E is a group with respect to the binary operation $\lambda_p * \lambda_q = \lambda_{p*q}$ for all $\lambda_p, \lambda_q \in E$. Define $Q/\lambda : E \rightarrow [0, 1]$ by $(Q/\lambda)(\lambda_p) = \lambda(p)$ for all $\lambda_p \in E$. Thus Q/λ is a fuzzy subgroup of E . Let $G = Supp(Q/\lambda)$. Define $\tilde{\sigma} : (G \times [0, 1]) \times \Sigma \times G \rightarrow [0, 1]$ by

$$\tilde{\sigma}((\lambda_p, \mu^t(\lambda_p)), x, \lambda_q) = \vee \{ \tilde{\delta}((a, \mu^t(a)), x, b) | a, b \in Q, \lambda_a = \lambda_p, \lambda_b = \lambda_q \}$$

for all $\lambda_p, \lambda_q \in G$. Define $f : Q \rightarrow G$ by $f(q) = \lambda_q$ for all $q \in Q$ and let $g : \Sigma \rightarrow \Sigma$ be identity map. Then we have

$$f(p * q) = \lambda_{p*q} = \lambda_p * \lambda_q = f(p) * f(q)$$

for all $p, q \in Q$.

Thus (f, g) is a homomorphism from \tilde{F} to $\tilde{H} = (G, \Sigma, \tilde{R}, Z, \omega, \tilde{\sigma}, F_1, F_2)$. \square

Definition 2.10. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be a general fuzzy automaton and let λ be a fuzzy kernel of \tilde{F} and $(Q, *)$ be a group. A fuzzy set ν of Q is called a fuzzy normal kernel of λ if the following conditions hold:

- i) $\nu \subseteq \lambda$ and ν is a fuzzy normal subgroup of λ ,
- ii) $\nu(p * r^{-1}) \geq \tilde{\delta}((q * k, \mu^t(q * k)), x, p) \wedge \tilde{\delta}((q, \mu^t(q)), x, r) \wedge \nu(k)$
for all $p, k, r \in Q, q \in Supp(\lambda), x \in \Sigma$.

Definition 2.11. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be a general fuzzy automaton and $(Q, *)$ be a group. \tilde{F} is called multiplicative if having the following properties:

- i) there exists the element x_0 of Σ such that $\tilde{\delta}((e, \mu^t(e)), x_0, e) > 0$
- ii) $\tilde{\delta}((q, \mu^t(q)), x, p * r) = \tilde{\delta}((q, \mu^t(q)), x_0, p) \wedge \tilde{\delta}((e, \mu^t(e)), x, r)$,
- iii) $\tilde{\delta}((p_1 * p_2^{-1}, \mu^t(p_1 * p_2^{-1})), x_0, q_1 * q_2^{-1}) = \tilde{\delta}((p_1, \mu^t(p_1)), x_0, q_1) \wedge \tilde{\delta}((p_2, \mu^t(p_2)), x_0, q_2)$
for all $p, q, r, p_1, p_2, q_1, q_2 \in Q, x \in \Sigma$.

Theorem 2.12. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be an adjustable multiplicative general fuzzy automaton and $(Q, *)$ be an abelian group. Let λ be a fuzzy kernel of \tilde{F} , ν be a fuzzy normal kernel of λ and $Supp(\lambda) \neq \Phi$. If ν is a fuzzy normal subgroup of Q , then ν is a fuzzy normal kernel of \tilde{F} .

Proof. Since \tilde{F} is multiplicative, for all $p, q, r, k \in Q, x \in \Sigma$, we have

$$\begin{aligned} & \tilde{\delta}((q * k, \mu^t(q * k)), x, p) \wedge \tilde{\delta}((q, \mu^t(q)), x, r) \\ &= \tilde{\delta}((q * k, \mu^t(q * k)), x, p * e) \wedge \tilde{\delta}((q, \mu^t(q)), x, r * e) \\ &= (\tilde{\delta}((q * k, \mu^t(q * k)), x_0, p) \wedge \tilde{\delta}((e, \mu^t(e)), x, e)) \wedge (\tilde{\delta}((q, \mu^t(q)), x_0, r) \wedge \tilde{\delta}((e, \mu^t(e)), x, e)) \\ &= (\tilde{\delta}((q * k, \mu^t(q * k)), x_0, p * e) \wedge \tilde{\delta}((e, \mu^t(e)), x, e)) \wedge (\tilde{\delta}((q, \mu^t(q)), x_0, r) \wedge \tilde{\delta}((e, \mu^t(e)), x, e)) \end{aligned}$$

$$= \tilde{\delta}((q, \mu^t(q)), x_0, p) \wedge \tilde{\delta}((k^{-1}, \mu^t(k^{-1})), x_0, e) \wedge \tilde{\delta}((e, \mu^t(e)), x, e) \wedge \tilde{\delta}((q, \mu^t(q)), x_0, r)$$

Now for any $q \in Q$, $b \in \text{Supp}(\lambda)$, $q = b * (q^{-1} * b)^{-1}$. Thus we have

$$\begin{aligned} & \tilde{\delta}((q * k, \mu^t(q * k)), x, p) \wedge \tilde{\delta}((q, \mu^t(q)), x, r) \\ &= \tilde{\delta}((b * (q^{-1} * b)^{-1} * k, \mu^t(b * (q^{-1} * b)^{-1} * k)), x, p * e) \wedge \tilde{\delta}((e * q, \mu^t(e * q)), x, r * e) \\ &= \tilde{\delta}((b, \mu^t(b)), x_0, p) \wedge \tilde{\delta}((q^{-1} * b, \mu^t(q^{-1} * b)), x_0, e) \wedge \\ & \tilde{\delta}((k^{-1}, \mu^t(k^{-1})), x_0, e) \wedge \tilde{\delta}((e, \mu^t(e)), x, e) \wedge \tilde{\delta}((q^{-1}, \mu^t(q^{-1})), x_0, r) \\ &= \tilde{\delta}((b, \mu^t(b)), x_0, p) \wedge \tilde{\delta}((k^{-1}, \mu^t(k^{-1})), x_0, e) \wedge \tilde{\delta}((e, \mu^t(e)), x, e) \wedge \\ & (\tilde{\delta}((q^{-1} * b, \mu^t(q^{-1} * b)), x_0, e) \wedge \tilde{\delta}((q^{-1}, \mu^t(q^{-1})), x_0, r)) \\ &= \tilde{\delta}((b, \mu^t(b)), x_0, p) \wedge \tilde{\delta}((k^{-1}, \mu^t(k^{-1})), x_0, e) \wedge \tilde{\delta}((e, \mu^t(e)), x, e) \wedge \tilde{\delta}((b, \mu^t(b)), x_0, r) \end{aligned}$$

Since ν is a fuzzy normal kernel of λ , for all $p, q, r \in Q$, $b \in \text{Supp}(\lambda)$, $x \in \Sigma$, we have

$$\begin{aligned} \nu(p * r^{-1}) &\geq \tilde{\delta}((b * k, \mu^t(b * k)), x, p) \wedge \tilde{\delta}((b, \mu^t(b)), x, r) \wedge \nu(k) \\ &= \tilde{\delta}((b * k, \mu^t(b * k)), x, p * e) \wedge \tilde{\delta}((b, \mu^t(b)), x, r * e) \wedge \nu(k) \\ &= \tilde{\delta}((b, \mu^t(b)), x_0, p) \wedge \tilde{\delta}((k^{-1}, \mu^t(k^{-1})), x_0, e) \wedge \\ & \tilde{\delta}((b, \mu^t(b)), x_0, r) \wedge \tilde{\delta}((e, \mu^t(e)), x, e) \wedge \nu(k) \\ &= \tilde{\delta}((q * k, \mu^t(q * k)), x, p) \wedge \tilde{\delta}((q, \mu^t(q)), x, r) \wedge \nu(k) \end{aligned}$$

Hence ν is a fuzzy normal kernel of \tilde{F} . \square

Theorem 2.13. Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be an adjustable multiplicative general fuzzy automaton, $(Q, *)$ be a group, λ be a fuzzy subset of Q and $\text{Supp}(\lambda) \neq \Phi$. Then the following statements are equivalent:

- i) λ is a fuzzy normal kernel of \tilde{F}
- ii) λ is a fuzzy normal subgroup of Q and $\lambda(q) \geq \tilde{\delta}((p, \mu^t(p)), x_0, q) \wedge \lambda(p)$, for all $p, q \in Q$.

Proof. ($i \Rightarrow ii$) Since λ is a fuzzy normal kernel of \tilde{F} , we have:

$$\begin{aligned} \lambda(q) &= \lambda(q * e^{-1}) \\ &\geq \tilde{\delta}((e * p, \mu^t(e * p)), x_0, q) \wedge \tilde{\delta}((e, \mu^t(e)), x_0, e) \wedge \lambda(p) \\ &= \tilde{\delta}((p, \mu^t(p)), x_0, q) \wedge \lambda(p) \end{aligned}$$

(Since \tilde{F} is adjustable, for all $p, q \in Q$, we have

$$\tilde{\delta}((p, \mu^t(p)), x_0, q) = \tilde{\delta}((e * p, \mu^t(e * p)), x_0, q) \leq \tilde{\delta}((e, \mu^t(e)), x_0, e))$$

(ii \Rightarrow i) Since \tilde{F} is multiplicative, for all $p, q, r, k \in Q$, $x \in \Sigma$, we have

$$\begin{aligned}
 & \tilde{\delta}((q * k, \mu^t(q * k)), x, p * r) \wedge \tilde{\delta}((q, \mu^t(q)), x, p) \wedge \lambda(k) \\
 &= \tilde{\delta}((q * k, \mu^t(q * k)), x, p * r * e) \wedge \tilde{\delta}((q, \mu^t(q)), x, p * e) \wedge \lambda(k) \\
 &= (\tilde{\delta}((q * k, \mu^t(q * k)), x_0, p * r) \wedge \tilde{\delta}((e, \mu^t(e)), x, e)) \wedge \\
 & \quad (\tilde{\delta}((q, \mu^t(q)), x_0, p) \wedge \tilde{\delta}((e, \mu^t(e)), x, e)) \wedge \lambda(k) \\
 &= \tilde{\delta}((q, \mu^t(q)), x_0, p) \wedge \tilde{\delta}((k^{-1}, \mu^t(k^{-1})), x_0, r^{-1}) \wedge \tilde{\delta}((e, \mu^t(e)), x, e) \wedge \lambda(k) \\
 &= \tilde{\delta}((q, \mu^t(q)), x_0, p) \wedge \tilde{\delta}((e * k^{-1}, \mu^t(e * k^{-1})), x_0, e * r^{-1}) \wedge \tilde{\delta}((e, \mu^t(e)), x, e) \wedge \lambda(k) \\
 & \quad = \tilde{\delta}((q, \mu^t(q)), x_0, p) \wedge \\
 & \quad (\tilde{\delta}((e, \mu^t(e)), x_0, e) \wedge \tilde{\delta}((k, \mu^t(k)), x_0, r)) \\
 & \quad \wedge \tilde{\delta}((e, \mu^t(e)), x, e) \wedge \lambda(k) \\
 & \leq \tilde{\delta}((k, \mu^t(k)), x_0, r) \wedge \lambda(k) \\
 & \leq \lambda(r) \leq \lambda(p * r * p^{-1})
 \end{aligned}$$

Now let $p_1, p_2, q, k \in Q$, $x \in \Sigma$. Since Q is a group, then there exists a unique element $r \in Q$ such that $p_1 = p_2 * r$. Thus

$$\begin{aligned}
 & \lambda(p_1 * p_2^{-1}) = \lambda(p_2 * r * p_2^{-1}) \\
 & \geq \tilde{\delta}((q * k, \mu^t(q * k)), x, p_2 * r) \wedge \tilde{\delta}((q, \mu^t(q)), x, p_2) \wedge \lambda(k) \\
 & = \tilde{\delta}((q * k, \mu^t(q * k)), x, p_1) \wedge \tilde{\delta}((q, \mu^t(q)), x, p_2) \wedge \lambda(k)
 \end{aligned}$$

Hence λ is a fuzzy normal kernel of \tilde{F} . \square

Theorem 2.14. *Let $\tilde{F} = (Q, \Sigma, \tilde{R}, Z, \omega, \tilde{\delta}, F_1, F_2)$ be an adjustable multiplicative general fuzzy automaton and $(Q, *)$ be a group. Then there exists a semigroup homomorphism $f : Q \rightarrow FP(Q)$ and there exists a function $g : \Sigma \rightarrow FP(Q)$ such that $\tilde{\delta}((q, \mu^t(q)), x, p) = f(q)(p) \wedge g(x)(e)$, for all $p, q \in Q$, $x \in \Sigma$.*

Proof. Define $f : Q \rightarrow FP(Q)$ by $f(q) = \mu_q$ for all $q \in Q$, where $\mu_q : Q \rightarrow [0, 1]$ is defined by $\mu_q(p) = \tilde{\delta}((q, \mu^t(q)), x_0, p)$ for all $p \in Q$. Since \tilde{F} is multiplicative, for all $p_1, p_2, q_1, q_2 \in Q$, we have

$$\begin{aligned} \mu_{q_1 * q_2}(p_1 * p_2) &= \tilde{\delta}((q_1 * q_2, \mu^t(q_1 * q_2)), x_0, p_1 * p_2) \\ &= \tilde{\delta}((q_1, \mu^t(q_1)), x_0, p_1) \wedge \tilde{\delta}((q_2^{-1}, \mu^t(q_2^{-1})), x_0, p_2^{-1}) \\ &= \tilde{\delta}((q_1, \mu^t(q_1)), x_0, p_1) \wedge \tilde{\delta}((q_2, \mu^t(q_2)), x_0, p_2) \\ &= \mu_{q_1}(p_1) \wedge \mu_{q_2}(p_2) \end{aligned}$$

Also we have

$$(\mu_{q_1} * \mu_{q_2})(p) = \vee \{ \mu_{q_1}(q) \wedge \mu_{q_2}(r) \mid q, r \in Q, p = q * r \} = \mu_{q_1 * q_2}(p)$$

for all $p \in Q$. Hence

$$f(q_1 * q_2) = f(q_1) * f(q_2)$$

For $x \in \Sigma$, define the fuzzy subset λ_x of Q by $\lambda_x(q) = \tilde{\delta}((e, \mu^t(e)), x, q)$ for all $q \in Q$. Define $g : \Sigma \rightarrow FP(Q)$ by $g(x) = \lambda_x$ for all $x \in \Sigma$. Since \tilde{F} is multiplicative, for all $p, q \in Q, x \in \Sigma$, we have

$$\begin{aligned} &f(q)(p) \wedge g(x)(e) \\ &= \mu_q(p) \wedge \lambda_x(e) = \tilde{\delta}((q, \mu^t(q)), x_0, p) \wedge \tilde{\delta}((e, \mu^t(e)), x, e) = \tilde{\delta}((q, \mu^t(q)), x, p) \end{aligned}$$

□

3. Conclusion

In this paper, we defined the concepts of fuzzy normal kernel of a general fuzzy automaton, fuzzy kernel of a general fuzzy automaton, adjustable, multiplicative. Then we obtained the relationships between them.

4. ACKNOWLEDGMENTS

The author wish to sincerely thank the referees for several useful comments.

REFERENCES

- [1] M. Doostfateme and S. C. Kremer, *New directions in fuzzy automata*, International Journal of Approximate Reasoning, **38** (2005), 175–214.
- [2] M. Horry and M. M. Zahedi, *Fuzzy subautomata of an invertible general fuzzy automaton*, Annals of fuzzy sets, fuzzy logic and fuzzy systems, **2(2)**(2013), 29–47.
- [3] J. Jin, Q. Li, Y. Li, *Algebraic properties of L-fuzzy finite automata*, Information Sciences, **234** (2013), 182-202.
- [4] Y. Li, W. Pedrycz, *Fuzzy finite automata and fuzzy regular expressions with membership values in lattice-ordered monoids*, Fuzzy Sets and Systems, **156** (2005), 68–92.
- [5] J. N. Mordeson and D. S. Malik, *Fuzzy automata and languages, theory and applications*, Chapman and Hall/CRC, London/Boca Raton, FL, 2002.
- [6] D. S. Malik, J. N. Mordeson and M. K. Sen, *On subsystems of fuzzy finite state machines*, Fuzzy Sets and Systems, **68** (1994), 83–92.
- [7] M. Mizumoto, J. Tanaka and K. Tanaka, *Some consideration on fuzzy automata*, J. Compute. Systems Sci. **3** (1969), 409–422.
- [8] W. Omlin, K. K. Giles and K. K. Thornber, *Equivalence in knowledge representation: automata, rnns, and dynamic fuzzy systems*, Proc. IEEE, **87(9)**(1999), 1623–1640.
- [9] W. Omlin, K. K. Thornber and K. K. Giles, *Fuzzy finite-state automata can be deterministically encoded into recurrent neural networks*, IEEE Trans. Fuzzy Syst. **5(1)**(1998), 76–89.
- [10] E. S. Santos, *Realization of fuzzy languages by probabilistic, max-prod and maximin automata*, Inform. Sci. **8** (1975), 39–53.
- [11] W. G. Wee, *On generalization of adaptive algorithm and application of the fuzzy sets concept to pattern classification*, Ph.D. dissertation Purdue University, IN, 1967.
- [12] M. M. Zahedi, M. Horry and Kh. Abolpor, *Bifuzzy (General) topology on max-min general fuzzy automata*, Advanced in Fuzzy Mathematics, **3(1)**(2008), 51–68.

Mohammad Horry

Department of mathematics,

Shahid Chamran university, Kerman, Iran.

`mhori@tvu.ac.ir`