



Research Paper

**ON THE ASSOCIATED PRIMES OF THE GENERALIZED d -LOCAL
COHOMOLOGY MODULES**

Z. RAHIMI-MOLAEI, SH. PAYROVI AND S. BABAEI

ABSTRACT. The first part of the paper is concerned to relationship between the sets of associated primes of the generalized d -local cohomology modules and the ordinary generalized local cohomology modules. Assume that R is a commutative Noetherian local ring, M and N are finitely generated R -modules and d, t are two integers. We prove that $\text{Ass } H_d^t(M, N) = \bigcup_{I \in \Phi} \text{Ass } H_I^t(M, N)$ whenever $H_d^i(M, N) = 0$ for all $i < t$ and $\Phi = \{I : I \text{ is an ideal of } R \text{ with } \dim R/I \leq d\}$. In the second part of the paper, we give some information about the non-vanishing of the generalized d -local cohomology modules. To be more precise, we prove that $H_d^i(M, R) \neq 0$ if and only if $i = n - d$ whenever R is a Gorenstein ring of dimension n and $\text{pd}_R(M) < \infty$. This result leads to an example which shows that $\text{Ass } H_d^{n-d}(M, R)$ is not necessarily a finite set.

DOI: 10.29252/as.2020.1615

MSC(2010): Primary 13D45; Secondary 14B15.

Keywords: Gorenstein Ring, Associated prime ideals, d -local cohomology modules.

Received: 27 April 2019, Accepted: 05 Aug 2019.

*Corresponding author

1. INTRODUCTION

Throughout this paper, R denotes a commutative Noetherian ring with non-zero identity. For an ideal I of R and R -modules M and N , the i th generalized local cohomology module of M and N with respect to I is defined as

$$H_I^i(M, N) \cong \varinjlim_{n \in \mathbb{N}} \text{Ext}_R^i(M/I^n M, N).$$

The reader can refer to [9, 18] for the basic properties of generalized local cohomology modules. An important problem in commutative algebra is determining when the set of associated primes of the i th local cohomology module $H_I^i(M)$ is finite. In [10], raised the following question: If M is a finitely generated R -module, then the set of associated primes of $H_I^i(M)$ is finite for all ideals I of R and all $i \geq 0$. In [11, 15, 17] the authors have given counterexamples to this conjecture. However, it is known that this conjecture is true in many situations; see [4, 5, 8, 13]. The purpose of this paper is to make a counterexample to above question in the context of general local cohomology modules. The theory of general local cohomology modules over commutative Noetherian ring introduced by M. H. Bijan-Zadeh in [3]. General local cohomology theory described as follows.

Let Φ be a non-empty set of ideals of R . We call Φ a system of ideals of R if, whenever $I, I' \in \Phi$, then there exists $J \in \Phi$ such that $J \subseteq II'$. Such a system of ideals gives rise to an additive, left exact functor

$$\Gamma_\Phi(M) = \{x \in M : Ix = 0 \text{ for some ideal } I \in \Phi\}$$

from the category of R -modules and R -homomorphisms to itself. $\Gamma_\Phi(-)$ is called the Φ -torsion functor. For each $i \geq 0$, the i th right derived functor of $\Gamma_\Phi(-)$ is denoted by $H_\Phi^i(-)$. For an ideal I of R , if $\Phi = \{I^n : n \in \mathbb{N}\}$, then $H_\Phi^i(-)$ coincides with the ordinary local cohomology functor $H_I^i(-)$. Let $d \geq 0$ be an integer. We denote $\Gamma_\Phi(-)$ and $H_\Phi^i(-)$ by $\Gamma_d(-)$ and $H_d^i(-)$ respectively, for the system of ideals $\Phi = \{I : I \text{ is an ideal of } R \text{ with } \dim R/I \leq d\}$. The functor $\Gamma_d(-)$ was originally define in [1] and the modules $H_d^i(M)$ were called d -local cohomology modules associated to M . For $i \geq 0$, we define $H_d^i(-, -) : \mathfrak{C}(R) \times \mathfrak{C}(R) \rightarrow \mathfrak{C}(R)$ with

$$H_d^i(M, N) = \varinjlim_{I \in \Phi} \text{Ext}_R^i(M/IM, N),$$

and call it the i th generalized d -local cohomology module of M and N . Then $H_d^i(-, -)$ is an additive, R -linear functor which is contravariant in the first variable and covariant in the second variable. After some preliminary results in section 2, for a finitely generated module M with $pd_R(M) < \infty$ and an integer t we prove that

$$\text{Ass}(H_d^t(M, N)) = \bigcup_{I \in \Phi} \text{Ass}(\text{Ext}_R^t(M/IM, N)) = \bigcup_{I \in \Phi} \text{Ass}(H_I^t(M, N)),$$

where $\Phi = \{I : I \text{ is an ideal of } R \text{ with } \dim R/I \leq d\}$ and $H_d^i(M, N) = 0$ for all $i < t$. In section 3, we shall provide some results concerning vanishing and non-vanishing of generalized d -local cohomology modules: we shall prove that, over a local ring R , if the non-zero finitely generated R -module M has Krull dimension n and $pd_R(M) < \infty$, then there exists an integer i with $0 \leq i \leq d$ such that $H_d^{n-i}(M, R) \neq 0$. We shall also prove that, when R is local Gorenstein of dimension n , then $H_d^i(M, R) \neq 0$ if and only if $i = n - d$. Furthermore, we shall prove that $H_d^{n-d}(M, R)$ is a non-Artinian module for which $\text{Ass}(H_d^{n-d}(M, R)) = \{\mathfrak{p} \in \text{Supp}(M) : \dim R/\mathfrak{p} = d\}$.

2. THE ASSOCIATED PRIMES

It is our intention in this section to present the relationship between the sets of associated primes of the generalized d -local cohomology modules and the ordinary generalized local cohomology modules. Its potential for use in arguments that employ the Grothendieck spectral sequence. So in this section, (R, \mathfrak{m}) is a local ring, I is an ideal of R , d is an integer and $\Phi = \{I : I \text{ is an ideal of } R \text{ with } \dim R/I \leq d\}$.

We say that M is a d -torsion module if $\Gamma_d(M) = M$, and it is d -torsion free if $\Gamma_d(M) = 0$.

Lemma 2.1. *Let M and N be finitely generated R -modules and t be an integer such that $H_d^i(M, N) = 0$ for all $i < t$. Then the following statements are true:*

- (i) $H_I^i(M, N) \subseteq H_d^i(M, N)$ for all $I \in \Phi$ and $i \leq t$.
- (ii) $H_c^i(M, N) \subseteq H_d^i(M, N)$ for all $c \leq d$ and $i \leq t$.

Proof. (i) It is clear that $\text{Hom}_R(R/I, \Gamma_d(M, N)) \cong \text{Hom}_R(M/IM, N)$ for $I \in \Phi$ so by [16, Theorem 11.38], the Grothendieck spectral sequence $E_2^{p,q} := \text{Ext}_R^p(R/I, H_d^q(M, N))$ converges to $E^{p+q} := \text{Ext}_R^{p+q}(M/IM, N)$. Because $E_\infty^{p,q}$ is a subquotient of $E_2^{p,q}$ so $E_\infty^{p,q} = 0$, for all $q < t$. On the other hand, there is a finite filtration

$$0 = F^{q+1}E^q \subseteq F^qE^q \subseteq \cdots \subseteq F^1E^q \subseteq F^0E^q = E^q$$

of E^q such that $E_\infty^{p,q-p} \cong F^pE^q/F^{p+1}E^q$, for all $p = 0, 1, \dots, q$. Thus $E_\infty^{0,q} \cong E^q$, for all $q \leq t$. Also, for all $q \leq t$, the sequence $0 \longrightarrow E_2^{0,q} \longrightarrow E_2^{2,q-1}$ and $E_2^{2,q-1} = 0$ imply that $E_\infty^{0,q} \cong E_2^{0,q}$, for all $q \leq t$. Thus $E^q \cong E_2^{0,q}$. Hence, $\text{Ext}_R^q(M/IM, N) \cong \text{Hom}_R(R/I, H_d^q(M, N))$. Therefore,

$$\Gamma_I(H_d^q(M, N)) \cong \varinjlim_{n \in \mathbb{N}} \text{Hom}(R/I^n, H_d^q(M, N)) \cong \varinjlim_{n \in \mathbb{N}} \text{Ext}_R^q(M/I^n, N) \cong H_I^q(M, N).$$

The proof is therefore complete.

- (ii) The proof is similar to that of (i). \square

Lemma 2.2. *Let M and N be finitely generated R -modules and t be an integer such that $H_d^i(M, N) = 0$, for all $i < t$. Then the following statements are true:*

- (i) $\text{Ass}(H_I^i(M, N)) = \text{Ass}(H_d^i(M, N)) \cap V(I)$, for all $I \in \Phi$ and $i \leq t$.
(ii) $\text{Ass}(H_J^i(M, N)) \subseteq \text{Ass}(H_I^i(M, N))$ for all $I, J \in \Phi$ with $I \subseteq J$ and $i \leq t$.

Proof. (i) Let $I \in \Phi$ and $i \leq t$. Then $\text{Hom}_R(R/I, H_d^i(M, N)) \cong \text{Hom}_R(R/I, H_I^i(M, N))$ and $\text{Ass}_R(\text{Hom}_R(R/I, H_d^t(M, N))) = \text{Ass}(\text{Hom}_R(R/I, H_I^t(M, N))) = \text{Ass}(H_I^t(M, N))$, by the proof of Lemma 2.1 (i) and [12, Proposition 1.1]. Thus $\text{Ass}(H_I^t(M, N)) = \text{Ass}(H_d^t(M, N)) \cap V(I)$.

(ii) It is obvious by (i). \square

Lemma 2.3. *If M is a d -torsion module, then $\text{Supp}(M) \subseteq \Phi$.*

Proof. Let $\mathfrak{p} \in \text{Supp}(M)$. Then there exists $x \in M$ such that $\text{Ann}(x) \subseteq \mathfrak{p}$. By assumption there is an ideal I of R such that $I \in \Phi$ and $Ix = 0$. Thus $I \subseteq \text{Ann}(x) \subseteq \mathfrak{p}$ and $\mathfrak{p} \in \Phi$. \square

Theorem 2.4. *Let M and N be finitely generated R -modules and t be an integer such that $H_d^i(M, N) = 0$ for all $i < t$. Then*

$$\text{Ass}(H_d^t(M, N)) = \bigcup_{I \in \Phi} \text{Ass}(\text{Ext}_R^t(M/IM, N)) = \bigcup_{I \in \Phi} \text{Ass}(H_I^t(M, N)).$$

Proof. First of all, we show that

$$\text{Ass}(H_d^t(M, N)) = \bigcup_{I \in \Phi} \text{Ass}(\text{Hom}_R(R/I, H_d^t(M, N))).$$

It is clear that $\text{Hom}_R(R/I, H_d^t(M, N)) \cong 0 :_{H_d^t(M, N)} I$. Therefore, $\text{Ass}(H_d^t(M, N)) \supseteq \bigcup_{I \in \Phi} \text{Ass}(\text{Hom}_R(R/I, H_d^t(M, N)))$. Let $\mathfrak{p} \in \text{Ass}(H_d^t(M, N))$. Then there exists $0 \neq m \in H_d^t(M, N)$ such that $\mathfrak{p} = \text{Ann}(m)$ now from Lemma 2.3 it follows that $\mathfrak{p} \in \Phi$. Thus $m \in 0 :_{H_d^t(M, N)} \mathfrak{p} \cong \text{Hom}_R(R/\mathfrak{p}, H_d^t(M, N))$ and therefore $\text{Ass}(H_d^t(M, N)) \subseteq \bigcup_{I \in \Phi} \text{Ass}(\text{Hom}_R(R/I, H_d^t(M, N)))$. On the other hand, by the proof of Lemma 2.1, $\text{Hom}_R(R/I, H_d^t(M, N)) \cong \text{Ext}_R^t(M/IM, N)$. Hence, the result follows. \square

Theorem 2.5. *Let M and N be finitely generated R -modules and t be an integer such that $H_d^i(M, N) = 0$ for all $i < t$. Then $\text{Ass}(H_d^t(M, N)) \subseteq \{\mathfrak{p} \in \text{Supp}(M) : \dim R/\mathfrak{p} = d\}$ if and only if $H_c^t(M, N) = 0$ for all integer c with $c < d$.*

Proof. Assume that $\text{Ass}(H_d^t(M, N)) \not\subseteq \{\mathfrak{p} \in \text{Supp}(M) : \dim R/\mathfrak{p} = d\}$. Thus there exists a prime ideal \mathfrak{p} belongs to $\text{Ass}(H_d^t(M, N))$ such that $\dim R/\mathfrak{p} = c < d$. So the exact sequence $0 \rightarrow \Gamma_c(R/\mathfrak{p}) \rightarrow \Gamma_c(H_d^t(M, N))$ and the fact that $\Gamma_c(R/\mathfrak{p}) = R/\mathfrak{p}$, $\Gamma_c(H_d^t(M, N)) \cong H_c^t(M, N)$ show that $\mathfrak{p} \in \text{Ass}(H_c^t(M, N))$. Hence, $H_c^t(M, N) \neq 0$. The converse is true by Lemmas 2.3 and 2.1. \square

3. THE VANISHING THEOREMS

In this section, we shall provide some results concerning the vanishing and non-vanishing of generalized d -local cohomology modules. Throughout R is a local ring with maximal ideal \mathfrak{m} and d is a non negative integer.

Theorem 3.1. *Let (R, \mathfrak{m}) be a local Gorenstein ring of dimension n and let M, N be finitely generated R -modules such that $\text{Ass}_R(M) \cap \text{Supp}(N) \neq \emptyset$ and $\text{pd}_R(M) < \infty$. Then there is at least one j with $0 \leq j \leq d$ for which $H_d^{n-j}(M, N) \neq 0$.*

Proof. It is easy to see that $\Gamma_{\mathfrak{m}}(\Gamma_d(M, N)) = \Gamma_{\mathfrak{m}}(M, N)$ so there is a Grothendieck spectral sequence

$$E_2^{i,j} = H_{\mathfrak{m}}^i(H_d^j(M, N)) \Rightarrow E^{i+j} = H_{\mathfrak{m}}^{i+j}(M, N),$$

see [16, Theorem 11.38]. By Lemma 2.3, $\text{Supp}(H_d^j(M, N)) \subseteq \Phi$. Thus $\dim H_d^j(M, N) \leq d$ and so $E_2^{i,j} = 0$ for all $i > d$, by [6, Theorem 6.1.2]. On the other hand, there is a finite filtration $0 = F^{n+1}E^n \subseteq F^n E^n \subseteq \dots \subseteq F^1 E^n \subseteq F^0 E^n = E^n$ with $E_{\infty}^{i,n-i} \cong F^i E^n / F^{i+1} E^n$. Then $F^{d+1} E^n = \dots = F^n E^n = 0$. If $E_{\infty}^{j,n-j} = 0$, for all j with $0 \leq j \leq d$, then $F^d E^n = \dots = F^0 E^n = E^n = 0$ contrary to [7, Lemma 2.4]. So suppose that $E_{\infty}^{j,n-j} \neq 0$ for some j with $0 \leq j \leq d$. Thus $E_2^{j,n-j} \neq 0$ and then $H_d^{n-j}(M, N) \neq 0$. \square

Theorem 3.2. *Let R be a Gorenstein local ring of dimension n and M be a R -module with $\text{pd}_R(M) < \infty$. Then the following statements are true:*

- (i) $H_d^i(M, R) = 0$ for all $0 \leq i < n - d$ and $H_d^i(M, R) \cong \text{Ext}_R^i(M, R)$ for all $n - d \leq i \leq n$.
- (ii) If $0 \leq d \leq \dim M$, then $\text{Ass}(H_d^{n-d}(M, R)) = \{\mathfrak{p} \in \text{Supp}(M) : \dim R/\mathfrak{p} = d\}$ and $H_d^{n-d}(M, R) \neq 0$.
- (iii) If $0 < d \leq \dim M$, then $H_d^{n-d}(M, R)$ is a Noetherian module which is not Artinian.

Proof. (i) Let $0 \rightarrow R \rightarrow E^0 \rightarrow E^1 \rightarrow \dots \rightarrow E^{n-1} \rightarrow E^n \rightarrow 0$ be a minimal injective resolution of R . Then by [14, Theorems 18.1 and 18.8] we have $E^i \cong \bigoplus_{\text{ht}(\mathfrak{p})=i} E(R/\mathfrak{p}) = \bigoplus_{\dim R/\mathfrak{p}=n-i} E(R/\mathfrak{p})$. If $i < n - d$, then $\dim R/\mathfrak{p} = n - i > d$ and so $\mathfrak{p} \notin \Phi$. Thus $\Gamma_d(E^i) = 0$ and so $\text{Hom}(M, \Gamma_d(E^i)) = 0$. It follows that $H_d^i(M, R) = 0$, for all $0 \leq i < n - d$. If $i \geq n - d$, then $\dim R/\mathfrak{p} = n - i \leq n - (n - d) = d$, and so $\mathfrak{p} \in \Phi$. Thus $\Gamma_d(E^i) = E^i$ and $\text{Hom}(M, \Gamma_d(E^i)) = \text{Hom}(M, E^i)$. It follows that $H_d^i(M, R) \cong \text{Ext}_R^i(M, R)$ for all $n - d \leq i \leq n$.

(ii) It is immediate that

$$\begin{aligned} \text{Ass}(H_d^{n-d}(M, R)) &\subseteq \text{Ass}(\text{Hom}_R(M, \bigoplus_{\dim R/\mathfrak{p}=d} E(R/\mathfrak{p}))) \\ &= \{\mathfrak{p} \in \text{Supp}(M) : \dim R/\mathfrak{p} = d\}. \end{aligned}$$

If $\mathfrak{p} \in \text{Supp}(M)$ and $\dim R/\mathfrak{p} = d$, then by [19, Lemma 2.1(5)], $(H_d^{n-d}(M, R))_{\mathfrak{p}} \cong H_0^{n-d}(M_{\mathfrak{p}}, R_{\mathfrak{p}}) \cong H_{\mathfrak{p}R_{\mathfrak{p}}}^{n-d}(M_{\mathfrak{p}}, R_{\mathfrak{p}})$ since every Gorenstein local ring is catenary and biequidimensional, see [14, Theorem 17.3]. Moreover, $H_{\mathfrak{p}R_{\mathfrak{p}}}^{n-d}(M_{\mathfrak{p}}, R_{\mathfrak{p}}) \neq 0$ by [7, Lemma 2.4]. Hence, \mathfrak{p} is a minimal element of $\text{Supp}(H_d^{n-d}(M, R))$ and therefore $\mathfrak{p} \in \text{Ass}(H_d^{n-d}(M, R))$.

(iii) By the proof of (ii), $\text{Ass}(H_d^{n-d}(M, R)) \not\subseteq \text{Max}(R)$ so that $H_d^{n-d}(M, R)$ is not Artinian.

□

Theorem 3.3. *Let (R, \mathfrak{m}) be a local ring and M, N two finitely generated R -modules with $pd(M) < \infty$. If t is a non-negative integer such that $pd(M) < t$, then the following statements are equivalent:*

- (i) $H_d^i(M, N)$ is finitely generated, for all $i \geq t$;
- (ii) $H_d^i(M, N) = 0$, for all $i \geq t$.

Proof. (i) \Rightarrow (ii). We use induction on $n = \dim N$. When $n = 0$, we have $H_d^i(M, N) = 0$, for all $i \geq t$, see [19, Theorem 4.1]. Assume inductively, that $n > 0$ and that the result has been proved for finitely generated R -modules of dimension $n - 1$. The exact sequence

$$0 \rightarrow \Gamma_d(N) \rightarrow N \rightarrow N/\Gamma_d(N) \rightarrow 0$$

yields the long exact sequence

$$\cdots \rightarrow H_d^i(M, \Gamma_d(N)) \rightarrow H_d^i(M, N) \rightarrow H_d^i(M, N/\Gamma_d(N)) \rightarrow \cdots.$$

By [19, Lemma 2.1(3)], $H_d^i(M, \Gamma_d(N)) \cong \text{Ext}^i(M, \Gamma_d(N))$, for all $i \geq 0$. Thus we have $H_d^i(M, \Gamma_d(N)) = 0$ and $H_d^i(M, N/\Gamma_d(N)) \cong H_d^i(M, N)$, for all $i > pd(M)$. Hence, we may assume, by replacing N with $N/\Gamma_d(N)$, that N is d -torsion free module. So there exists $x \in \mathfrak{m}$ which is a non-zero divisor on N . The exact sequence

$$0 \rightarrow N \xrightarrow{x} N \rightarrow N/xN \rightarrow 0$$

induces the long exact sequence

$$\cdots \rightarrow H_d^i(M, N) \xrightarrow{x} H_d^i(M, N) \rightarrow H_d^i(M, N/xN) \rightarrow H_d^{i+1}(M, N) \rightarrow \cdots.$$

which implies that $H_d^i(M, N/xN)$ is finitely generated, for all $i \geq t$. Since N/xN is a finitely generated R -module with $\dim N/xN = n - 1$ thus by the inductive hypothesis $H_d^i(M, N/xN) = 0$, for all $i \geq t$. Therefore, $H_d^i(M, N) \cong xH_d^i(M, N)$, for all $i \geq t$. Hence, $H_d^i(M, N) = 0$, for all $i \geq t$, by Nakayma's Lemma.

(ii) \Rightarrow (i). It is clear. □

4. THE ARTINIANNES THEOREMS

Theorem 4.1. *Let M be a finitely generated R -module and N be an Artinian R -module, then $H_d^i(M, N)$ is Artinian, for all $i \geq 0$.*

Proof. We are going to argue by induction on i . In the case where $i = 0$, we see from $\Gamma_d(M, N) = \text{Hom}(M, \Gamma_d(N))$ that $H_d^0(M, N)$ is Artinian. Now suppose, inductively, that $i > 0$ and that the result has been proved for all integers less than i . Let $E(N)$ be the injective envelope of N . We have the exact sequence

$$0 \rightarrow N \rightarrow E(N) \rightarrow E(N)/N \rightarrow 0$$

which induces a long exact sequence

$$\cdots \rightarrow H_d^{i-1}(M, E(N)/N) \rightarrow H_d^i(M, N) \rightarrow H_d^i(M, E(N)) \rightarrow \cdots.$$

Since $H_d^i(M, E(N)) = 0$, for all $i > 0$, thus $H_d^{i-1}(M, E(N)/N) \cong H_d^i(M, N)$, for all $i > 1$. By hypothesis N is Artinian so $E(N)$ is Artinian thus by induction hypothesis $H_d^{i-1}(M, E(N)/N)$ is Artinian. Hence, $H_d^i(M, N)$ is Artinian. This completes the inductive step. \square

Theorem 4.2. *Let (R, \mathfrak{m}) be a local ring and M, N be two finitely generated R -modules with $r = \text{pd}(M)$ and $n = \dim(N)$. Then*

$$H_d^{r+n}(M, N) \cong \text{Ext}_R^r(M, H_d^n(N)).$$

In particular, $H_d^{r+n}(M, N)$ is an Artinian R -module.

Proof. By [16, Theorem 11.38], there is a Grothendieck spectral sequence,

$$E_2^{p,q} := \text{Ext}_R^p(M, H_d^q(N)) \Rightarrow E^{p+q} := H_d^{p+q}(M, N).$$

We have $H_d^q(N) = 0$, for all $q > n$, see [3, Lemma 2.1] and [6, Theorem 6.1.2]. Then $E_2^{p,q} = 0$, for all $p > r$ or $q > n$. We have $E_k^{r-k, n+k-1} = E_k^{r+k, n+1-k} = 0$, for all $k \geq 2$, and homomorphisms of the spectral

$$E_k^{r-k, n+k-1} \rightarrow E_k^{r,n} \rightarrow E_k^{r+k, n+1-k}$$

so $E_2^{r,n} = E_3^{r,n} = \cdots = E_\infty^{r,n}$. It is enough to prove that $E_\infty^{r,n} \cong H_d^{r+n}(M, N)$. There is a filtration

$$0 = F^{r+n+1}H^{r+n} \subseteq \cdots \subseteq F^1H^{r+n} \subseteq F^0H^{r+n} = H_d^{r+n}(M, N)$$

and

$$E_\infty^{i, r+n-i} \cong F^iH^{r+n}/F^{i+1}H^{r+n}$$

for all $0 \leq i \leq r+n$. Thus $E_2^{i, r+n-i} = \text{Ext}_R^i(M, H_d^{r+n-i}(N)) = 0$ for all $i \neq r$. Hence

$$F^{r+1}H^{r+n} = F^{r+2}H^{r+n} = \dots = F^{r+n+1}H^{r+n} = 0$$

and

$$F^r H^{r+n} = F^{r-1} H^{r+n} = \dots = F^n H^{r+n} = H_d^{r+n}(M, N).$$

This gives

$$E_\infty^{r,n} \cong F^r H^{r+n} / F^{r+1} H^{r+n} \cong H_d^{r+n}(M, N).$$

So $\text{Ext}_R^r(M, H_d^n(N)) \cong H_d^{r+n}(M, N)$. By [2, Theorem 3.1], $H_d^n(N)$ is an Artinian R -module, therefore $H_d^{r+n}(M, N)$ is also Artinian. \square

Theorem 4.3. *Let M be a finitely generated R -module and N be an R -module. Let t be a non-negative integer such that $H_d^i(N)$ is Artinian for all $i < t$. Then the following statements are true:*

- (i) $H_d^i(M, N)$ is Artinian, for all $i < t$.
- (ii) $\text{Ext}_R^i(R/\mathfrak{a}, N)$ is Artinian, for all $i < t$ and for all $\mathfrak{a} \in \Phi$.

Proof. (i) We use induction on t . When $t = 1$, $\text{Hom}(M, \Gamma_d(N))$ is Artinian so that $\Gamma_d(M, N)$ is Artinian since $\Gamma_d(M, N) = \text{Hom}(M, \Gamma_d(N))$, see [19, Lemma 2.1(1)]. Now suppose that $t > 1$ and that the result has been proved for each integer less than t . Let $E(N)$ be the injective envelope of N . Then we have the exact sequence

$$0 \rightarrow N \rightarrow E(N) \rightarrow E(N)/N \rightarrow 0.$$

Applying the functors $\Gamma_d(-)$ and $\Gamma_d(M, -)$, we get isomorphisms $H_d^i(E(N)/N) \cong H_d^{i+1}(N)$ and $H_d^i(M, E(N)/N) \cong H_d^{i+1}(M, N)$ for all $i > 0$. From the hypothesis, $H_d^i(N)$ is Artinian for all $i < t$, thus $H_d^i(E(N)/N)$ is Artinian for all $i < t-1$. By the inductive hypothesis on $E(N)/N$, $H_d^i(M, E(N)/N)$ is Artinian, for all $i < t-1$. Then by second isomorphism $H_d^i(M, N)$ is Artinian, for all $i < t$.

(ii) We use induction on t . When $t = 1$, we have the exact sequence

$$0 \rightarrow \Gamma_d(N) \rightarrow N \rightarrow N/\Gamma_d(N) \rightarrow 0$$

thus there is a long exact sequence

$$0 \rightarrow \text{Hom}(R/\mathfrak{a}, \Gamma_d(N)) \rightarrow \text{Hom}(R/\mathfrak{a}, N) \rightarrow \text{Hom}(R/\mathfrak{a}, N/\Gamma_d(N)) \rightarrow \dots$$

As $N/\Gamma_d(N)$ is Φ -torsion-free, $\text{Hom}(R/\mathfrak{a}, N/\Gamma_d(N)) = 0$ and

$$\text{Hom}(R/\mathfrak{a}, N) \cong \text{Hom}(R/\mathfrak{a}, \Gamma_d(N)).$$

Since $\Gamma_d(N)$ is Artinian R -module, then $\text{Hom}(R/\mathfrak{a}, \Gamma_d(N))$ is Artinian R -module. The proof for $t > 1$ is similar to that of (i). \square

5. ACKNOWLEDGMENT

The authors are deeply grateful to the referee for his/her careful reading of the manuscript and very helpful suggestions.

REFERENCES

- [1] C. Banica and M. Stoia, *Singular sets of a module and local cohomology modules*, Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat., **16** (1976) 923-934.
- [2] M. H. Bijan-Zadeh, *On the Artinian property of certain general local cohomology modules*, J. London Math. Soc., **32** (1985) 399-403.
- [3] M. H. Bijan-Zadeh, *A common generalization of local cohomology theories*, Glasgow Math. J., **21** (1980) 173-181.
- [4] M. P. Brodmann and A. Lashgari Faghani, *A finiteness result for associated primes of local cohomology modules*, Proc. Amer. Math. Soc., **128** (2000) 2851-2853.
- [5] M. P. Brodmann, C. Rotthaus and R. Y. Sharp, *On annihilators and associated primes of local cohomology module*, J. Pure Appl. Algebra **135** (2000) 197-227.
- [6] M. P. Brodmann and R. Y. Sharp, *Local cohomology an algebraic introduction with geometric applications*, Cambridge University Press, Cambridge, 1998.
- [7] K. Divaani Aazar, R. Sazeeleh and M. Tousi, *On vanishing of generalized local cohomology modules*, Algebra Colloq., **12** (2005) 213-218.
- [8] M. Hellus, *On the set of associated primes of a local cohomology modules*, J. Algebra, **237** (2001) 406-419.
- [9] J. Herzog, *Komplexe Suflosungen und Dualiat in der lokalen Algebra*, Universitat Regensburg, Preprint, 1974.
- [10] C. Huneke, *Problems on local cohomology modules. Free Resolution in Commutative Algebra and Algebraic Geometry*. (Sundance, UT, 1990) Res. Notes Math., 2, Jones and Bartlett, Boston, MA, (1992) 93-108.
- [11] M. Katzman, *An example of an infinite set of associated primes of local cohomology modules*, J. Algebra, **252** (2002) 161-166.
- [12] T. Marley, *Associated primes of local cohomology modules over rings of small dimension*, Manuscripta Math., **104** (2001) 519-525.
- [13] T. Marley, *The associated primes of local cohomology modules over rings of small dimension*, Manuscripta Math., **104** (2001) 519-525.
- [14] H. Matsumura, *Commutative ring theory*, Cambridge University Press, Cambridge, 1986.
- [15] Z. Rahimi, Sh. Payrovi and S. Babaei, *On the associated primes of the d -local cohomology modules*, Int. Elec. J. Alg., **25** (2019) 55-63.
- [16] J. Rotman, *Introduction to homological algebra*, Academic Press, Orlando, 1979.
- [17] A. K. Singh, *p -Tortion element in local cohomology modules*, Math. Res. Lett., **7** (2000) 165-176.

- [18] N. Suzuki, *On the generalized local cohomology and its duality*, J. Math. Kyoto Univ., **18** (1978) 71-78.
- [19] N. Zamani, M. H. Bijan-Zadeh and M. S. Sayedsadeghi, *Cohomology with supported of dimension $\leq d$* , J. Alg. Appl., **15** (2016) 1650042.

Z. Rahimi-Molaei

Department of Mathematics, Imam Khomeini International University,

Postal Code: 34149-1-6818, Qazvin, Iran.

`z.rahimi@edu.ikiu.ac.ir`

Sh. Payrovi

Department of Mathematics, Imam Khomeini International University,

Postal Code: 34149-1-6818, Qazvin, Iran.

`shpayrovi@sci.ikiu.ac.ir`

S. Babaei

Department of Mathematics, Imam Khomeini International University,

Postal Code: 34149-1-6818, Qazvin, Iran.

`sakinehbabaei@gmail.com`