

## Research Paper

### $r$ -SUBMODULES AND $uz$ -MODULES

ROSTAM MOHAMADIAN

**ABSTRACT.** In this article we study and investigate the behavior of  $r$ -submodules (a proper submodule  $N$  of an  $R$ -module  $M$  in which  $am \in N$  with  $\text{Ann}_M(a) = (0)$  implies that  $m \in N$  for each  $a \in R$  and  $m \in M$ ). We show that every simple submodule, direct summand, divisible submodule, torsion submodule and the socle of a module is an  $r$ -submodule and if  $R$  is a domain, then the singular submodule is an  $r$ -submodule. We also introduce the concepts of  $uz$ -module (i.e., an  $R$ -module  $M$  such that either  $\text{Ann}_M(a) \neq (0)$  or  $aM = M$ , for every  $a \in R$ ) and strongly  $uz$ -module (i.e., an  $R$ -module  $M$  such that  $aM \subseteq a^2M$ , for every  $a \in R$ ) in the category of modules over commutative rings. We show that every Von Neumann regular module is a strongly  $uz$ -module and every Artinian  $R$ -module is a  $uz$ -module. It is observed that if  $M$  is a faithful cyclic  $R$ -module, then  $M$  is a  $uz$ -module if and only if every its cyclic submodule is an  $r$ -submodule. In addition, in this case,  $R$  is a domain if and only if the only  $r$ -submodule of  $M$  is zero submodule. Finally, we prove that  $R$  is a  $uz$ -ring if and only if every faithful cyclic  $R$ -module is a  $uz$ -module.

---

DOI: 10.22034/as.2020.1858

MSC(2010): Primary:13A18; Secondary 13C60.

Keywords:  $r$ -ideal,  $r$ -submodule, Strongly  $uz$ -module,  $uz$ -module.

Received: 03 March 2020, Accepted: 03 August 2020.

\*Corresponding author

## 1. INTRODUCTION

Throughout this paper  $R$  is a commutative ring with  $1 \neq 0$  and  $M$  is a unitary  $R$ -module. For  $S \subseteq R$  and  $N \subseteq M$  we define  $\text{Ann}_R(S) = \{a \in R : aS = (0)\}$ ,  $\text{Ann}_M(S) = \{m \in M : mS = (0)\}$  and  $\text{Ann}_R(N) = \{a \in R : aN = (0)\}$ . For simplicity of notation, in the case  $S = \{a\}$  and  $N = \{m\}$ , we write  $\text{Ann}_R(a)$ ,  $\text{Ann}_M(a)$  and  $\text{Ann}_R(m)$  instead of  $\text{Ann}_R(\{a\})$ ,  $\text{Ann}_M(\{a\})$  and  $\text{Ann}_R(\{m\})$ , respectively. An element  $a \in R$  is said to be regular if  $\text{Ann}_R(a) = (0)$ , otherwise, it is called a zerodivisor element, and is said to be regular (resp., zerodivisor) element relative to an  $R$ -module  $M$  if  $\text{Ann}_M(a) = (0)$  (resp.,  $\text{Ann}_M(a) \neq (0)$ ). By  $r(R)$ ,  $\text{zd}(R)$  and  $u(R)$  we mean the set of all regular elements, zerodivisor elements and unit elements of  $R$ , respectively. We call a ring  $R$  a *uz*-ring if for every  $a \in R$  either  $a \in \text{zd}(R)$  or  $a \in u(R)$ . Also we denote the set of all regular elements of  $R$  relative to  $M$ , by  $r_M(R)$ , that is  $r_M(R) = \{a \in R : \text{Ann}_M(a) = (0)\}$ . An ideal  $I$  of  $R$  is called a) a nonregular ideal if  $I \subseteq \text{zd}(R)$ ; b) an  $r$ -ideal if  $ab \in I$ , with  $\text{Ann}_R(a) = (0)$ , implies that  $b \in I$ , for each  $a, b \in R$ . For  $m \in M$  (resp.,  $a \in R$ ),  $Rm$  (resp.,  $Ra$ ) denotes the cyclic submodule (resp., principal ideal) generated by  $m \in M$  (resp.,  $a \in R$ ). A homomorphism of an  $R$ -module  $M$  to itself is called an endomorphism. The set of all endomorphisms of  $M$  is a ring, which is denoted by  $\text{End}_R(M)$ . For each  $R$ -module  $M$ , the Jacobson (resp., socle), by definition, is the intersection (resp., the sum) of all maximal (resp., minimal) submodules of  $M$ , which will be denoted by  $J(M)$  (resp.,  $\text{soc}(M)$ ). An  $R$ -module  $M$  is said to be a) a simple module if it is nonzero and it has no nontrivial submodule; b) semisimple if every submodule of  $M$  is a direct summand; c) divisible if for each  $m \in M$  and  $0 \neq a \in R$ , there exists  $x \in M$  such that  $m = ax$ ; d) faithful if  $\text{Ann}_R(M) = (0)$ ; e) Von Neumann regular module if every its cyclic submodule is a direct summand. Also a nonzero submodule  $N$  of an  $R$ -module  $M$  is said to be essential if for every nonzero submodule  $K$  of  $M$  we have  $N \cap K \neq (0)$ . For more information about the aforementioned submodules in the category of  $R$ -modules, we refer the reader to [1, 6, 10, 11]. We also refer the reader to [12] and [9] for the necessary information about  $r$ -ideals and  $r$ -submodules, respectively. Finally, for more details and undefined terms and notations, see [2, 3, 5, 7].

2.  $r$ -submodules

Our aim in this section is to study the behavior of  $r$ -submodules. The concept of  $r$ -ideal was introduced and study in [12]. Recall from [9] the following definition.

**Definition 2.1.** Let  $R$  be a ring and  $M$  be an  $R$ -module. A proper submodule  $N$  of  $M$  is called an  $r$ -submodule if  $am \in N$  with  $\text{Ann}_M(a) = (0)$  implies that  $m \in N$  for each  $a \in R$  and  $m \in M$ .

Let  $R$  be any ring and let us consider  $R$  as a module over itself. Since that the submodules of  $R$  are ideals in  $R$ , one can easily show that  $I$  is an  $r$ -ideal if and only if  $I$  as a submodule is an  $r$ -submodule.

Some preliminary properties of  $r$ -submodules are as follows:

**Remark 2.2.** Let  $M$  be an  $R$ -module.

- (i) The zero submodule of  $M$  is an  $r$ -submodule.
- (ii) The intersection of any family of  $r$ -submodules of  $M$  is an  $r$ -submodule.
- (iii)  $\text{Ann}_M(I)$  is an  $r$ -submodule of  $M$  for any ideal  $I$  of  $R$ .
- (iv) If  $f \in \text{End}_R(M)$ , then  $\ker(f) = \{m \in M : f(m) = 0\}$  is an  $r$ -submodule of  $M$ .

Let  $M$  be an  $R$ -module. Recall that if  $N$  is a submodule of  $M$  and  $a \in R$ , then  $(N : a) = \{m \in M : am \in N\}$  is a submodule of  $M$  which contains  $N$ . Also if  $S$  is a multiplicatively closed subset of  $R$ , then  $S^{-1}R$  (resp.,  $S^{-1}M$ ) is a ring (resp., an  $S^{-1}R$ -module), which is called the ring (resp., module) of fractions of  $R$  (resp.,  $M$ ) with respect to  $S$ . Clearly,  $S = r_M(R)$  is a multiplicatively closed subset in  $R$ . For more information about the above concepts, see [13]. In the following proposition we give several equivalent definitions for  $r$ -submodules. For the proof see Proposition 4 in [9].

**Proposition 2.3.** Let  $M$  be an  $R$ -module and  $N$  be a submodule of  $M$ . Then the following statements are equivalent.

- (i)  $N$  is an  $r$ -submodule.
- (ii)  $aM \cap N = aN$ , for each  $a \in r_M(R)$ .
- (iii)  $N = (N : a)$ , for each  $a \in r_M(R)$ .
- (iv)  $N = \mathcal{N}^c$ , where  $\mathcal{N}$  is a submodule in  $S^{-1}M$  and  $S = r_M(R)$ .

**Proposition 2.4.** Let  $N \subseteq K$  be two submodules of an  $R$ -module  $M$ . If  $N$  is an  $r$ -submodule of  $M$  and  $\frac{K}{N}$  is an  $r$ -submodule of  $R$ -module  $\frac{M}{N}$ , then  $K$  is an  $r$ -submodule of  $M$ .

*Proof.* Let  $a \in R$ ,  $m \in M$  and  $am \in K$  with  $\text{Ann}_M(a) = (0)$ . Clearly,  $a(m + N) \in \frac{K}{N}$  and also  $\text{Ann}_{\frac{M}{N}}(a) = (0)$ . To see this, let  $m + N \in \text{Ann}_{\frac{M}{N}}(a)$ . Hence  $a(m + N) = am + N = N$  which implies that  $am \in N$ . On the other hand, since  $\text{Ann}_M(a) = (0)$  and  $N$  is an  $r$ -submodule, we have  $m \in N$  whence  $m + N = N$ . Therefore by our hypothesis, we have  $m + N \in \frac{K}{N}$  and so  $m \in K$ . This shows that  $K$  is an  $r$ -submodule.  $\square$

If  $f : M \rightarrow N$  is an  $R$ -module isomorphism, then it is clear that  $\text{Ann}_M(a) = (0)$  if and only if  $\text{Ann}_N(a) = (0)$ , for any  $a \in R$ .

**Proposition 2.5.**  *$r$ -submodules are invariant under isomorphisms.*

*Proof.* Let  $M$  and  $N$  be  $R$ -modules and  $f : M \rightarrow N$  be an  $R$ -module isomorphism. We are to show that whenever  $K$  is an  $r$ -submodule of  $M$ , then  $f(K)$  is an  $r$ -submodule of  $N$ . To see this, suppose that  $a \in R$ ,  $n \in N$  and  $an \in f(K)$  with  $\text{Ann}_N(a) = (0)$ . Take  $m \in K$  and  $m_1 \in M$  such that  $an = f(m)$  and  $n = f(m_1)$ . Clearly,  $an = af(m_1) = f(am_1) = f(m)$ , whence  $f(am_1 - m) = 0$  and so  $am_1 - m \in \ker(f) = (0)$ . Therefore  $am_1 = m \in K$ . Since  $\text{Ann}_M(a) = (0)$ , we infer that  $m_1 \in K$  and hence  $n = f(m_1) \in f(K)$ .  $\square$

In the following two theorems we observe that every simple submodule and the socle of a module are  $r$ -submodules.

**Theorem 2.6.** *Every simple submodule of a module is an  $r$ -submodule.*

*Proof.* Assume that  $N$  is a simple submodule of an  $R$ -module  $M$ . Therefore there exists  $0 \neq m \in N$  such that  $N = Rm$ . Now let  $a \in R$ ,  $x \in M$  and  $ax \in N$  with  $\text{Ann}_M(a) = (0)$ . If  $ax = 0$ , then  $x = 0 \in N$ . In case  $ax \neq 0$ , we have  $N = Rax$ . Since  $am \neq 0$ , we infer that  $N = Ram$ . Consequently,  $N = Rax = Ram$  and hence  $ax \in Ram$ . Therefore there exists  $s \in R$  such that  $ax = sam$ , whence  $x - sm \in \text{Ann}_M(a) = (0)$ . Thus  $x = sm \in Rm = N$  which completes the proof.  $\square$

The following corollary is now an immediate consequence of Theorem 2.6.

**Corollary 2.7.** *If  $M$  is a very semisimple  $R$ -module (i.e., its every cyclic submodule is simple), then every cyclic submodule of  $M$  is an  $r$ -submodule.*

**Theorem 2.8.** *Let  $M$  be any  $R$ -module. Then  $\text{soc}(M)$  is an  $r$ -submodule.*

*Proof.* Suppose that  $\{N_i : i \in A\}$  be the set of all minimal submodules of  $M$ . By definition, we have  $\text{soc}(M) = \bigoplus_{i \in A} N_i$ . Now let  $a \in R$ ,  $m \in M$  and  $am \in \text{soc}(M)$  with  $\text{Ann}_M(a) = (0)$ . Hence  $am = \sum_{k=1}^n a_{i_k}$ , where  $a_{i_k} \in N_{i_k}$ , for  $i_1, \dots, i_n \in A$ . Without loss of generality, we can assume that  $aa_{i_k} \neq 0$ , for each  $i_k$ . Consequently,  $Raa_{i_k} = N_{i_k}$  therefore  $am = \sum_{k=1}^n ar_{i_k}a_{i_k}$ , where  $r_{i_k} \in R$ , for  $k = 1, \dots, n$ . This implies that  $m - \sum_{k=1}^n r_{i_k}a_{i_k} \in \text{Ann}_M(a) = (0)$ , hence  $m = \sum_{k=1}^n r_{i_k}a_{i_k}$ . Therefore  $m \in \text{soc}(M)$  and we are done.  $\square$

**Proposition 2.9.** (i) *Every divisible submodule of a module is an  $r$ -submodule.*

(ii) *Every direct summand of a module is an  $r$ -submodule.*

*Proof.* (i) Assume that  $N$  is a divisible submodule of an  $R$ -module  $M$ . Let  $a \in R$ ,  $m \in M$  and  $am \in N$  with  $\text{Ann}_M(a) = (0)$ . Since  $N$  is divisible, there exists  $n \in N$  such that  $am = an$ . Hence  $m - n \in \text{Ann}_M(a) = (0)$  and therefore  $m = n \in N$ .

(ii) Suppose that  $N$  is a direct summand of an  $R$ -module  $M$ . By Lemma 5.6 in [1], there exists  $e \in \text{End}_R(M)$  such that  $N = \ker(1 - e)$  and  $e^2 = e$ . This means that  $N$  is an  $r$ -submodule.  $\square$

**Example 2.10.** In view of Proposition 2.9, injective submodules in any  $R$ -module, a fortiori  $\mathbb{Z}(p^\infty)$  as a  $\mathbb{Z}$ -submodule of  $\frac{\mathbb{Q}}{\mathbb{Z}}$  are  $r$ -submodules.

Using the previous proposition we have the next corollary.

**Corollary 2.11.** (i) *If  $M$  is a semisimple  $R$ -module, then every submodule of  $M$  is an  $r$ -submodule.*

(ii) *If  $M$  is a Von Neumann regular  $R$ -module, then every finitely generated submodule of  $M$  is a direct summand and therefore it is an  $r$ -submodule as well, see Lemma 1 in [8].*

We recall that if  $M$  is an  $R$ -module, then  $t(M) = \{m \in M : \text{Ann}_R(m) \neq (0)\}$  is called torsion submodule of  $M$ . If  $t(M) = M$  (resp.,  $t(M) = (0)$ ), then  $M$  is called torsion (resp., torsion free) module. We also recall that  $\mathcal{Z}(M) = \{m \in M : \text{Ann}_R(m) \text{ is an essential ideal in } R\}$  is a submodule of  $M$ , which is called singular submodule. If  $\mathcal{Z}(M) = M$ , (resp.,  $\mathcal{Z}(M) = (0)$ ) then  $M$  is called singular (resp., nonsingular) module. In the following results we show that  $t(M)$  is always an  $r$ -submodule of  $M$  and if  $R$  is a domain, then  $\mathcal{Z}(M)$  is also an  $r$ -submodule of  $M$ .

**Proposition 2.12.** *Let  $M$  be an  $R$ -module. Then the following statements hold.*

(i) *The torsion submodule of  $M$  is an  $r$ -submodule.*

(ii) *If  $R$  is a domain then the singular submodule of  $M$  is an  $r$ -submodule.*

*Proof.* (i) Suppose that  $am \in t(M)$  with  $\text{Ann}_M(a) = (0)$ , where  $a \in R$  and  $m \in M$ . By definition of the torsion submodule, we have  $\text{Ann}_R(am) \neq (0)$  whence there exists  $0 \neq s \in R$  such that  $s(am) = a(sm) = 0$ . Therefore  $sm \in \text{Ann}_M(a) = (0)$  and hence  $0 \neq s \in \text{Ann}_R(m)$ , that is,  $\text{Ann}_R(m) \neq (0)$ . This means that  $m \in t(M)$ .

(ii) Assume that  $am \in \mathcal{Z}(M)$  with  $\text{Ann}_M(a) = (0)$ , where  $a \in R$  and  $m \in M$ . The definition of singular submodule implies that  $\text{Ann}_R(am) \cap Rx \neq (0)$ , for any  $0 \neq x \in R$ . Hence there exists  $0 \neq s \in R$  such that  $(sxm)a = 0$ , and thus  $sxm \in \text{Ann}_M(a) = (0)$ , so  $0 \neq sx \in \text{Ann}_R(m) \cap Rx$ , i.e.,  $m \in \mathcal{Z}(M)$ .  $\square$

The direct sum of two  $r$ -submodules may not be an  $r$ -submodule, see Example 5.14 in [12]. For a nontrivial idempotent  $e$  in  $R$ ,  $eM$  is clearly an  $r$ -submodule of  $M$ , for manifestly  $eM$  is a summand of  $M$ . Now the following proposition shows certain direct sum of  $r$ -submodules in a module  $M$ , which are not necessarily simple submodules is in fact an  $r$ -submodule.

**Proposition 2.13.** *Let  $M$  be an  $R$ -module and  $\{e_i : i \in A\}$  be a set of orthogonal idempotents in  $R$  and no finite subset of these idempotents generate  $R$ , in the sense that  $1 \neq \sum_{i \in B} e_i$ , where  $B$  is a finite subset of  $A$ . Then  $N = \oplus_{i \in A} e_i M$  is an  $r$ -submodule.*

*Proof.* Let  $am \in N$ , where  $a \in R$ ,  $m \in M$  with  $\text{Ann}_M(a) = (0)$ . We are to show that  $m \in N$ . Clearly,  $am = \sum_{k=1}^n e_{i_k} m_{i_k}$ , where  $i_k \in A$  and  $m_{i_k} \in M$ , for  $k = 1, \dots, n$ . Let us put  $x = \prod_{k=1}^n (1 - e_{i_k})$ . It is manifest that  $amx = 0$  and hence  $mx = 0$ . It is now evident that  $x = 1 - y$ , where  $y = \sum_{k=1}^n e_{i_k}$ . Therefore  $m(1 - y) = 0$ , so  $m = my$ . This implies that  $m \in N$ .  $\square$

**Definition 2.14.** Let  $R$  be a ring and  $M$  be an  $R$ -module. Then

- (i)  $a \in R$  is said to be  $m$ -regular relative to  $M$ , if  $\text{Ann}_R(a) = (0)$  implies that  $\text{Ann}_M(a) = (0)$ ;
- (ii)  $a \in R$  is said to be  $R$ -regular relative to  $M$ , if  $\text{Ann}_M(a) = (0)$  implies that  $\text{Ann}_R(a) = (0)$ .

For example if we consider  $R[x]$  as a module over  $R$ , then every  $a \in R$  is an  $m$ -regular element relative to  $R[x]$  if and only if it is an  $R$ -regular element relative to  $R[x]$ . Also one can easily see that, if  $M$  is an  $R$ -module and  $\text{Ann}_M(a) = (0)$ , for every  $a \in R$ , then  $\text{Ann}_R(M)$  is an  $r$ -ideal in  $R$ . Note that, in this case, there is no any essential  $r$ -submodule in  $M$ .

**Lemma 2.15.** *Suppose that  $M$  is an  $R$ -module. Then the following statements hold.*

- (i) *If  $M$  is a faithful  $R$ -module, then every  $a \in R$  is an  $R$ -regular element relative to  $M$ .*
- (ii) *If  $M$  is a finitely generated free  $R$ -module, then every  $a \in R$  is an  $m$ -regular element relative to  $M$ .*

*Proof.* (i) Assume that  $a \in R$  with  $\text{Ann}_M(a) = (0)$  and  $s \in \text{Ann}_R(a)$ . Hence  $sa = 0$  and it is evident that  $sam = 0$ , for any  $m \in M$ . Thus  $sm \in \text{Ann}_M(a) = (0)$  and therefore  $sm = 0$ . This implies that  $s \in \text{Ann}_R(M) = (0)$ , i.e.,  $\text{Ann}_R(a) = (0)$ .

(ii) Assume that  $X = \{x_1, \dots, x_n\}$  is a base for  $M$ ,  $a \in R$  with  $\text{Ann}_R(a) = (0)$  and  $m \in M$ . Now suppose that  $m \in \text{Ann}_M(a)$ , hence  $am = 0$ . On the other hand, there exist  $s_1, \dots, s_n \in R$  such that  $m = s_1 x_1 + \dots + s_n x_n$ . Therefore  $as_1 x_1 + \dots + as_n x_n = 0$ , and consequently  $as_i = 0$ , for  $i = 1, \dots, n$ . This conclude that  $s_1, \dots, s_n \in \text{Ann}_R(a) = (0)$ , therefore  $s_i = 0$ , for  $i = 1, \dots, n$  and hence  $m = 0$ . This implies that  $\text{Ann}_M(a) = (0)$ .  $\square$

We should emphasize that any cyclic submodule need not be an  $r$ -submodule. For example, the principle ideal  $I = \mathbb{Z}4$  in  $\mathbb{Z}$  is not an  $r$ -ideal and so it is not an  $r$ -submodule of  $\mathbb{Z}$  as a  $\mathbb{Z}$ -module. Whenever  $M$  is a finitely generated free  $R$ -module and  $I$  is an  $r$ -ideal in  $R$ , we have the following fact.

**Proposition 2.16.** *Let  $M$  be a finitely generated free  $R$ -module with a base  $X$  and  $I$  be an ideal in  $R$ . Then  $I$  is an  $r$ -ideal in  $R$  if and only if  $IX$  is an  $r$ -submodule of  $M$ .*

*Proof.* Suppose that  $X = \{x_1, \dots, x_n\}$  and  $am \in IX$  with  $\text{Ann}_M(a) = (0)$ , where  $a \in R$  and  $m \in M$ . Take  $s_1, \dots, s_n \in R$  and  $t_1, \dots, t_n \in I$  such that  $m = s_1x_1 + \dots + s_nx_n$  and  $am = t_1x_1 + \dots + t_nx_n$ . Hence  $as_1x_1 + \dots + as_nx_n = t_1x_1 + \dots + t_nx_n$ . Therefore  $as_i = t_i \in I$ , for  $i = 1, \dots, n$ . Now by part (i) of the above lemma we have  $\text{Ann}_R(a) = (0)$  and so by our hypothesis, we conclude that  $s_i \in I$ , for  $i = 1, \dots, n$ . This means that  $m \in IX$ . Conversely, suppose that  $ax \in I$ , with  $\text{Ann}_R(a) = (0)$ , where  $a, x \in R$  and  $0 \neq m \in M$ . Clearly,  $axm \in IX$ . Now using part (ii) of the above lemma, we have  $\text{Ann}_M(a) = (0)$ , whence by our hypothesis, we have  $xm \in IX$ . This yields that  $x \in I$ .  $\square$

We remind the reader that a submodule  $N$  of a module  $M$  is called prime (resp., primary) if for each  $a \in R$  and  $m \in M$ ,  $am \in N$  implies that  $m \in N$  or  $aM \subseteq N$  (resp.,  $a^nM \subseteq N$  for some  $n \in \mathbb{N}$ ). Also  $N$  is called semiprime, if  $a^2m \in N$  implies that  $am \in N$ . Clearly, every submodule is prime if and only if it is both primary and semiprime. Furthermore, if  $N$  is a prime  $r$ -submodule of  $M$ , then  $am \in N$  implies that  $m \in N$ , for every  $m \in M$  and  $a \in r_M(R)$ . For otherwise, we have  $aM \subseteq N$  and so by part (ii) of Proposition 2.3 we conclude that  $aM = aN$ . This immediately implies that  $M = N$  which is not true.

Now similarly to the notion of nonregular ideal, we may define a nonregular submodule.

**Definition 2.17.** A proper submodule  $N$  of an  $R$ -module  $M$  is called nonregular, if  $aM \subseteq N$  implies that  $\text{Ann}_M(a) \neq (0)$ , for each  $a \in R$ .

If we consider  $R$  as an  $R$ -module, then our definition agrees with the concept of nonregular ideal.

**Remark 2.18.** (i) Every  $r$ -submodule of a module is nonregular.

(ii) Every prime nonregular submodule of a module is an  $r$ -submodule.

The converse of part (i) of the above remark is not true, in general. For example, consider  $\mathbb{Q}$  as a  $\mathbb{Z}$ -module. Then the submodule  $N = \mathbb{Z}\frac{1}{2}$  is a nonregular submodule but it is not an  $r$ -submodule. To see this, it is clear that  $2 \cdot \frac{3}{4} = \frac{1}{2} \cdot 3 \in N$  and  $\text{Ann}_{\mathbb{Q}}(2) = (0)$  but  $\frac{3}{4} \notin N$ .

We conclude this section with the following proposition.

**Proposition 2.19.** *Every maximal  $r$ -submodule is a prime submodule.*

*Proof.* Assume that  $N$  is a maximal  $r$ -submodule of an  $R$ -module  $M$ . We are to show that  $N$  is prime. To see this, let  $a \in R$ ,  $m \in M$  and  $am \in N$ . Since  $N$  is an  $r$ -submodule,  $(N : a)$  is an  $r$ -submodule and it is evident that  $N \subseteq (N : a)$ . Now maximality of  $N$  implies that  $(N : a) = N$  and hence we have  $m \in N$ , i.e.,  $N$  is prime.  $\square$

### 3. $uz$ -modules

This section is devoted to the introduction of the  $uz$ -modules and strongly  $uz$ -modules. We begin with the following definitions.

**Definition 3.1.** An  $R$ -module  $M$  is called a

- (i)  $uz$ -module, if for every  $a \in R$  either  $\text{Ann}_M(a) \neq (0)$  or  $aM = M$ ;
- (ii) strongly  $uz$ -module if for every  $a \in R$  we have  $aM \subseteq a^2M$  (in fact,  $aM = a^2M$ ).

For instance, the modules  $\frac{\mathbb{Q}}{\mathbb{Z}}$ ,  $\mathbb{Q}$  and  $\mathbb{Z}(p^\infty)$  over  $\mathbb{Z}$  are strongly  $uz$ -modules but  $\mathbb{Z}$  as  $\mathbb{Z}$ -module is not a strongly  $uz$ -module. Every strongly  $uz$ -module is a  $uz$ -module, but the converse is not true, in general. For example,  $\mathbb{Z}_4$  as a  $\mathbb{Z}_4$ -module is a  $uz$ -module, but is not a strongly  $uz$ -module. The ring of  $C(X)$ , i.e., the ring of all real-valued continuous functions on a completely regular Hausdorff space  $X$  is a strongly  $uz$ -module as a  $C(X)$ -module. Recall that it is possible that  $\text{Ann}_M(a) = (0)$  or  $aM = M$  for every  $0 \neq a \in R$ . For example, if we consider  $\mathbb{Q}$  as a  $\mathbb{Z}$ -module then both  $\text{Ann}_{\mathbb{Q}}(a) = (0)$  and  $a\mathbb{Q} = \mathbb{Q}$  for every  $0 \neq a \in \mathbb{Z}$ .

It is clear that a) every simple module is a strongly  $uz$ -module; b) a ring  $R$  is a  $uz$ -ring (resp., Von Neumann regular ring) if and only if as a module over itself is a  $uz$ -module (resp., strongly  $uz$ -module); c) if  $M$  is a strongly  $uz$ -module, then every primary submodule of  $M$  is prime and  $\text{Ann}_R(M)$  is a semiprime ideal.

**Remark 3.2.** Let  $M$  be an  $R$ -module. Then the following statements hold.

- (i) If  $a \in R$ , then  $\text{Ann}_M(a) = (0)$  if and only if  $\text{Ann}_M(a^n) = (0)$ , for any  $n \in \mathbb{N}$ .
- (ii) The zero submodule of  $M$  is prime if and only if  $\text{Ann}_M(a) = (0)$ , for any  $0 \neq a \in R$ .

**Remark 3.3.** Let  $M$  be a faithful  $R$ -module. Then the following statements hold.

- (i) If  $M$  is a strongly  $uz$ -module, then  $R$  is a reduced ring. In particular, every Von Neumann regular ring is reduced.
- (ii) If  $M$  is an Artinian module and the zero submodule of  $M$  is prime, then  $aM = M$ , for any  $0 \neq a \in R$ . In this case, clearly  $M$  is a strongly  $uz$ -module. As a consequence we have the well known fact that every Artinian domain is a field.

The next result states that every Von Neumann regular (resp., Artinian) module is a strongly  $uz$ -module (resp.,  $uz$ -module).

**Theorem 3.4.** (i) *Every Von Neumann regular  $R$ -module is a strongly  $uz$ -module.*  
(ii) *Every Artinian  $R$ -module is a  $uz$ -module.*

*Proof.* (i) Assume that  $M$  is a Von Neumann regular  $R$ -module and  $a \in R$ . We must show that  $aM \subseteq a^2M$ . Let  $m \in M$ , it is sufficient to show that  $am \in a^2M$ . Put  $N = Ram$ . Clearly,  $N$  is a submodule of  $M$  and hence it is a direct summand. Thus there exists a submodule  $K$  of  $M$  such that  $M = N \oplus K$ . Hence there exist  $r \in R$  and  $x \in K$  such that  $m = ram + x$ . Consequently,  $ax = (1 - ra)am \in N \cap K = (0)$ . Therefore  $am = ra^2m \in a^2M$ .

(ii) If  $\text{Ann}_M(a) \neq (0)$ , for any  $a \in R$ , then we are done. Hence suppose that there exists  $a_0 \in R$  such that  $\text{Ann}_M(a_0) = (0)$ . Since  $a_0M \supseteq a_0^2M \supseteq a_0^3M \supseteq \dots$ , it follows that there exists  $n_0 \in \mathbb{N}$  such that  $a_0^nM = a_0^{n+1}M$ , for any  $n \geq n_0$ . Now take an arbitrary  $m \in M$ . Hence there exists  $x \in M$  such that  $a_0^{n_0}m = a_0^{n_0+1}x$ . Therefore  $a_0^{n_0}(m - a_0x) = 0$  and so  $m - a_0x \in \text{Ann}_M(a_0^{n_0}) = (0)$ . Thus  $m = a_0x \in a_0M$ , i.e.,  $M = a_0M$ , which completes the proof.  $\square$

Part (i) of the previous theorem conclude that every semisimple module is a strongly  $uz$ -module. Also the converse of parts (i) and (ii) is not true, in general. For example  $\mathbb{Z}(p^\infty)$  as  $\mathbb{Z}$ -module is a strongly  $uz$ -module but is not a Von Neumann regular  $\mathbb{Z}$ -module and  $\mathbb{Q}$  as  $\mathbb{Z}$ -module is a strongly  $uz$ -module but is not a Artinian  $\mathbb{Z}$ -module

Recall that an  $R$ -module  $M$  is called multiplication module if for each submodule  $N$  of  $M$ ,  $N = IM$  for some ideal  $I$  of  $R$ . In view of Proposition 2.16, it is easy to show that if  $M$  is a cyclic free multiplication  $R$ -module, then  $R$  is a  $uz$ -ring if and only if every submodule of  $M$  is an  $r$ -submodule.

**Proposition 3.5.** *Let  $M$  be a strongly  $uz$ -module. Then the following statements hold.*

- (i) *Every primary submodule of  $M$  is prime.*
- (ii) *Every semiprime submodule of  $M$  is an  $r$ -submodule.*
- (iii) *If  $N$  is a semiprime submodule of  $M$  and  $am \in N$ , where  $a \in R$  and  $m \in M$ , then either  $m \in N$  or  $\text{Ann}_M(a) \neq (0)$ .*

*Proof.* (i) It is evident.

(ii) Let  $am \in N$ , where  $a \in R$ ,  $m \in M$  with  $\text{Ann}_M(a) = (0)$ . Since  $aM = a^2M$ , there exists  $x \in M$  such that  $am = a^2x \in N$ . Consequently,  $m - ax \in \text{Ann}_M(a) = (0)$ , implies  $m = ax$ . On the other hand, since  $N$  is semiprime, we have  $ax \in N$  and consequently,  $m \in N$ .

(iii) It is evident.  $\square$

An infinite  $R$ -module  $M$  is called Jónsson module if every proper submodule of  $M$  has smaller cardinality than  $M$ . Clearly, every simple module is a Jónsson module. It is well known that if  $M$  is a Jónsson module, then either  $aM = M$  or  $aM = (0)$ , for each  $a \in R$ , and moreover  $\text{Ann}_R(M)$  is a prime ideal of  $R$ , see Proposition 2.5 in [4]. One can easily show that every Jónsson module is a strongly  $uz$ -module. For more details about Jónsson modules, see [4].

In the following result, we observe that for any faithful cyclic  $R$ -module  $M$ , every submodule of  $M$  is an  $r$ -submodule if and only if  $M$  is a  $uz$ -module.

**Proposition 3.6.** *Let  $M$  be a faithful cyclic  $R$ -module. Then the following conditions are equivalent.*

- (i)  $M$  is a  $uz$ -module.
- (ii) Every submodule of  $M$  is an  $r$ -submodule.
- (iii) Every cyclic submodule of  $M$  is an  $r$ -submodule.

*Proof.* ( $i \Rightarrow ii$ ) Suppose that  $0 \neq m \in M$  and  $M = Rm$ . Let  $N$  be a submodule of  $M$ ,  $a \in R$ ,  $x \in M$  and  $ax \in N$  with  $\text{Ann}_M(a) = (0)$ . By our hypothesis, we have  $aM = M$ , that is,  $Ram = Rm$ . Hence there exists  $s \in R$  such that  $m = asm$ . Therefore  $(1 - as) \in \text{Ann}_R(m) = (0)$ , so  $1 = as$ . Thus we conclude that  $x = s(ax) \in N$ , i.e.,  $N$  is an  $r$ -submodule.

( $ii \Rightarrow i$ ) If  $aM = M$ , for any  $a \in R$ , then we are done. Otherwise, let there exists  $a_0 \in R$  such that  $a_0M \neq M$ . We are to show that  $\text{Ann}_M(a_0) \neq (0)$ . Since  $a_0M \neq M$ , there exists  $m_0 \in M$  such that  $m_0 \notin a_0M$ . By taking that  $N = a_0M$ , we have  $a_0m_0 \in N$  and  $m_0 \notin N$ . Now if  $\text{Ann}_M(a_0) = (0)$ , then  $N$  is not an  $r$ -submodule, which is a contradiction.

( $ii \Rightarrow iii$ ) It is evident.

( $iii \Rightarrow ii$ ) Let  $N$  be a submodule of  $M$ ,  $a \in R$ ,  $x \in M$  and  $ax \in N$  with  $\text{Ann}_M(a) = (0)$ . Now by our hypothesis, the submodule  $K = Rax$  is an  $r$ -submodule and it is obvious that  $K \subseteq N$ . Clearly,  $x \in K$  and so  $x \in N$ , that is,  $N$  is an  $r$ -submodule.  $\square$

The condition of “ $M$  being a cyclic  $R$ -module” for implication ( $i \Rightarrow ii$ ) is essential, i.e., ( $i \Rightarrow ii$ ) is not true, in general. For example,  $\mathbb{Q}$  as  $\mathbb{Z}$ -module is a  $uz$ -module, but the submodule  $N = \mathbb{Z}_2^1$  of  $\mathbb{Q}$  is not an  $r$ -submodule. It is worth to point out that the implication ( $ii \Rightarrow i$ ) is valid for every  $R$ -module. Furthermore, if in the above proposition we consider  $R$  as an  $R$ -module, then Proposition 3.4 in [12] is an immediate consequence of Proposition 3.6.

**Remark 3.7.** (i) A ring  $R$  is a domain if it has a  $uz$ -module whose every nonzero submodule is faithful. To see this, let  $M$  be an  $R$ -module which is a  $uz$ -module with the property mentioned above. Hence, in view of Definition 3.1, we must have  $aM = M$ , for all

$a \in R$  (note, by our assumption we can not have  $\text{Ann}_M(a) \neq (0)$ ). This immediately implies that  $R$  is a domain.

(ii) A ring  $R$  is a field if and only if it has a  $uz$ -cyclic module whose every nonzero submodule is faithful. To see this, if  $M$  is cyclic which is a  $uz$ -module with the property mentioned above, it is evident that it is isomorphic to  $R$  itself, i.e.,  $R$  becomes a domain which is at the same time a  $uz$ -module over itself (note, the property of being a  $uz$ -module or a strongly  $uz$ -module is preserved under isomorphism). Consequently,  $aR = R$  for all  $a \in R$  which is the same thing as  $R$  being a field, and we are done. The converse is evident, for it is sufficient to consider  $R$  as an  $R$ -module.

It is well known that a ring  $R$  is a domain if and only if the only  $r$ -ideal of  $R$  is zero ideal, see Proposition 2.8 in [12]. By replacing the  $r$ -ideals with the  $r$ -submodules, we get the next interesting fact.

**Proposition 3.8.** *Let  $M$  be a faithful cyclic  $R$ -module. Then the following statements are equivalent.*

- (i)  $R$  is a domain.
- (ii) The only  $r$ -submodule of  $M$  is zero submodule.
- (iii)  $\text{Ann}_M(ab) = \text{Ann}_M(a) \cup \text{Ann}_M(b)$ , for every  $a, b \in R$ .

*Proof.* ( $i \Rightarrow ii$ ) Assume that  $0 \neq m \in M$  and  $M = Rm$ . Let  $(0) \neq N$  be an  $r$ -submodule of  $M$  and  $0 \neq n \in N$ . Hence there exists  $0 \neq a \in R$  such that  $n = am$ . We claim that  $\text{Ann}_M(a) = (0)$ . To see this, let  $0 \neq x \in M$  such that  $ax = 0$ . Thus there exists  $0 \neq b \in R$  such that  $x = bm$ . Therefore we have  $abm = 0$ , so  $ab \in \text{Ann}_R(m) = (0)$ . This implies that  $ab = 0$ , which is not true, for  $R$  is a domain. Now since  $N$  is an  $r$ -submodule, we infer that  $m \in N$ , that is,  $M = N$ , which is a contradiction.

( $ii \Rightarrow iii$ ) Since  $\text{Ann}_M(a)$  is an  $r$ -submodule, for every  $a \in R$ , the proof is evident.

( $iii \Rightarrow i$ ) Suppose that  $a, b \in R$  and  $ab = 0$ . Hence  $M = \text{Ann}_M(0) = \text{Ann}_M(ab) = \text{Ann}_M(a) \cup \text{Ann}_M(b)$ . Therefore  $M = \text{Ann}_M(a)$  or  $M = \text{Ann}_M(b)$ . That is  $aM = (0)$  or  $bM = (0)$ . Thus  $a \in \text{Ann}(M) = (0)$  or  $b \in \text{Ann}(M) = (0)$ . This means that  $a = 0$  or  $b = 0$ .  $\square$

Propositions 2.12 and 3.8 state that every faithful cyclic module over a domain is both nonsingular and torsion free.

In the following theorem we observe two equivalent conditions for  $uz$ -rings. For the other equivalent conditions in terms of  $r$ -ideals, see Proposition 3.4 in [12].

**Theorem 3.9.** *Let  $R$  be a ring. Then the following statements are equivalent.*

- (i)  $R$  is a  $uz$ -ring.
- (ii) Every faithful  $R$ -module is a  $uz$ -module.
- (iii) Every faithful cyclic  $R$ -module is a  $uz$ -module.

*Proof.* ( $i \Rightarrow ii$ ) Let  $M$  be a faithful  $R$ -module. In view of Proposition 3.6, it is enough to show that every submodule of  $M$  is an  $r$ -submodule. Suppose that  $N$  is a submodule of  $M$  and  $am \in N$  with  $\text{Ann}_M(a) = (0)$ , where  $a \in R$  and  $m \in M$ . In case  $a \in u(R)$ , we have  $m = a^{-1}am \in N$ . If  $\text{Ann}_R(a) \neq (0)$ , then there exists  $0 \neq b \in R$  such that  $ab = 0$ . Now assume that  $x \in M$  is an arbitrary element. Clearly,  $abx = 0$ , whence  $bx \in \text{Ann}_M(a) = (0)$ . Hence  $bx = 0$ , i.e.,  $0 \neq b \in \text{Ann}_R(M) = (0)$ , which is a contradiction.

( $ii \Rightarrow iii$ ) It is evident.

( $iii \Rightarrow i$ ) Let  $M = Rm$  be a faithful cyclic  $R$ -module and  $a \in R$ . If  $\text{Ann}_M(a) \neq (0)$ , then there exists  $0 \neq n \in M$  such that  $an = 0$ . On the other hand, there exists  $0 \neq b \in R$  such that  $n = bm$ , hence  $abm = 0$ , whence  $ab \in \text{Ann}_R(m) = (0)$ , that is  $ab = 0$ . This means that  $a \in \text{zd}(R)$ . If  $aM = M$ , then there exists  $m_1 \in M$  such that  $m = am_1$ . Also there is  $t \in R$  such that  $m_1 = mt$ . Therefore  $m = amt$ , implies  $(1 - at)m = 0$ , hence  $1 - at \in \text{Ann}_R(m) = (0)$ , that is,  $at = 1$ . This implies that  $a \in u(R)$ .  $\square$

We conclude the paper by the following corollary which introduces some  $r$ -submodules of a  $uz$ -module.

**Corollary 3.10.** *Let  $R$ -module  $M$  be a  $uz$ -module. Then the following statements hold.*

- (i) Every prime submodule of  $M$  is nonregular, and hence it is an  $r$ -submodule.
- (ii)  $J(M)$  is an  $r$ -submodule of  $M$ .

#### 4. ACKNOWLEDGMENTS

The author would like to thank the referee for reading the article carefully and giving useful comments. Also the author acknowledges the financial support (GN:SCU.MM98.648) from the Research Council of Shahid Chamran University of Ahvaz.

#### REFERENCES

- [1] F. W. Anderson, K. R. Fuller, *Rings and Categories of Modules*, Graduate Texts in Mathematics, Vol. 13, 2nd Ed., Springer-Verlag, Berlin/Heidelberg, New York, 1992.
- [2] M. F. Atiyah, I. G. Macdonald, *Introduction to Commutative Algebra*, Addison-Wesely, Reading Mass, 1969.
- [3] R. Gilmer, *Multiplicative Ideal Theory*, Marcel Dekker Inc, New York, 1972.
- [4] R. Gilmer, W. Heinzer, *On Jónsson modules over a commutative ring*, Acta Sci. Math., **46** (1983) 3-15.
- [5] J. A. Huckabo, *Commutative Ring with Zero Divisors*, Marcel Dekker Inc, 1988.

- [6] T. W. Hungerford, *Algebra*, Graduate Texts in Mathematics, Vol. 73, Springer-Verlag, Berlin/Heidelberg, New York, 1989.
- [7] I. Kaplansky, *Commutative Rings*, Allyn and Bacon, Boston, 1970.
- [8] O. A. S. Karamzadeh, *Module whose countably generated submodules are epimorphic images*, Colloq. Math., **46** (1982) 143-146.
- [9] S. Koc, U. Tekir, *r-Submodules and special r-Submodules*, Turk. J. Math., **42** (2018) 1863-1876.
- [10] T. Y. Lam, *Lecture on Modules and Rings*, Springer, 1999.
- [11] J. Lambek, *Lecture on Rings and Modules*, Waltham-Toronto-London: Blaisdell, 1966.
- [12] R. Mohamadian, *r-ideals in commutative rings*, Turk. J. Math., **39** (2015) 733-749.
- [13] R. Y. Sharp, *Steps in Commutative Algebra*, Cambridge university press, 1990.

**Rostam Mohamadian**

Department of Mathematics, Faculty of Mathematical Sciences and Computer

Shahid Chamran University of Ahvaz

Ahvaz, Iran.

mohamadian\_r@scu.ac.ir