

Research Paper

**SPECTRAL ASPECTS OF COMMUTING CONJUGACY CLASS GRAPH
 OF FINITE GROUPS**

PARTHAJIT BHOWAL AND RAJAT KANTI NATH*

ABSTRACT. The commuting conjugacy class graph of a non-abelian group G , denoted by $CCC(G)$, is a simple undirected graph whose vertex set is the set of conjugacy classes of the non-central elements of G and two distinct vertices x^G and y^G are adjacent if there exists some elements $x' \in x^G$ and $y' \in y^G$ such that $x'y' = y'x'$. In this paper we compute various spectra and energies of commuting conjugacy class graph of the groups $D_{2n}, Q_{4m}, U_{(n,m)}, V_{8n}$ and SD_{8n} . Our computation shows that $CCC(G)$ is super integral for these groups. We compare various energies and as a consequence it is observed that $CCC(G)$ satisfy E-LE Conjecture of Gutman et al. We also provide negative answer to a question posed by Dutta et al. comparing Laplacian and Signless Laplacian energy. Finally, we conclude this paper by characterizing the above mentioned groups G such that $CCC(G)$ is hyperenergetic, L-hyperenergetic or Q-hyperenergetic.

DOI: 10.29252/as.2021.1979

MSC(2010): Primary:20D99, 05C50, 15A18, 05C25.

Keywords: commuting conjugacy class graph, spectrum, energy, finite group.

Received: 26 August 2020, Accepted: 15 January 2021.

*Corresponding author

1. INTRODUCTION

The commuting graph of a finite non-abelian group G with center $Z(G)$ is a simple undirected graph whose vertex set is $G \setminus Z(G)$ and two distinct vertices x and y are adjacent if $xy = yx$. This graph was first considered by Brauer and Fowler [3]. Later on, many mathematicians have extended this graph by considering nilpotent graphs, solvable graphs, commuting conjugacy class graphs etc. The commuting conjugacy class graph of a non-abelian group G , denoted by $\mathcal{CCC}(G)$, is a simple undirected graph whose vertex set is the set of conjugacy classes of the non-central elements of G and two distinct vertices x^G and y^G are adjacent if there exists some elements $x' \in x^G$ and $y' \in y^G$ such that $x'y' = y'x'$. The notion of commuting conjugacy class graph of groups was introduced by Herzog, Longobardi and Maj [14] in the year 2009. However, in their definition of $\mathcal{CCC}(G)$, the vertex set is considered to be the set of all non-identity conjugacy classes of G . In the year 2016, Mohammadian et al. [18] have classified all finite groups such that $\mathcal{CCC}(G)$ is triangle-free. Recently, in [20], Salahshour and Ashrafi have obtain the structure of $\mathcal{CCC}(G)$ considering G to be the following groups:

$$D_{2n} = \langle x, y : x^n = y^2 = 1, yxy = x^{-1} \rangle \text{ for } n \geq 3,$$

$$Q_{4m} = \langle x, y : x^{2m} = 1, x^m = y^2, y^{-1}xy = x^{-1} \rangle \text{ for } m \geq 2,$$

$$U_{(n,m)} = \langle x, y : x^{2n} = y^m = 1, x^{-1}yx = y^{-1} \rangle \text{ for } m \geq 2 \text{ and } n \geq 2,$$

$$V_{8n} = \langle x, y : x^{2n} = y^4 = 1, yx = x^{-1}y^{-1}, y^{-1}x = x^{-1}y \rangle \text{ for } n \geq 2,$$

$$SD_{8n} = \langle x, y : x^{4n} = y^2 = 1, yxy = x^{2n-1} \rangle \text{ for } n \geq 2 \text{ and}$$

$$G(p, m, n) = \langle x, y : x^{p^m} = y^{p^n} = [x, y]^p = 1, [x, [x, y]] = [y, [x, y]] = 1 \rangle,$$

where p is any prime, $m \geq 1$ and $n \geq 1$.

In this paper we compute various spectra and energies of commuting conjugacy class graph of the first five groups listed above due to the similar nature of their commuting conjugacy class graphs. In a subsequent paper we shall consider commuting conjugacy class graph of $G(p, m, n)$. Computation of various spectra is helpful to check whether $\mathcal{CCC}(G)$ is super integral. Recall that a graph \mathcal{G} is called super integral if it is integral, L-integral and Q-integral. In the year 1974, Harary and Schwenk [13] introduced the concept of integral graphs. Several results on these graphs can be found in [1, 2, 4, 15, 17, 21]. It is observed that $\mathcal{CCC}(G)$ is super integral for the groups mentioned above. In Section 4, using the energies computed in Section 3, we determine whether the inequalities in [5, Conjecture 1] and [5, Question 1] satisfy for $\mathcal{CCC}(G)$. In Section 5, we determine whether $\mathcal{CCC}(G)$ is hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic or Q-borderenergetic. It is worth

mentioning that various spectra and energies of commuting graphs of finite groups have been computed in [6, 7, 19, 8, 5].

2. DEFINITIONS AND USEFUL RESULTS

Let $A(\mathcal{G})$ and $D(\mathcal{G})$ denote the adjacency matrix and degree matrix of a graph \mathcal{G} respectively. Then the Laplacian matrix and Signless Laplacian matrix of \mathcal{G} are given by $L(\mathcal{G}) = D(\mathcal{G}) - A(\mathcal{G})$ and $Q(\mathcal{G}) = D(\mathcal{G}) + A(\mathcal{G})$ respectively. We write $\text{Spec}(\mathcal{G})$, $\text{L-spec}(\mathcal{G})$ and $\text{Q-spec}(\mathcal{G})$ to denote the spectrum, Laplacian spectrum and Signless Laplacian spectrum of \mathcal{G} . Also, $\text{Spec}(\mathcal{G}) = \{\alpha_1^{a_1}, \alpha_2^{a_2}, \dots, \alpha_l^{a_l}\}$, $\text{L-spec}(\mathcal{G}) = \{\beta_1^{b_1}, \beta_2^{b_2}, \dots, \beta_m^{b_m}\}$ and $\text{Q-spec}(\mathcal{G}) = \{\gamma_1^{c_1}, \gamma_2^{c_2}, \dots, \gamma_n^{c_n}\}$ where $\alpha_1, \alpha_2, \dots, \alpha_n$ are the eigenvalues of $A(\mathcal{G})$ with multiplicities a_1, a_2, \dots, a_l ; $\beta_1, \beta_2, \dots, \beta_m$ are the eigenvalues of $L(\mathcal{G})$ with multiplicities b_1, b_2, \dots, b_m and $\gamma_1, \gamma_2, \dots, \gamma_n$ are the eigenvalues of $Q(\mathcal{G})$ with multiplicities c_1, c_2, \dots, c_n respectively. A graph \mathcal{G} is called integral or L-integral or Q-integral according as $\text{Spec}(\mathcal{G})$ or $\text{L-spec}(\mathcal{G})$ or $\text{Q-spec}(\mathcal{G})$ contains only integers. Following theorem is helpful in computing various spectra.

Theorem 2.1. *If $\mathcal{G} = l_1K_{m_1} \sqcup l_2K_{m_2}$, where $l_iK_{m_i}$ denotes the disjoint union of l_i copies of the complete graph K_{m_i} on m_i vertices for $i = 1, 2$, then*

$$\begin{aligned} \text{Spec}(\mathcal{G}) &= \left\{ (-1)^{\sum_{i=1}^2 l_i(m_i-1)}, (m_1 - 1)^{l_1}, (m_2 - 1)^{l_2} \right\} \\ \text{L-spec}(\mathcal{G}) &= \left\{ 0^{l_1+l_2}, m_1^{l_1(m_1-1)}, m_2^{l_2(m_2-1)} \right\} \text{ and} \\ \text{Q-spec}(\mathcal{G}) &= \left\{ (2m_1 - 2)^{l_1}, (m_1 - 2)^{l_1(m_1-1)}, (2m_2 - 2)^{l_2}, (m_2 - 2)^{l_2(m_2-1)} \right\}. \end{aligned}$$

Depending on the various spectra of a graph, there are various energies called *energy*, *Laplacian energy* and *Signless Laplacian energy* denoted by $E(\mathcal{G})$, $LE(\mathcal{G})$ and $LE^+(\mathcal{G})$ respectively. These energies are defined as follows:

$$\begin{aligned} (1) \quad E(\mathcal{G}) &= \sum_{\lambda \in \text{Spec}(\mathcal{G})} |\lambda|, \\ (2) \quad LE(\mathcal{G}) &= \sum_{\mu \in \text{L-spec}(\mathcal{G})} \left| \mu - \frac{2|e(\Gamma)|}{|V(\Gamma)|} \right|, \\ (3) \quad LE^+(\mathcal{G}) &= \sum_{\nu \in \text{Q-spec}(\mathcal{G})} \left| \nu - \frac{2|e(\Gamma)|}{|V(\mathcal{G})|} \right|, \end{aligned}$$

where $V(\mathcal{G})$ and $e(\mathcal{G})$ denote the sets of vertices and edges of Γ .

In 2008, Gutman et al. [12] posed the following conjecture comparing $E(\mathcal{G})$ and $LE(\mathcal{G})$.

Conjecture 2.2. (E-LE Conjecture) $E(\mathcal{G}) \leq LE(\mathcal{G})$ for any graph \mathcal{G} .

However, in the same year, Stevanović et al. [22] disproved the above conjecture. In 2009, Liu and Lin [16] also disproved Conjecture 2.2 by providing some counter examples. Following Gutman et al. [12], recently Dutta et al. have posed the following question in [5] comparing Laplacian and singless Laplacian energies of graphs.

Question 2.3. Is $LE(\mathcal{G}) \leq LE^+(\mathcal{G})$ for all graphs \mathcal{G} ?

3. VARIOUS SPECTRA AND ENERGIES

In this section we compute various spectra and energies of commuting conjugacy class graphs of the groups mentioned in the introduction.

Theorem 3.1. *If $G = D_{2n}$ then*

$$(i) \text{ Spec}(\text{CCC}(G)) = \begin{cases} \left\{ (-1)^{\frac{n-3}{2}}, 0^1, \left(\frac{n-3}{2}\right)^1 \right\}, & \text{if } n \text{ is odd} \\ \left\{ (-1)^{\frac{n}{2}-2}, 0^2, \left(\frac{n}{2} - 2\right)^1 \right\}, & \text{if } n \text{ and } \frac{n}{2} \text{ are even} \\ \left\{ (-1)^{\frac{n}{2}-1}, 1^1, \left(\frac{n}{2} - 2\right)^1 \right\}, & \text{if } n \text{ is even and } \frac{n}{2} \text{ is odd} \end{cases}$$

$$\text{and } E(\text{CCC}(G)) = \begin{cases} n - 3, & \text{if } n \text{ is odd} \\ n - 4, & \text{if } n \text{ and } \frac{n}{2} \text{ are even} \\ n - 2, & \text{if } n \text{ is even and } \frac{n}{2} \text{ is odd.} \end{cases}$$

$$(ii) \text{ L-spec}(\text{CCC}(G)) = \begin{cases} \left\{ 0^2, \left(\frac{n-1}{2}\right)^{\frac{n-3}{2}} \right\}, & \text{if } n \text{ is odd} \\ \left\{ 0^3, \left(\frac{n}{2} - 1\right)^{\frac{n}{2}-2} \right\}, & \text{if } n \text{ and } \frac{n}{2} \text{ are even} \\ \left\{ 0^2, 2^1, \left(\frac{n}{2} - 1\right)^{\frac{n}{2}-2} \right\}, & \text{if } n \text{ is even and } \frac{n}{2} \text{ is odd} \end{cases}$$

$$\text{and } LE(\text{CCC}(G)) = \begin{cases} \frac{2(n-1)(n-3)}{n+1}, & \text{if } n \text{ is odd} \\ \frac{3(n-2)(n-4)}{n+2}, & \text{if } n \text{ and } \frac{n}{2} \text{ are even} \\ 4, & \text{if } n = 6 \\ \frac{(n-4)(3n-10)}{n+2}, & \text{if } n \text{ is even, } n \geq 10 \text{ and } \frac{n}{2} \text{ is odd.} \end{cases}$$

$$(iii) \text{ Q-spec}(\text{CCC}(G)) = \begin{cases} \left\{ 0^1, (n-3)^1, \left(\frac{n-5}{2}\right)^{\frac{n-3}{2}} \right\}, & \text{if } n \text{ is odd} \\ \left\{ 0^2, (n-4)^1, \left(\frac{n}{2} - 3\right)^{\frac{n}{2}-2} \right\}, & \text{if } n \text{ and } \frac{n}{2} \text{ are even} \\ \left\{ 2^1, 0^1, (n-4)^1, \left(\frac{n}{2} - 3\right)^{\frac{n}{2}-2} \right\}, & \text{if } n \text{ is even and } \frac{n}{2} \text{ is odd} \end{cases}$$

$$\text{and } LE^+(CCC(G)) = \begin{cases} \frac{(n-3)(n+3)}{n+1}, & \text{if } n \text{ is odd} \\ \frac{(n-4)(n+6)}{n+2}, & \text{if } n = 4, 8 \\ \frac{2(n-2)(n-4)}{n+2}, & \text{if } n, \frac{n}{2} \text{ are even and } n \geq 12 \\ 4, & \text{if } n = 6 \\ \frac{22}{3}, & \text{if } n = 10 \\ \frac{2(n-2)(n-6)}{n+2}, & \text{if } n \text{ is even, } n \geq 14 \text{ and } \frac{n}{2} \text{ is odd.} \end{cases}$$

Proof. We shall prove the result by considering the following cases.

Case 1. n is odd.

By [20, Proposition 2.1] we have $CCC(G) = K_1 \sqcup K_{\frac{n-1}{2}}$. Therefore, by Theorem 2.1, it follows that

$$\text{Spec}(CCC(G)) = \left\{ (-1)^{\frac{n-3}{2}}, 0^1, \left(\frac{n-3}{2}\right)^1 \right\}, \quad \text{L-spec}(CCC(G)) = \left\{ 0^2, \left(\frac{n-1}{2}\right)^{\frac{n-3}{2}} \right\}$$

$$\text{and } \text{Q-spec}(CCC(G)) = \left\{ 0^1, (n-3)^1, \left(\frac{n-5}{2}\right)^{\frac{n-3}{2}} \right\}.$$

Hence, by (1), we get

$$E(CCC(G)) = \frac{n-3}{2} + \frac{n-3}{2} = n-3.$$

We have $|V(CCC(G))| = \frac{n+1}{2}$ and $|e(CCC(G))| = \frac{(n-1)(n-3)}{8}$. Therefore, $\frac{2|e(CCC(G))|}{|V(CCC(G))|} = \frac{(n-1)(n-3)}{2(n+1)}$. Also,

$$\left| 0 - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| 0 - \frac{(n-1)(n-3)}{2(n+1)} \right| = \frac{(n-1)(n-3)}{2(n+1)} \quad \text{and}$$

$$\left| \frac{n-1}{2} - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| \frac{n-1}{2} - \frac{(n-1)(n-3)}{2(n+1)} \right| = \frac{2(n-1)}{n+1}.$$

Now, by (2), we have

$$LE(CCC(G)) = 2 \times \frac{(n-1)(n-3)}{2(n+1)} + \frac{n-3}{2} \times \frac{2(n-1)}{n+1} = \frac{2(n-1)(n-3)}{n+1}.$$

Again,

$$\left| n-3 - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| n-3 - \frac{(n-1)(n-3)}{2(n+1)} \right| = \frac{(n-3)(n+3)}{2(n+1)} \quad \text{and}$$

$$\left| \frac{n-5}{2} - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| \frac{n-5}{2} - \frac{(n-1)(n-3)}{2(n+1)} \right| = \left| \frac{-4}{n+1} \right| = \frac{4}{n+1}.$$

By (3), we have

$$LE^+(CCC(G)) = \frac{(n-1)(n-3)}{2(n+1)} + \frac{(n-3)(n+3)}{2(n+1)} + \frac{n-3}{2} \times \frac{4}{n+1} = \frac{(n-3)(n+3)}{n+1}.$$

Case 2. n is even.

Consider the following subcases.

Subcase 2.1 $\frac{n}{2}$ is even.

By [20, Proposition 2.1] we have $\mathcal{CCC}(G) = 2K_1 \sqcup K_{\frac{n}{2}-1}$. Therefore, by Theorem 2.1, it follows that

$$\text{Spec}(\mathcal{CCC}(G)) = \left\{ (-1)^{\frac{n}{2}-2}, 0^2, \left(\frac{n}{2} - 2\right)^1 \right\}, \quad \text{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^3, \left(\frac{n}{2} - 1\right)^{\frac{n}{2}-2} \right\}$$

$$\text{and Q-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, (n-4)^1, \left(\frac{n}{2} - 3\right)^{\frac{n}{2}-2} \right\}.$$

Hence, by (1), we get

$$E(\mathcal{CCC}(G)) = \frac{n}{2} - 2 + \frac{n}{2} - 2 = n - 4.$$

We have $|V(\mathcal{CCC}(G))| = \frac{n}{2} + 1$ and $|e(\mathcal{CCC}(G))| = \frac{(n-2)(n-4)}{8}$. So, $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(n-2)(n-4)}{2(n+2)}$.

Also,

$$\begin{aligned} \left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| 0 - \frac{(n-2)(n-4)}{2(n+2)} \right| = \frac{(n-2)(n-4)}{2(n+2)} \quad \text{and} \\ \left| \frac{n}{2} - 1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| \frac{n}{2} - 1 - \frac{(n-2)(n-4)}{2(n+2)} \right| = \frac{3(n-2)}{n+2}. \end{aligned}$$

Now, by (2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(n-2)(n-4)}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{3(n-2)}{n+2} = \frac{3(n-2)(n-4)}{n+2}.$$

Again,

$$\left| n - 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n - 4 - \frac{(n-2)(n-4)}{2(n+2)} \right| = \frac{(n-4)(n+6)}{2(n+2)} \quad \text{and}$$

$$\left| \frac{n}{2} - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n}{2} - 3 - \frac{(n-2)(n-4)}{2(n+2)} \right| = \left| \frac{n-10}{n+2} \right| = \begin{cases} \frac{-n+10}{n+2}, & \text{if } n = 4, 8 \\ \frac{n-10}{n+2}, & \text{if } n \geq 12. \end{cases}$$

By (3), we have

$$LE^+(\mathcal{CCC}(G)) = 2 \times \frac{(n-2)(n-4)}{2(n+2)} + \frac{(n-4)(n+6)}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{-n+10}{n+2} = \frac{(n-4)(n+6)}{n+2},$$

if $n = 4, 8$. If $n \geq 12$ then

$$LE^+(\mathcal{CCC}(G)) = 2 \times \frac{(n-2)(n-4)}{2(n+2)} + \frac{(n-4)(n+6)}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{n-10}{n+2} = \frac{2(n-2)(n-4)}{n+2}.$$

Subcase 2.2 $\frac{n}{2}$ is odd.

By [20, Proposition 2.1] we have $\mathcal{CCC}(G) = K_2 \sqcup K_{\frac{n}{2}-1}$. Therefore, by Theorem 2.1, it follows that

$$\text{Spec}(\mathcal{CCC}(G)) = \left\{ (-1)^{\frac{n}{2}-1}, 1^1, \left(\frac{n}{2} - 2\right)^1 \right\}, \quad \text{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, 2^1, \left(\frac{n}{2} - 1\right)^{\frac{n}{2}-2} \right\}$$

$$\text{and Q-spec}(\mathcal{CCC}(G)) = \left\{ 2^1, 0^1, (n-4)^1, \left(\frac{n}{2} - 3\right)^{\frac{n}{2}-2} \right\}.$$

Hence, by (1), we get

$$E(\mathcal{CCC}(G)) = \frac{n}{2} - 1 + 1 + \frac{n}{2} - 2 = n - 2.$$

We have $|V(\mathcal{CCC}(G))| = \frac{n}{2} + 1$ and $|e(\mathcal{CCC}(G))| = \frac{(n-2)(n-4)+8}{8}$. Therefore, $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(n-2)(n-4)+8}{2(n+2)}$. Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(n-2)(n-4)+8}{2(n+2)} \right| = \frac{(n-2)(n-4)+8}{2(n+2)},$$

$$\begin{aligned} \left| 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| 2 - \frac{(n-2)(n-4)+8}{2(n+2)} \right| \\ &= \left| \frac{-n^2 + 10n - 8}{2(n+2)} \right| = \begin{cases} 1, & \text{if } n = 6 \\ \frac{n^2 - 10n + 8}{2(n+2)}, & \text{if } n \geq 10 \end{cases} \end{aligned}$$

and

$$\left| \frac{n}{2} - 1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| \frac{n}{2} - 1 - \frac{(n-2)(n-4)+8}{2(n+2)} \right| = \frac{3n-10}{n+2}.$$

Now, by (2), we have

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{(n-2)(n-4)+8}{2(n+2)} + 1 + \left(\frac{n}{2} - 2\right) \times \frac{3n-10}{n+2} = 4,$$

if $n = 6$. If $n \geq 10$ then

$$\begin{aligned} LE(\mathcal{CCC}(G)) &= 2 \times \frac{(n-2)(n-4)+8}{2(n+2)} + \frac{n^2 - 10n + 8}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{3n-10}{n+2} \\ &= \frac{3n^2 - 22n - 40}{n+2} = \frac{(n-4)(3n-10)}{n+2}. \end{aligned}$$

Again,

$$\left| n - 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n - 4 - \frac{(n-2)(n-4)+8}{2(n+2)} \right| = \frac{n^2 + 2n - 32}{2(n+2)} \quad \text{and}$$

$$\begin{aligned} \left| \frac{n}{2} - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| \frac{n}{2} - 3 - \frac{(n-2)(n-4)+8}{2(n+2)} \right| \\ &= \left| \frac{n-14}{n+2} \right| = \begin{cases} \frac{-n+14}{n+2}, & \text{if } n = 6, 10 \\ \frac{n-14}{n+2}, & \text{if } n \geq 14. \end{cases} \end{aligned}$$

By (3), we have

$$LE^+(\mathcal{CCC}(G)) = 1 + \frac{(n-2)(n-4)+8}{2(n+2)} + \frac{n^2 + 2n - 32}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{-n+14}{n+2} = 4,$$

if $n = 6$. If $n = 10$ then

$$\begin{aligned} LE^+(CCC(G)) &= \frac{n^2 - 10n + 8}{2(n+2)} + \frac{(n-2)(n-4) + 8}{2(n+2)} + \frac{n^2 + 2n - 32}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{-n + 14}{n+2} \\ &= \frac{22}{3}. \end{aligned}$$

If $n \geq 14$ then

$$\begin{aligned} LE^+(CCC(G)) &= \frac{n^2 - 10n + 8}{2(n+2)} + \frac{(n-2)(n-4) + 8}{2(n+2)} + \frac{n^2 + 2n - 32}{2(n+2)} + \left(\frac{n}{2} - 2\right) \times \frac{n - 14}{n+2} \\ &= \frac{2(n-2)(n-6)}{n+2}. \end{aligned}$$

This completes the proof. \square

Theorem 3.2. *If $G = Q_{4m}$ then*

$$(i) \text{ Spec}(CCC(G)) = \begin{cases} \{(-1)^{m-1}, 1^1, (m-2)^1\}, & \text{if } m \text{ is odd} \\ \{(-1)^{m-2}, 0^2, (m-2)^1\}, & \text{if } m \text{ is even} \end{cases}$$

$$\text{and } E(CCC(G)) = \begin{cases} 2m - 2, & \text{if } m \text{ is odd} \\ 2m - 4, & \text{if } m \text{ is even.} \end{cases}$$

$$(ii) \text{ L-spec}(CCC(G)) = \begin{cases} \{0^2, 2^1, (m-1)^{m-2}\}, & \text{if } m \text{ is odd} \\ \{0^3, (m-1)^{m-2}\}, & \text{if } m \text{ is even} \end{cases}$$

$$\text{and } LE(CCC(G)) = \begin{cases} 4, & \text{if } m = 3 \\ \frac{2(m-2)(3m-5)}{m+1}, & \text{if } m \text{ is odd and } m \geq 5 \\ \frac{6(m-1)(m-2)}{m+1}, & \text{if } m \text{ is even.} \end{cases}$$

$$(iii) \text{ Q-spec}(CCC(G)) = \begin{cases} \{2^1, 0^1, (2m-4)^1, (m-3)^{m-2}\}, & \text{if } m \text{ is odd} \\ \{0^2, (2m-4)^1, (m-3)^{m-2}\}, & \text{if } m \text{ is even.} \end{cases}$$

$$\text{and } LE^+(CCC(G)) = \begin{cases} 4, & \text{if } m = 3 \\ \frac{22}{3}, & \text{if } m = 5 \\ \frac{4(m-1)(m-3)}{m+1}, & \text{if } m \text{ is odd and } m \geq 7 \\ \frac{2(m-2)(m+3)}{m+1}, & \text{if } m = 2, 4 \\ \frac{4(m-1)(m-2)}{m+1}, & \text{if } m \text{ is even and } m \geq 6. \end{cases}$$

Proof. We shall prove the result by considering the following cases.

Case 1. m is odd.

By [20, Proposition 2.2] we have $CCC(G) = K_2 \sqcup K_{m-1}$. Therefore, by Theorem 2.1, it follows that

$$\text{Spec}(CCC(G)) = \{(-1)^{m-1}, 1^1, (m-2)^1\}, \quad \text{L-spec}(CCC(G)) = \{0^2, 2^1, (m-1)^{m-2}\}$$

and $\text{Q-spec}(CCC(G)) = \{2^1, 0^1, (2m-4)^1, (m-3)^{m-2}\}$.

Hence, by (1), we get

$$E(CCC(G)) = m - 1 + 1 + m - 2 = 2m - 2.$$

We have $|V(CCC(G))| = m + 1$ and $|e(CCC(G))| = \frac{(m-1)(m-2)+2}{2}$. Therefore, $\frac{2|e(CCC(G))|}{|V(CCC(G))|} = \frac{(m-1)(m-2)+2}{m+1}$. Also,

$$\left|0 - \frac{2|e(CCC(G))|}{|V(CCC(G))|}\right| = \left|0 - \frac{(m-1)(m-2)+2}{m+1}\right| = \frac{(m-1)(m-2)+2}{m+1},$$

$$\left|2 - \frac{2|e(CCC(G))|}{|V(CCC(G))|}\right| = \left|2 - \frac{(m-1)(m-2)+2}{m+1}\right|$$

$$= \left|\frac{-m^2+5m-2}{m+1}\right| = \begin{cases} 1, & \text{if } m = 3 \\ \frac{m^2-5m+2}{m+1}, & \text{if } m \geq 5 \end{cases}$$

and

$$\left|m - 1 - \frac{2|e(CCC(G))|}{|V(CCC(G))|}\right| = \left|m - 1 - \frac{(m-1)(m-2)+2}{m+1}\right| = \frac{3m-5}{m+1}.$$

Now, by (2), we have

$$LE(CCC(G)) = 2 \times \frac{(m-1)(m-2)+2}{m+1} + 1 + (m-2) \times \frac{3m-5}{m+1} = 4,$$

if $m = 3$. If $m \geq 5$ then

$$LE(CCC(G)) = 2 \times \frac{(m-1)(m-2)+2}{m+1} + \frac{m^2-5m+2}{m+1} + (m-2) \times \frac{3m-5}{m+1}$$

$$= \frac{2(m-2)(3m-5)}{m+1}.$$

Again,

$$\left|2m - 4 - \frac{2|e(CCC(G))|}{|V(CCC(G))|}\right| = \left|2m - 4 - \frac{(m-1)(m-2)+2}{m+1}\right| = \frac{m^2+m-8}{m+1}$$

and

$$\begin{aligned} \left| m - 3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| m - 3 - \frac{(m-1)(m-2) + 2}{m+1} \right| \\ &= \left| \frac{m-7}{m+1} \right| = \begin{cases} \frac{-m+7}{m+1}, & \text{if } m = 3, 5 \\ \frac{m-7}{m+1}, & \text{if } m \geq 7. \end{cases} \end{aligned}$$

By (3), we have

$$LE^+(\mathcal{CCC}(G)) = 1 + \frac{(m-1)(m-2) + 2}{m+1} + \frac{m^2 + m - 8}{m+1} + (m-2) \times \frac{-m+7}{m+1} = 4,$$

if $m = 3$. If $m = 5$ then

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) &= \frac{m^2 - 5m + 2}{m+1} + \frac{(m-1)(m-2) + 2}{m+1} + \frac{m^2 + m - 8}{m+1} + (m-2) \times \frac{-m+7}{m+1} \\ &= \frac{22}{3}. \end{aligned}$$

If $m \geq 7$ then

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) &= \frac{m^2 - 5m + 2}{m+1} + \frac{(m-1)(m-2) + 2}{m+1} + \frac{m^2 + m - 8}{m+1} + (m-2) \times \frac{m-7}{m+1} \\ &= \frac{4(m-1)(m-3)}{m+1}. \end{aligned}$$

Case 2. m is even.

By [20, Proposition 2.2] we have $\mathcal{CCC}(G) = 2K_1 \sqcup K_{m-1}$. Therefore, by Theorem 2.1, it follows that

$$\begin{aligned} \text{Spec}(\mathcal{CCC}(G)) &= \{(-1)^{m-2}, 0^2, (m-2)^1\}, \quad \text{L-spec}(\mathcal{CCC}(G)) = \{0^3, (m-1)^{m-2}\} \\ \text{and Q-spec}(\mathcal{CCC}(G)) &= \{0^2, (2m-4)^1, (m-3)^{m-2}\}. \end{aligned}$$

Hence, by (1), we get

$$E(\mathcal{CCC}(G)) = m - 2 + m - 2 = 2m - 4.$$

We have $|V(\mathcal{CCC}(G))| = m + 1$ and $|e(\mathcal{CCC}(G))| = \frac{(m-1)(m-2)}{2}$. Therefore, $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(m-1)(m-2)}{m+1}$. Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(m-1)(m-2)}{m+1} \right| = \frac{(m-1)(m-2)}{m+1}$$

and

$$\left| m - 1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| m - 1 - \frac{(m-1)(m-2)}{m+1} \right| = \frac{3(m-1)}{m+1}.$$

Now, by (2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(m-1)(m-2)}{m+1} + (m-2) \times \frac{3(m-1)}{m+1} = \frac{6(m-1)(m-2)}{m+1}.$$

Again,

$$\left| 2m - 4 - \frac{2|e(\text{CCC}(G))|}{|V(\text{CCC}(G))|} \right| = \left| 2m - 4 - \frac{(m-1)(m-2)}{m+1} \right| = \frac{(m-2)(m+3)}{m+1}$$

and

$$\left| m - 3 - \frac{2|e(\text{CCC}(G))|}{|V(\text{CCC}(G))|} \right| = \left| m - 3 - \frac{(m-1)(m-2)}{m+1} \right| = \left| \frac{m-5}{m+1} \right| = \begin{cases} \frac{-m+5}{m+1}, & \text{if } m = 2, 4 \\ \frac{m-5}{m+1}, & \text{if } m \geq 6. \end{cases}$$

By (3), we have

$$\begin{aligned} LE^+(\text{CCC}(G)) &= 2 \times \frac{(m-1)(m-2)}{m+1} + \frac{(m-2)(m+3)}{m+1} + (m-2) \times \frac{-m+5}{m+1} \\ &= \frac{2(m-2)(m+3)}{m+1}, \end{aligned}$$

if $m = 2, 4$. If $m \geq 6$ then

$$\begin{aligned} LE^+(\text{CCC}(G)) &= 2 \times \frac{(m-1)(m-2)}{m+1} + \frac{(m-2)(m+3)}{m+1} + (m-2) \times \frac{m-5}{m+1} \\ &= \frac{4(m-1)(m-2)}{m+1}. \end{aligned}$$

This completes the proof. \square

Theorem 3.3. If $G = \left\{ \left\{ \binom{n(m+1)-4}{(n,m)} \text{ then } \left(\frac{n(m-1)-2}{2} \right)^1, (n-1)^1 \right\}, \text{ if } m \text{ is odd and } n \geq 2 \right.$
 (i) $\text{Spec}(\text{CCC}(G)) = \left. \left\{ \left\{ (-1)^{\frac{n(m+2)-6}{2}}, \left(\frac{n(m-2)-2}{2} \right)^1, (n-1)^2 \right\}, \text{ if } m \text{ is even and } n \geq 2 \right. \right\}$

and

$$E(\text{CCC}(G)) = \begin{cases} n(m+1) - 4, & \text{if } m \text{ is odd and } n \geq 2 \\ 4(n-1), & \text{if } m = 2 \text{ and } n \geq 2 \\ n(m+2) - 6, & \text{if } m \text{ is even, } m \geq 4 \text{ and } n \geq 2. \end{cases}$$

$$(ii) \text{L-spec}(\text{CCC}(G)) = \begin{cases} \left\{ 0^2, \left(\frac{n(m-1)}{2} \right)^{\frac{n(m-1)-2}{2}}, n^{n-1} \right\}, & \text{if } m \text{ is odd and } n \geq 2 \\ \left\{ 0^3, \left(\frac{n(m-2)}{2} \right)^{\frac{n(m-2)-2}{2}}, n^{2n-2} \right\}, & \text{if } m \text{ is even and } n \geq 2 \end{cases}$$

and

$$LE(CCC(G)) = \begin{cases} 4(n-1), & \text{if } m = 3 \text{ and } n \geq 2 \\ \frac{2(2n-1)(n+3)}{3}, & \text{if } m = 5 \text{ and } n \geq 2 \\ \frac{m^2n^2 - 4mn^2 + m^2n + 3n^2 - 2mn - 2m + 5n - 2}{m+1}, & \text{if } m \text{ is odd, } m \geq 7 \\ & \text{and } n \geq 2 \\ 4(n-1), & \text{if } m = 2 \text{ and } n \geq 2 \\ 6(n-1), & \text{if } m = 4 \text{ and } n \geq 2 \\ \frac{2m^2n^2 - 12mn^2 + m^2n + 16n^2 - 4mn - 2m + 12n - 4}{m+2}, & \text{if } m \text{ is even, } m \geq 6 \\ & \text{and } n \geq 2. \end{cases}$$

(iii) $Q\text{-spec}(CCC(G))$

$$= \begin{cases} \left\{ (n(m-1) - 2)^1, \left(\frac{n(m-1)-4}{2} \right)^{\frac{n(m-1)-2}{2}}, (2n-2)^1, (n-2)^{n-1} \right\}, \\ \text{if } m \text{ is odd and } n \geq 2 \\ \left\{ (n(m-2) - 2)^1, \left(\frac{n(m-2)-4}{2} \right)^{\frac{n(m-2)-2}{2}}, (2n-2)^2, (n-2)^{2n-2} \right\}, \\ \text{if } m \text{ is even and } n \geq 2 \end{cases}$$

$$\text{and } LE^+(CCC(G)) = \begin{cases} 4(n-1), & \text{if } m = 3 \text{ and } n \geq 2 \\ \frac{22}{3}, & \text{if } m = 5 \text{ and } n = 2 \\ \frac{2(2n+3)(n-1)}{3}, & \text{if } m = 5 \text{ and } n \geq 3 \\ \frac{n^2(m-1)(m-3)}{m+1}, & \text{if } m \text{ is odd, } m \geq 7 \text{ and } n \geq 2 \\ 4(n-1), & \text{if } m = 2 \text{ and } n \geq 2 \\ 6(n-1), & \text{if } m = 4 \text{ and } n \geq 2 \\ 2(n+2)(n-1), & \text{if } m = 6 \text{ and } n \geq 2 \\ \frac{2n^2(m-2)(m-4)}{m+2}, & \text{if } m \text{ is even, } m \geq 8 \text{ and } n \geq 2. \end{cases}$$

Proof. We shall prove the result by considering the following cases.

Case 1. m is odd.

By [20, Proposition 2.3] we have $CCC(G) = K_{\frac{n(m-1)}{2}} \sqcup K_n$. Therefore, by Theorem 2.1, it follows that

$$\begin{aligned} \text{Spec}(CCC(G)) &= \left\{ (-1)^{\frac{n(m+1)-4}{2}}, \left(\frac{n(m-1)-2}{2} \right)^1, (n-1)^1 \right\}, \\ \text{L-spec}(CCC(G)) &= \left\{ 0^2, \left(\frac{n(m-1)}{2} \right)^{\frac{n(m-1)-2}{2}}, n^{n-1} \right\} \\ \text{and Q-spec}(CCC(G)) &= \left\{ (n(m-1)-2)^1, \left(\frac{n(m-1)-4}{2} \right)^{\frac{n(m-1)-2}{2}}, (2n-2)^1, (n-2)^{n-1} \right\}. \end{aligned}$$

Hence, by (1), we get

$$E(CCC(G)) = \frac{n(m+1)-4}{2} + \frac{n(m-1)-2}{2} + n - 1 = n(m+1) - 4.$$

We have $|V(CCC(G))| = \frac{n(m+1)}{2}$ and $|e(CCC(G))| = \frac{n^2(m-1)^2 - 2n(m-2n+1)}{8}$. Therefore, $\frac{2|e(CCC(G))|}{|V(CCC(G))|} = \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)}$. Also,

$$\left| 0 - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| 0 - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)},$$

since $n(m-1)^2 - 2(m-2n+1) = m^2n - 2m(n+1) + 5n - 2 > 0$;

$$\left| \frac{n(m-1)}{2} - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| \frac{n(m-1)}{2} - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \frac{n(m-3) + m + 1}{m + 1}$$

and

$$\left| n - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| n - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \left| \frac{-f_1(m, n)}{2(m+1)} \right|,$$

where $f_1(m, n) = n(m^2 + 3) - (4mn + 2m + 2)$. For $m = 3$ and $n \geq 2$ we have $f_1(3, n) = -8$. For $m = 5$ and $n \geq 2$ we have $f_1(5, n) = 8n - 12 > 0$. For $m \geq 7$ and $n \geq 2$ we have $m^2 + 3 > m^2 > 4m + 2m + 2$. Therefore, $n(m^2 + 3) > 4mn + (2m + 2)n > 4mn + 2m + 2$ and so $f_1(m, n) > 0$. Hence,

$$\left| \frac{-f_1(m, n)}{2(m+1)} \right| = \begin{cases} 1, & \text{if } m = 3 \text{ and } n \geq 2 \\ \frac{2n-3}{3}, & \text{if } m = 5 \text{ and } n \geq 2 \\ \frac{n(m^2+3)-(4mn+2m+2)}{2(m+1)}, & \text{if } m \geq 7 \text{ and } n \geq 2. \end{cases}$$

Now, by (2), we have

$$\begin{aligned} LE(CCC(G)) &= 2 \times \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} + \frac{n(m-1)-2}{2} \times \frac{n(m-3) + m + 1}{m + 1} \\ &\quad + (n-1) \times 1 \\ &= 4(n-1), \end{aligned}$$

if $m = 3$ and $n \geq 2$. If $m = 5$ and $n \geq 2$ then

$$\begin{aligned} LE(\mathcal{CCC}(G)) &= 2 \times \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} + \frac{n(m-1)-2}{2} \times \frac{n(m-3)+m+1}{m+1} \\ &\quad + (n-1) \times \frac{2n-3}{3} \\ &= \frac{2(2n-1)(n+3)}{3}. \end{aligned}$$

If $m \geq 7$ and $n \geq 2$ then

$$\begin{aligned} LE(\mathcal{CCC}(G)) &= 2 \times \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} + \frac{n(m-1)-2}{2} \times \frac{n(m-3)+m+1}{m+1} \\ &\quad + (n-1) \times \frac{n(m^2+3) - (4mn+2m+2)}{2(m+1)} \\ &= \frac{m^2n^2 - 4mn^2 + m^2n + 3n^2 - 2mn - 2m + 5n - 2}{m+1}. \end{aligned}$$

Again,

$$\begin{aligned} \left| n(m-1) - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| n(m-1) - 2 - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| \\ &= \left| \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} \right| \\ &= \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)}, \end{aligned}$$

since $n(m-1)(m+3) - 2(m+2n+1) = n(m^2-4) - 2 + n(m-3) + m(n-2) > 0$;

$$\begin{aligned} \left| \frac{n(m-1)-4}{2} - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| \frac{n(m-1)-4}{2} - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| \\ &= \left| \frac{f_2(m,n)}{2(m+1)} \right|, \end{aligned}$$

where $f_2(m,n) = n(m-6) - 2 + m(n-2)$. Clearly, for $m \geq 7$ and $n \geq 2$ we have $f_2(m,n) \geq 0$. For $m = 3$ and $n \geq 2$ we have $f_2(3,n) = -8$. Also for $m = 5$ and $n \geq 2$ we have $f_2(5,n) = 4n - 12$. Therefore, $f_2(5,2) = -4$ and $f_2(5,n) \geq 0$ for $n \geq 3$. Hence,

$$\left| \frac{f_2(m,n)}{2(m+1)} \right| = \begin{cases} 1, & \text{if } m = 3 \text{ and } n \geq 2 \\ \frac{1}{3}, & \text{if } m = 5 \text{ and } n = 2 \\ \frac{n-3}{3}, & \text{if } m = 5 \text{ and } n \geq 3 \\ \frac{n(m-3)-m-1}{m+1}, & \text{if } m \geq 7 \text{ and } n \geq 2. \end{cases}$$

$$\left| 2n - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n - 2 - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \left| -\frac{f_3(m,n)}{2(m+1)} \right|,$$

where $f_3(m, n) = mn(m - 6) + 2m + n + 2$. Clearly, $f_3(m, n) > 0$ if $m \geq 7$ and $n \geq 2$. For $m = 3$ and $n \geq 2$ we have $f_3(3, n) = -8n + 8 < 0$. For $m = 5$ and $n \geq 2$ we have $f_3(5, n) = -4n + 12$. Therefore, $f_3(5, 2) = 4$ and $f_3(5, n) \leq 0$ if $n \geq 3$. Hence,

$$\left| -\frac{f_3(m, n)}{2(m+1)} \right| = \begin{cases} n-1, & \text{if } m=3 \text{ and } n \geq 2 \\ \frac{1}{3}, & \text{if } m=5 \text{ and } n=2 \\ \frac{n-3}{3}, & \text{if } m=5 \text{ and } n \geq 3 \\ \frac{mn(m-6)+2m+n+2}{2(m+1)}, & \text{if } m \geq 7 \text{ and } n \geq 2. \end{cases}$$

$$\left| n-2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n-2 - \frac{n(m-1)^2 - 2(m-2n+1)}{2(m+1)} \right| = \left| -\frac{f_4(m, n)}{2(m+1)} \right|,$$

where $f_4(m, n) = mn(m - 2) + 2 - (m(n - 2) + n(m - 3))$. For $m = 3$ and $n \geq 2$ we have $f_4(3, n) = 8$. Also, for $m \geq 5$ and $n \geq 2$ we have

$$mn(m - 2) - 2mn + 2 = mn(m - 4) + 2 > -2m - 3n.$$

Therefore,

$$mn(m - 2) + 2 > 2mn - 2m - 3n = m(n - 2) + n(m - 3)$$

and so $f_4(m, n) > 0$ for $m \geq 5$ and $n \geq 2$. Hence,

$$\left| -\frac{f_4(m, n)}{2(m+1)} \right| = \frac{f_4(m, n)}{2(m+1)} = \frac{mn(m-2) + 2 - m(n-2) - n(m-3)}{2(m+1)}.$$

By (3), we have

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) &= \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times 1 + (n-1) \\ &\quad + (n-1) \times \frac{mn(m-2) + 2 - m(n-2) - n(m-3)}{2(m+1)} \\ &= 4(n-1), \end{aligned}$$

if $m = 3$ and $n \geq 2$. If $m = 5$ and $n = 2$ then

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) &= \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times \frac{1}{3} + \frac{1}{3} \\ &\quad + (n-1) \times \frac{mn(m-2) + 2 - m(n-2) - n(m-3)}{2(m+1)} \\ &= \frac{22}{3}. \end{aligned}$$

If $m = 5$ and $n \geq 3$ then

$$\begin{aligned} LE^+(CCC(G)) &= \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times \frac{n-3}{3} + \frac{n-3}{3} \\ &\quad + (n-1) \times \frac{mn(m-2) + 2 - m(n-2) - n(m-3)}{2(m+1)} \\ &= \frac{2(2n^2 + n - 3)}{3} = \frac{2(2n+3)(n-1)}{3}. \end{aligned}$$

If $m \geq 7$ and $n \geq 2$ then

$$\begin{aligned} LE^+(CCC(G)) &= \frac{n(m-1)(m+3) - 2(m+2n+1)}{2(m+1)} + \frac{n(m-1) - 2}{2} \times \frac{n(m-3) - m - 1}{m+1} \\ &\quad + \frac{mn(m-6) + 2m + n + 2}{2(m+1)} \\ &\quad + (n-1) \times \frac{mn(m-2) + 2 - m(n-2) - n(m-3)}{2(m+1)} \\ &= \frac{n^2(m-1)(m-3)}{m+1}. \end{aligned}$$

Case 2. m is even.

By [20, Proposition 2.3] we have $CCC(G) = K_{\frac{n(m-2)}{2}} \sqcup 2K_n$. Therefore, by Theorem 2.1, it follows that

$$\begin{aligned} \text{Spec}(CCC(G)) &= \left\{ (-1)^{\frac{n(m+2)-6}{2}}, \left(\frac{n(m-2)-2}{2} \right)^1, (n-1)^2 \right\}, \\ \text{L-spec}(CCC(G)) &= \left\{ 0^3, \left(\frac{n(m-2)}{2} \right)^{\frac{n(m-2)-2}{2}}, n^{2n-2} \right\} \\ \text{and Q-spec}(CCC(G)) &= \left\{ (n(m-2)-2)^1, \left(\frac{n(m-2)-4}{2} \right)^{\frac{n(m-2)-2}{2}}, (2n-2)^2, (n-2)^{2n-2} \right\}. \end{aligned}$$

We have

$$\left| \frac{n(m-2)-2}{2} \right| = \begin{cases} 1, & \text{if } m = 2 \\ \frac{n(m-2)-2}{2}, & \text{if } m \geq 4. \end{cases}$$

Therefore, by (1), we have

$$E(CCC(G)) = \frac{n(m+2)-6}{2} + 1 + 2(n-1) = 4(n-1),$$

if $m = 2$. If $m \geq 4$ then

$$E(CCC(G)) = \frac{n(m+2)-6}{2} + \frac{n(m-2)-2}{2} + 2(n-1) = n(m+2) - 6.$$

We have $|V(CCC(G))| = \frac{n(m+2)}{2}$ and $|e(CCC(G))| = \frac{n^2(m-2)^2 - 2n(m-4n+2)}{8}$. Therefore, $\frac{2|e(CCC(G))|}{|V(CCC(G))|} = \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)}$. Also,

$$\left| 0 - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| 0 - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{-f_5(m, n)}{2(m+2)} \right|,$$

where $f_5(m, n) = m(n(m-4) - 2) + 12n - 4$. Note that for $m \geq 6$ we have $f_5(m, n) > 0$ since $n(m-4) > 2$ and $12n - 4 > 0$. For $m = 2$ and $n \geq 2$ we have $f_5(2, n) = 8n - 8 > 0$. For $m = 4$ and $n \geq 2$ we have $f_5(4, n) = 12n - 12 > 0$. Therefore, for all $m \geq 2$ and $n \geq 2$, we have

$$\left| \frac{-f_5(m, n)}{2(m+2)} \right| = \left| \frac{f_5(m, n)}{2(m+2)} \right| = \frac{m(n(m-4) - 2) + 12n - 4}{2(m+2)}.$$

$$\left| \frac{n(m-2)}{2} - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| \frac{n(m-2)}{2} - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{f_6(m, n)}{m+2} \right|,$$

where $f_6(m, n) = 2n(m-4) + m + 2$. Clearly, $f_6(m, n) > 0$ if $m \geq 4$ and $n \geq 2$. For $m = 2$ and $n \geq 2$ we have $f_6(2, n) = -4n + 4 < 0$. Therefore,

$$\left| \frac{f_6(m, n)}{m+2} \right| = \begin{cases} n-1, & \text{if } m=2 \text{ and } n \geq 2 \\ \frac{2n(m-4)+m+2}{m+2}, & \text{if } m \geq 4 \text{ and } n \geq 2. \end{cases}$$

$$\left| n - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| n - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{-f_7(m, n)}{2(m+2)} \right|,$$

where $f_7(m, n) = mn(m-6) - 2m + 8n - 4$. For $m = 2$ and $n \geq 2$ we have $f_7(2, n) = -8$. For $m = 4$ and $n \geq 2$ we have $f_7(4, n) = -12$. For $m = 6$ and $n \geq 2$ we have $f_7(6, n) = 8n - 16 \geq 0$. Also, for $m \geq 8$ and $n \geq 2$ we have $m^2 \geq 8m$ which gives $m(m-6) \geq 2m$ and so $mn(m-6) \geq 2mn > 2m$. Therefore, $mn(m-6) - 2m > 0$ and so $f_7(m, n) > 0$ since $8n - 4 > 0$. Hence,

$$\left| \frac{-f_7(m, n)}{2(m+2)} \right| = \begin{cases} 1, & \text{if } m=2, 4 \text{ and } n \geq 2 \\ \frac{mn(m-6)-2m+8n-4}{2(m+2)}, & \text{if } m \geq 6 \text{ and } n \geq 2. \end{cases}$$

Now, by (2), we have

$$\begin{aligned} LE(CCC(G)) &= 3 \times \frac{m(n(m-4) - 2) + 12n - 4}{2(m+2)} + \frac{n(m-2) - 2}{2} \times (n-1) + (2n-2) \times 1 \\ &= 4(n-1), \end{aligned}$$

if $m = 2$ and $n \geq 2$. If $m = 4$ and $n \geq 2$ then

$$\begin{aligned} LE(CCC(G)) &= 3 \times \frac{m(n(m-4) - 2) + 12n - 4}{2(m+2)} + \frac{n(m-2) - 2}{2} \times \frac{2n(m-4) + m + 2}{m+2} \\ &\quad + (2n-2) \times 1 \\ &= 6(n-1). \end{aligned}$$

If $m \geq 6$ and $n \geq 2$ then

$$\begin{aligned} LE(\mathcal{CCC}(G)) &= 3 \times \frac{m(n(m-4)-2) + 12n - 4}{2(m+2)} + \frac{n(m-2) - 2}{2} \times \frac{2n(m-4) + m + 2}{m+2} \\ &\quad + (2n-2) \times \frac{mn(m-6) - 2m + 8n - 4}{2(m+2)} \\ &= \frac{2m^2n^2 - 12mn^2 + m^2n + 16n^2 - 4mn - 2m + 12n - 4}{m+2}. \end{aligned}$$

Again,

$$\left| n(m-2) - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| n(m-2) - 2 - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{f_8(m, n)}{2(m+2)} \right|,$$

where $f_8(m, n) = n(m^2 - 20) + 2m(n - 1) + 2mn - 4$. For $m = 2$ and $n \geq 2$ we have $f_8(2, n) = -8n - 8 < 0$. For $m = 4$ and $n \geq 2$ we have $f_8(4, n) = 12n - 12 > 0$. For $m \geq 6$ and $n \geq 2$ we have $f_8(m, n) > 0$. Therefore,

$$\left| \frac{f_8(m, n)}{2(m+2)} \right| = \begin{cases} n+1, & \text{if } m=2 \text{ and } n \geq 2 \\ n-1, & \text{if } m=4 \text{ and } n \geq 2 \\ \frac{n(m^2-20)+2m(n-1)+2mn-4}{2(m+2)}, & \text{if } m \geq 6 \text{ and } n \geq 2. \end{cases}$$

$$\begin{aligned} \left| \frac{n(m-2) - 4}{2} - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| \frac{n(m-2) - 4}{2} - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| \\ &= \left| \frac{f_9(m, n)}{m+2} \right|, \end{aligned}$$

where $f_9(m, n) = n(m-8) + m(n-1) - 2$. For $m = 2$ and $n \geq 2$ we have $f_9(2, n) = -4n - 4 < 0$. For $m = 4$ and $n \geq 2$ we have $f_9(4, n) = -6$. For $m = 6$ and $n \geq 2$ we have $f_9(6, n) = 4n - 8 \geq 0$. Further, if For $m \geq 8$ and $n \geq 2$ then $f_9(m, n) > 0$ since $n(m-8) \geq 0$ and $m(n-1) - 2 > 0$. Hence,

$$\left| \frac{f_9(m, n)}{m+2} \right| = \begin{cases} n+1, & \text{if } m=2 \text{ and } n \geq 2 \\ 1, & \text{if } m=4 \text{ and } n \geq 2 \\ \frac{n(m-8)+m(n-1)-2}{m+2}, & \text{if } m \geq 6 \text{ and } n \geq 2. \end{cases}$$

$$\left| 2n - 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n - 2 - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{-f_{10}(m, n)}{2(m+2)} \right|,$$

where $f_{10}(m, n) = n(m^2 - 8m + 4) + 2m + 4$. Clearly, $f_{10}(m, n) > 0$ for $m \geq 8$ and $n \geq 2$. For $m = 2$ and $n \geq 2$ we have $f_{10}(2, n) = -8n + 8 < 0$. For $m = 4$ and $n \geq 2$ we have

$f_{10}(4, n) = -12n + 12 < 0$. For $m = 6$ and $n \geq 2$ we have $f_{10}(6, n) = -8n + 16 \leq 0$. Hence,

$$\left| \frac{f_{10}(m, n)}{m + 2} \right| = \begin{cases} n - 1, & \text{if } m = 2 \text{ and } n \geq 2 \\ n - 1, & \text{if } m = 4 \text{ and } n \geq 2 \\ \frac{n-2}{2}, & \text{if } m = 6 \text{ and } n \geq 2 \\ \frac{n(m^2-8m+4)+2m+4}{2(m+2)}, & \text{if } m \geq 8 \text{ and } n \geq 2. \end{cases}$$

$$\left| n - 2 - \frac{2|e(CCC(G))|}{|V(CCC(G))|} \right| = \left| n - 2 - \frac{n(m-2)^2 - 2(m-4n+2)}{2(m+2)} \right| = \left| \frac{-f_{11}(m, n)}{2(m+2)} \right|,$$

where $f_{11}(m, n) = n(m-2)(m-4) + 2m + 4$. Note that for $m \geq 4$ and $n \geq 2$ we have $f_{11}(m, n) > 0$. For $m = 2$ and $n \geq 2$ we have $f_{11}(m, n) = 8$. Therefore,

$$\left| \frac{-f_{11}(m, n)}{2(m+2)} \right| = \frac{f_{11}(m, n)}{2(m+2)} = \frac{n(m-2)(m-4) + 2m + 4}{2(m+2)}.$$

By (3), we have

$$\begin{aligned} LE^+(CCC(G)) &= n + 1 + \frac{n(m-2) - 2}{2} \times (n + 1) + 2 \times (n - 1) \\ &\quad + (2n - 2) \times \frac{n(m-2)(m-4) + 2m + 4}{2(m+2)} \\ &= 4(n - 1), \end{aligned}$$

if $m = 2$ and $n \geq 2$. If $m = 4$ and $n \geq 2$ then

$$\begin{aligned} LE^+(CCC(G)) &= n - 1 + \frac{n(m-2) - 2}{2} \times 1 + 2 \times (n - 1) \\ &\quad + (2n - 2) \times \frac{n(m-2)(m-4) + 2m + 4}{2(m+2)} \\ &= 6(n - 1). \end{aligned}$$

If $m = 6$ and $n \geq 2$ then

$$\begin{aligned} LE^+(CCC(G)) &= \frac{n(m^2 - 20) + 2m(n - 1) + 2mn - 4}{2(m + 2)} \\ &\quad + \frac{n(m - 2) - 2}{2} \times \frac{n(m - 8) + m(n - 1) - 2}{m + 2} \\ &\quad + 2 \times \frac{n - 2}{2} + (2n - 2) \times \frac{n(m - 2)(m - 4) + 2m + 4}{2(m + 2)} \\ &= 2(n + 2)(n - 1). \end{aligned}$$

If $m \geq 8$ and $n \geq 2$ then

$$\begin{aligned} LE^+(CCC(G)) &= \frac{n(m^2 - 20) + 2m(n - 1) + 2mn - 4}{2(m + 2)} \\ &\quad + \frac{n(m - 2) - 2}{2} \times \frac{n(m - 8) + m(n - 1) - 2}{m + 2} \end{aligned}$$

$$\begin{aligned}
& + 2 \times \frac{n(m^2 - 8m + 4) + 2m + 4}{2(m + 2)} \\
& + (2n - 2) \times \frac{n(m - 2)(m - 4) + 2m + 4}{2(m + 2)} \\
& = \frac{2n^2(m - 2)(m - 4)}{m + 2}.
\end{aligned}$$

This completes the proof. \square

Theorem 3.4. *If $G = V_{8n}$ then*

$$(i) \text{ Spec}(\mathcal{CCC}(G)) = \begin{cases} \{(-1)^{2n-2}, 0^2, (2n-2)^1\}, & \text{if } n \text{ is odd} \\ \{(-1)^{2n-1}, 1^2, (2n-3)^1\}, & \text{if } n \text{ is even} \end{cases}$$

$$\text{and } E(\mathcal{CCC}(G)) = \begin{cases} 4n - 4, & \text{if } n \text{ is odd} \\ 4n - 2, & \text{if } n \text{ is even.} \end{cases}$$

$$(ii) \text{ L-spec}(\mathcal{CCC}(G)) = \begin{cases} \{0^3, (2n-1)^{2n-2}\}, & \text{if } n \text{ is odd} \\ \{0^3, 2^2, (2n-2)^{2n-3}\}, & \text{if } n \text{ is even} \end{cases}$$

$$\text{and } LE(\mathcal{CCC}(G)) = \begin{cases} \frac{6(2n-1)(2n-2)}{2n+1}, & \text{if } n \text{ is odd} \\ 6, & \text{if } n = 2 \\ \frac{2(2n-3)(5n-7)}{n+1}, & \text{if } n \text{ is even and } n \geq 4. \end{cases}$$

$$(iii) \text{ Q-spec}(\mathcal{CCC}(G)) = \begin{cases} \{0^2, (4n-4)^1, (2n-3)^{2n-2}\}, & \text{if } n \text{ is odd} \\ \{2^2, 0^2, (4n-6)^1, (2n-4)^{2n-3}\}, & \text{if } n \text{ is even} \end{cases}$$

$$\text{and } LE^+(\mathcal{CCC}(G)) = \begin{cases} \frac{4(2n-1)(2n-2)}{2n+1}, & \text{if } n \text{ is odd} \\ 6, & \text{if } n = 2 \\ \frac{16(n-1)(n-2)}{n+1}, & \text{if } n \text{ is even and } n \geq 4. \end{cases}$$

Proof. We shall prove the result by considering the following cases.

Case 1. n is odd.

By [20, Proposition 2.4] we have $\mathcal{CCC}(G) = 2K_1 \sqcup K_{2n-1}$. Therefore, by Theorem 2.1, it follows that

$$\begin{aligned}
\text{Spec}(\mathcal{CCC}(G)) &= \{(-1)^{2n-2}, 0^2, (2n-2)^1\}, \quad \text{L-spec}(\mathcal{CCC}(G)) = \{0^3, (2n-1)^{2n-2}\} \\
\text{and } \text{Q-spec}(\mathcal{CCC}(G)) &= \{0^2, (4n-4)^1, (2n-3)^{2n-2}\}.
\end{aligned}$$

Hence, by (1), we get

$$E(\mathcal{CCC}(G)) = 2n - 2 + 2n - 2 = 4n - 4.$$

We have $|V(\mathcal{CCC}(G))| = 2n + 1$ and $|e(\mathcal{CCC}(G))| = \frac{(2n-1)(2n-2)}{2}$. Therefore,

$$\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(2n-1)(2n-2)}{2n+1}.$$

Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{(2n-1)(2n-2)}{2n+1}$$

and

$$\left| 2n-1 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n-1 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{3(2n-1)}{2n+1}.$$

Now, by (2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(2n-1)(2n-2)}{2n+1} + (2n-2) \times \frac{3(2n-1)}{2n+1} = \frac{6(2n-1)(2n-2)}{2n+1}.$$

Again,

$$\left| 4n-4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 4n-4 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{(2n-2)(2n+3)}{2n+1}$$

and

$$\left| 2n-3 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n-3 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{2n-5}{2n+1}.$$

By (3), we have

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) &= 2 \times \frac{(2n-1)(2n-2)}{2n+1} + \frac{(2n-2)(2n+3)}{2n+1} + (2n-2) \times \frac{2n-5}{2n+1} \\ &= \frac{4(2n-1)(2n-2)}{2n+1}. \end{aligned}$$

Case 2. n is even.

By [20, Proposition 2.4] we have $\mathcal{CCC}(G) = 2K_2 \sqcup K_{2n-2}$. Therefore, by Theorem 2.1, it follows that

$$\text{Spec}(\mathcal{CCC}(G)) = \left\{ (-1)^{2n-1}, 1^2, (2n-3)^1 \right\}, \quad \text{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^3, 2^2, (2n-2)^{2n-3} \right\}$$

$$\text{and } \text{Q-spec}(\mathcal{CCC}(G)) = \left\{ 2^2, 0^2, (4n-6)^1, (2n-4)^{2n-3} \right\}.$$

Hence, by (1), we get

$$E(\mathcal{CCC}(G)) = 2n-1 + 2 + 2n-3 = 4n-2.$$

We have $|V(\mathcal{CCC}(G))| = 2n + 2$ and $|e(\mathcal{CCC}(G))| = \frac{(2n-2)(2n-3)+4}{2}$. Therefore, $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(n-1)(2n-3)+2}{n+1}$. Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(n-1)(2n-3)+2}{n+1} \right| = \frac{(n-1)(2n-3)+2}{n+1},$$

$$\begin{aligned}
\left| 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| 2 - \frac{(n-1)(2n-3)+2}{n+1} \right| \\
&= \left| \frac{-(2n-1)(n-3)}{n+1} \right| \\
&= \begin{cases} 1, & \text{if } n = 2 \\ \frac{(2n-1)(n-3)}{n+1}, & \text{if } n \geq 4 \end{cases}
\end{aligned}$$

and

$$\left| 2n-2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n-2 - \frac{(n-1)(2n-3)+2}{n+1} \right| = \frac{5n-7}{n+1}.$$

Now, by (2), we have

$$LE(\mathcal{CCC}(G)) = 3 \times \frac{(n-1)(2n-3)+2}{n+1} + 2 \times 1 + (2n-3) \times \frac{5n-7}{n+1} = 6,$$

if $n = 2$. If $n \geq 4$ then

$$\begin{aligned}
LE(\mathcal{CCC}(G)) &= 3 \times \frac{(n-1)(2n-3)+2}{n+1} + 2 \times \frac{(2n-1)(n-3)}{n+1} + (2n-3) \times \frac{5n-7}{n+1} \\
&= \frac{2(10n^2 - 29n + 21)}{n+1} = \frac{2(2n-3)(5n-7)}{n+1}.
\end{aligned}$$

Again,

$$\left| 4n-6 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 4n-6 - \frac{(n-1)(2n-3)+2}{n+1} \right| = \frac{2n^2+3n-11}{n+1}$$

and

$$\begin{aligned}
\left| 2n-4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| 2n-4 - \frac{(n-1)(2n-3)+2}{n+1} \right| \\
&= \left| \frac{3n-9}{n+1} \right| \\
&= \begin{cases} 1, & \text{if } n = 2 \\ \frac{3n-9}{n+1}, & \text{if } n \geq 4. \end{cases}
\end{aligned}$$

By (3), we have

$$LE^+(\mathcal{CCC}(G)) = 2 \times 1 + 2 \times \frac{(n-1)(2n-3)+2}{n+1} + \frac{2n^2+3n-11}{n+1} + (2n-3) \times 1 = 6,$$

if $n = 2$. If $n \geq 4$ then

$$\begin{aligned} LE^+(CCC(G)) &= 2 \times \frac{(2n-1)(n-3)}{n+1} + 2 \times \frac{(n-1)(2n-3)+2}{n+1} + \frac{2n^2+3n-11}{n+1} \\ &\quad + (2n-3) \times \frac{3n-9}{n+1} \\ &= \frac{16(n-1)(n-2)}{n+1}. \end{aligned}$$

This completes the proof. \square

Theorem 3.5. *If $G = SD_{8n}$ then*

$$(i) \text{ Spec}(CCC(G)) = \begin{cases} \{(-1)^{2n}, 3^1, (2n-3)^1\}, & \text{if } n \text{ is odd} \\ \{(-1)^{2n-2}, 0^2, (2n-2)^1\}, & \text{if } n \text{ is even} \end{cases}$$

$$\text{and } E(CCC(G)) = \begin{cases} 4n, & \text{if } n \text{ is odd} \\ 4n-4, & \text{if } n \text{ is even.} \end{cases}$$

$$(ii) \text{ L-spec}(CCC(G)) = \begin{cases} \{0^2, 4^3, (2n-2)^{2n-3}\}, & \text{if } n \text{ is odd} \\ \{0^3, (2n-1)^{2n-2}\}, & \text{if } n \text{ is even} \end{cases}$$

$$\text{and } LE(CCC(G)) = \begin{cases} 12, & \text{if } n = 3 \\ \frac{2(2n-3)(5n-11)}{n+1}, & \text{if } n \text{ is odd and } n \geq 5 \\ \frac{6(2n-1)(2n-2)}{2n+1}, & \text{if } n \text{ is even.} \end{cases}$$

$$(iii) \text{ Q-spec}(CCC(G)) = \begin{cases} \{6^1, 2^3, (4n-6)^1, (2n-4)^{2n-3}\}, & \text{if } n \text{ is odd} \\ \{0^2, (4n-4)^1, (2n-3)^{2n-2}\}, & \text{if } n \text{ is even} \end{cases}$$

$$\text{and } LE^+(CCC(G)) = \begin{cases} 12, & \text{if } n = 3 \\ 22, & \text{if } n = 5 \\ \frac{16(n-1)(n-3)}{n+1}, & \text{if } n \text{ is odd and } n \geq 7 \\ \frac{28}{5}, & \text{if } n = 2 \\ \frac{4(2n-1)(2n-2)}{2n+1}, & \text{if } n \text{ is even and } n \geq 4. \end{cases}$$

Proof. We shall prove the result by considering the following cases.

Case 1. n is odd.

By [20, Proposition 2.5] we have $\mathcal{CCC}(G) = K_4 \sqcup K_{2n-2}$. Therefore, by Theorem 2.1, it follows that

$$\text{Spec}(\mathcal{CCC}(G)) = \left\{ (-1)^{2n}, 3^1, (2n-3)^1 \right\}, \quad \text{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^2, 4^3, (2n-2)^{2n-3} \right\}$$

$$\text{and } \text{Q-spec}(\mathcal{CCC}(G)) = \left\{ 6^1, 2^3, (4n-6)^1, (2n-4)^{2n-3} \right\}.$$

Hence, by (1), we get

$$E(\mathcal{CCC}(G)) = 2n + 3 + 2n - 3 = 4n.$$

We have $|V(\mathcal{CCC}(G))| = 2n + 2$ and $|e(\mathcal{CCC}(G))| = \frac{(2n-2)(2n-3)+12}{2}$. Therefore, $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(n-1)(2n-3)+6}{n+1}$. Also,

$$\left| 0 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 0 - \frac{(n-1)(2n-3)+6}{n+1} \right| = \frac{(n-1)(2n-3)+6}{n+1},$$

$$\left| 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 4 - \frac{(n-1)(2n-3)+6}{n+1} \right| = \left| \frac{-2n^2+9n-5}{n+1} \right| = \begin{cases} 1, & \text{if } n = 3 \\ \frac{2n^2-9n+5}{n+1}, & \text{if } n \geq 5 \end{cases}$$

and

$$\left| 2n-2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2n-2 - \frac{(n-1)(2n-3)+6}{n+1} \right| = \frac{5n-11}{n+1}.$$

Now, by (2), we have

$$LE(\mathcal{CCC}(G)) = 2 \times \frac{(n-1)(2n-3)+6}{n+1} + 3 \times 1 + (2n-3) \times \frac{5n-11}{n+1} = 12,$$

if $n = 3$. If $n \geq 5$ then

$$\begin{aligned} LE(\mathcal{CCC}(G)) &= 2 \times \frac{(n-1)(2n-3)+6}{n+1} + 3 \times \frac{2n^2-9n+5}{n+1} + (2n-3) \times \frac{5n-11}{n+1} \\ &= \frac{2(10n^2-37n+33)}{n+1} = \frac{2(2n-3)(5n-11)}{n+1}. \end{aligned}$$

Again,

$$\begin{aligned} \left| 6 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| 6 - \frac{(n-1)(2n-3)+6}{n+1} \right| \\ &= \left| \frac{-2n^2+11n-3}{n+1} \right| \\ &= \begin{cases} \frac{-2n^2+11n-3}{n+1}, & \text{if } n = 3, 5 \\ \frac{2n^2-11n+3}{n+1}, & \text{if } n \geq 7, \end{cases} \end{aligned}$$

$$\left| 2 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 2 - \frac{(n-1)(2n-3)+6}{n+1} \right| = \frac{2n^2-7n+7}{n+1},$$

$$\left| 4n-6 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| = \left| 4n-6 - \frac{(n-1)(2n-3)+6}{n+1} \right| = \frac{2n^2+3n-15}{n+1}$$

and

$$\begin{aligned} \left| 2n - 4 - \frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} \right| &= \left| 2n - 4 - \frac{(n-1)(2n-3) + 6}{n+1} \right| \\ &= \left| \frac{3n-13}{n+1} \right| \\ &= \begin{cases} 1, & \text{if } n = 3 \\ \frac{3n-13}{n+1}, & \text{if } n \geq 5. \end{cases} \end{aligned}$$

By (3), we have

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) &= \frac{-2n^2 + 11n - 3}{n+1} + 3 \times \frac{2n^2 - 7n + 7}{n+1} + \frac{2n^2 + 3n - 15}{n+1} + (2n-3) \times 1 \\ &= 12, \end{aligned}$$

if $n = 3$. If $n = 5$ then

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) &= \frac{-2n^2 + 11n - 3}{n+1} + 3 \times \frac{2n^2 - 7n + 7}{n+1} + \frac{2n^2 + 3n - 15}{n+1} \\ &\quad + (2n-3) \times \frac{3n-13}{n+1} \\ &= 22. \end{aligned}$$

If $n \geq 7$ then

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) &= \frac{2n^2 - 11n + 3}{n+1} + 3 \times \frac{2n^2 - 7n + 7}{n+1} + \frac{2n^2 + 3n - 15}{n+1} + (2n-3) \times \frac{3n-13}{n+1} \\ &= \frac{16(n-1)(n-3)}{n+1}. \end{aligned}$$

Case 2. n is even.

By [20, Proposition 2.5] we have $\mathcal{CCC}(G) = 2K_1 \sqcup K_{2n-1}$. Therefore, by Theorem 2.1, it follows that

$$\begin{aligned} \text{Spec}(\mathcal{CCC}(G)) &= \left\{ (-1)^{2n-2}, 0^2, (2n-2)^1 \right\}, \quad \text{L-spec}(\mathcal{CCC}(G)) = \left\{ 0^3, (2n-1)^{2n-2} \right\} \\ \text{and Q-spec}(\mathcal{CCC}(G)) &= \left\{ 0^2, (4n-4)^1, (2n-3)^{2n-2} \right\}. \end{aligned}$$

Hence, by (1), we get

$$E(\mathcal{CCC}(G)) = 2n - 2 + 2n - 2 = 4n - 4.$$

We have $V(\mathcal{CCC}(G)) = 2n + 1$ and $e(\mathcal{CCC}(G)) = \frac{(2n-1)(2n-2)}{2}$. So, $\frac{2|e(\mathcal{CCC}(G))|}{|V(\mathcal{CCC}(G))|} = \frac{(2n-1)(2n-2)}{2n+1}$.

Also,

$$\left| 0 - \frac{2|e(\text{CCC}(G))|}{|V(\text{CCC}(G))|} \right| = \left| 0 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{(2n-1)(2n-2)}{2n+1}$$

and

$$\left| 2n-1 - \frac{2|e(\text{CCC}(G))|}{|V(\text{CCC}(G))|} \right| = \left| 2n-1 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{3(2n-1)}{2n+1}.$$

Now, by (2), we have

$$LE(\text{CCC}(G)) = 3 \times \frac{(2n-1)(2n-2)}{2n+1} + (2n-2) \times \frac{3(2n-1)}{2n+1} = \frac{6(2n-1)(2n-2)}{2n+1}.$$

Again,

$$\left| 4n-4 - \frac{2|e(\text{CCC}(G))|}{|V(\text{CCC}(G))|} \right| = \left| 4n-4 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \frac{(2n-2)(2n+3)}{2n+1} \quad \text{and}$$

$$\left| 2n-3 - \frac{2|e(\text{CCC}(G))|}{|V(\text{CCC}(G))|} \right| = \left| 2n-3 - \frac{(2n-1)(2n-2)}{2n+1} \right| = \left| \frac{2n-5}{2n+1} \right| = \begin{cases} \frac{1}{5}, & \text{if } n=2 \\ \frac{2n-5}{2n+1}, & \text{if } n \geq 4. \end{cases}$$

By (3), we have

$$LE^+(\text{CCC}(G)) = 2 \times \frac{(2n-1)(2n-2)}{2n+1} + \frac{(2n-2)(2n+3)}{2n+1} + (2n-2) \times \frac{1}{5} = \frac{28}{5},$$

if $n=2$. If $n \geq 4$ then

$$\begin{aligned} LE^+(\text{CCC}(G)) &= 2 \times \frac{(2n-1)(2n-2)}{2n+1} + \frac{(2n-2)(2n+3)}{2n+1} + (2n-2) \times \frac{2n-5}{2n+1} \\ &= \frac{4(2n-1)(2n-2)}{2n+1}. \end{aligned}$$

This completes the proof. \square

We conclude this section with the following corollary.

Corollary 3.6. *If G is isomorphic to $D_{2n}, Q_{4m}, U_{(n,m)}, V_{8n}$ or SD_{8n} then $\text{CCC}(G)$ is super integral.*

4. COMPARING VARIOUS ENERGIES

In this section we compare various energies of $\text{CCC}(G)$ obtained in Section 3 and derive the following relations.

Theorem 4.1. *Let $G = D_{2n}$.*

- (i) *If $n = 3, 4, 6$ then $E(\text{CCC}(G)) = LE^+(\text{CCC}(G)) = LE(\text{CCC}(G))$.*
- (ii) *If $n = 5$ then $E(\text{CCC}(G)) < LE^+(\text{CCC}(G)) = LE(\text{CCC}(G))$.*
- (iii) *If $n = 10$ then $LE^+(\text{CCC}(G)) < E(\text{CCC}(G)) < LE(\text{CCC}(G))$.*
- (iv) *If $n \geq 7$ but $n \neq 10$ then $E(\text{CCC}(G)) < LE^+(\text{CCC}(G)) < LE(\text{CCC}(G))$.*

Proof. We shall prove the result by considering the following cases.

Case 1. n is odd.

If $n = 3$ then, by Theorem 3.1, we have

$$E(CCC(G)) = LE^+(CCC(G)) = LE(CCC(G)) = 0.$$

If $n = 5$ then, by Theorem 3.1, we have

$$E(CCC(G)) - LE^+(CCC(G)) = n - 3 - \frac{(n-3)(n+3)}{n+1} = -\frac{4}{5} < 0$$

and $LE^+(CCC(G)) = LE(CCC(G)) = \frac{8}{3}$. Therefore, $E(CCC(G)) < LE^+(CCC(G)) = LE(CCC(G))$.

If $n \geq 7$ then, by Theorem 3.1, we have

$$E(CCC(G)) - LE^+(CCC(G)) = n - 3 - \frac{(n-3)(n+3)}{n+1} = -\frac{2(n-3)}{n+1} < 0$$

and

$$LE^+(CCC(G)) - LE(CCC(G)) = \frac{(n-3)(n+3)}{n+1} - \frac{2(n-1)(n-3)}{n+1} = -\frac{(n-3)(n-5)}{n+1} < 0.$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$.

Case 2. n is even.

Consider the following subcases.

Subcase 2.1 $\frac{n}{2}$ is even.

If $n = 4$ then, by Theorem 3.1, we have

$$E(CCC(G)) = LE^+(CCC(G)) = LE(CCC(G)) = 0.$$

If $n = 8$ then, by Theorem 3.1, we have

$$E(CCC(G)) - LE^+(CCC(G)) = n - 4 - \frac{(n-4)(n+6)}{n+2} = -\frac{8}{5} < 0$$

and

$$LE^+(CCC(G)) - LE(CCC(G)) = \frac{(n-4)(n+6)}{n+2} - \frac{3(n-2)(n-4)}{n+2} = -\frac{8}{5} < 0.$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$.

If $n \geq 12$ then, by Theorem 3.1, we have

$$E(CCC(G)) - LE^+(CCC(G)) = n - 4 - \frac{2(n-2)(n-4)}{n+2} = -\frac{(n-4)(n-6)}{n+2} < 0$$

and

$$\begin{aligned} LE^+(CCC(G)) - LE(CCC(G)) &= \frac{2(n-2)(n-4)}{n+2} - \frac{3(n-2)(n-4)}{n+2} \\ &= -\frac{(n-2)(n-4)}{n+2} < 0. \end{aligned}$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$.

Subcase 2.2 $\frac{n}{2}$ is odd.

If $n = 6$ then, by Theorem 3.1, we have

$$E(CCC(G)) = LE^+(CCC(G)) = LE(CCC(G)) = 4.$$

If $n = 10$ then, by Theorem 3.1, we have

$$LE^+(CCC(G)) - E(CCC(G)) = \frac{22}{3} - (n-2) = -\frac{2}{3} < 0$$

and

$$E(CCC(G)) - LE(CCC(G)) = n - 2 - \frac{(n-4)(3n-10)}{n+2} = -2 < 0.$$

Therefore, $LE^+(CCC(G)) < E(CCC(G)) < LE(CCC(G))$.

If $n \geq 14$ then, by Theorem 3.1, we have

$$\begin{aligned} E(CCC(G)) - LE^+(CCC(G)) &= n - 2 - \frac{2(n-2)(n-6)}{n+2} \\ &= -\frac{(n-2)(n-10)}{n+2} < 0 \end{aligned}$$

and

$$\begin{aligned} LE^+(CCC(G)) - LE(CCC(G)) &= \frac{2(n-2)(n-6)}{n+2} - \frac{(n-4)(3n-10)}{n+2} \\ &= -\frac{n^2 - 6n + 16}{n+2} \\ &= -\frac{n(n-14) + 8n + 10}{n+2} < 0. \end{aligned}$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$. Hence, the result follows. \square

Theorem 4.2. Let $G = Q_{4m}$.

- (i) If $m = 2, 3$ then $E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G))$.
- (ii) If $m = 5$ then $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.
- (iii) If $m = 7$ then $LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.
- (iv) If $m = 4, 6$ or $m \geq 8$ then $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.

Proof. We shall prove the result by considering the following cases.

Case 1. m is odd.

If $m = 3$ then, by Theorem 3.2, we have

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4.$$

If $m = 5$ then, by Theorem 3.2, we have

$$LE^+(\mathcal{CCC}(G)) - E(\mathcal{CCC}(G)) = \frac{22}{3} - (2m - 2) = -\frac{2}{3} < 0$$

and

$$E(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = 2m - 2 - \frac{2(m - 2)(3m - 5)}{m + 1} = -2 < 0.$$

Therefore, $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.

If $m = 7$ then, by Theorem 3.2, we have $LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) = 12$ and

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) &= \frac{4(m - 1)(m - 3)}{m + 1} - \frac{2(m - 2)(3m - 5)}{m + 1} \\ &= -\frac{2(m + 4)(m - 1)}{m + 1} < 0. \end{aligned}$$

Therefore, $LE^+(\mathcal{CCC}(G)) = E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.

If $m \geq 9$ then, by Theorem 3.2, we have

$$E(\mathcal{CCC}(G)) - LE^+(\mathcal{CCC}(G)) = 2m - 2 - \frac{4(m - 1)(m - 3)}{m + 1} = -\frac{2(m - 1)(m - 7)}{m + 1} < 0$$

and

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) &= \frac{4(m - 1)(m - 3)}{m + 1} - \frac{2(m - 2)(3m - 5)}{m + 1} \\ &= -\frac{2(m + 4)(m - 1)}{m + 1} < 0. \end{aligned}$$

Therefore, $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.

Case 2. m is even.

If $m = 2$ then, by Theorem 3.2, we have

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 0.$$

If $m = 4$ then, by Theorem 3.2, we have

$$E(\mathcal{CCC}(G)) - LE^+(\mathcal{CCC}(G)) = 2m - 4 - \frac{2(m - 2)(m + 3)}{m + 1} = -\frac{8}{5} < 0$$

and

$$LE^+(CCC(G)) - LE(CCC(G)) = \frac{2(m-2)(m+3)}{m+1} - \frac{6(m-1)(m-2)}{m+1} = -\frac{8}{5} < 0.$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$.

If $m \geq 6$ then, by Theorem 3.2, we have

$$E(CCC(G)) - LE^+(CCC(G)) = 2m - 4 - \frac{4(m-1)(m-2)}{m+1} = -\frac{2(m-2)(m-3)}{m+1} < 0$$

and

$$\begin{aligned} LE^+(CCC(G)) - LE(CCC(G)) &= \frac{4(m-1)(m-2)}{m+1} - \frac{6(m-1)(m-2)}{m+1} \\ &= -\frac{2(m-1)(m-2)}{m+1}. \end{aligned}$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$. Hence, the result follows. \square

Theorem 4.3. *Let $G = U_{(n,m)}$.*

(i) *If $m = 2, 3, 4$ and $n \geq 2$ then $LE^+(CCC(G)) = E(CCC(G)) = LE(CCC(G))$.*

(ii) *If $m = 5$ and $n = 2, 3$; or $m = 6$ and $n = 2$ then*

$$LE^+(CCC(G)) < E(CCC(G)) < LE(CCC(G)).$$

(iii) *If $m = 5$ and $n \geq 4$; $m \geq 6$ and $n \geq 3$; or $m \geq 8$ and $n \geq 2$ then*

$$E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G)).$$

(iv) *If $m = 7$ and $n = 2$ then $E(CCC(G)) = LE^+(CCC(G)) < LE(CCC(G))$.*

Proof. We shall prove the result by considering the following cases.

Case 1. If m is odd and $n \geq 2$.

If $m = 3$ and $n \geq 2$ then, by Theorem 3.3, we have

$$LE^+(CCC(G)) = E(CCC(G)) = LE(CCC(G)) = 4(n-1).$$

If $m = 5$ and $n = 2$ then, by Theorem 3.3, we have

$$LE^+(CCC(G)) - E(CCC(G)) = \frac{2n^2 + 10n - 6}{3} - (n(m+1) - 4) = -\frac{2}{3} < 0$$

and

$$E(CCC(G)) - LE(CCC(G)) = n(m+1) - 4 - \frac{2(2n-1)(n+3)}{3} = -2 < 0.$$

Therefore, $LE^+(CCC(G)) < E(CCC(G)) < LE(CCC(G))$.

If $m = 5$ and $n = 3$ then, by Theorem 3.3, we have

$$LE^+(CCC(G)) - E(CCC(G)) = \frac{2(2n+3)(n-1)}{3} - (n(m+1) - 4) = -2 < 0$$

and

$$E(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = n(m + 1) - 4 - \frac{2(2n - 1)(n + 3)}{3} = -4 < 0.$$

Therefore, $LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.

If $m = 5$ and $n \geq 4$ then, by Theorem 3.3, we have

$$\begin{aligned} E(\mathcal{CCC}(G)) - LE^+(\mathcal{CCC}(G)) &= n(m + 1) - 4 - \frac{2(2n + 3)(n - 1)}{3} \\ &= \frac{-2(2n^2 - 8n + 3)}{3} \\ &= \frac{-2(2n(n - 4) + 3)}{3} < 0 \end{aligned}$$

and

$$LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) = \frac{2(2n + 3)(n - 1)}{3} - \frac{2(2n - 1)(n + 3)}{3} = \frac{-8n}{3} < 0.$$

Therefore, $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.

If $m \geq 7$ and $n \geq 2$ then, by Theorem 3.3, we have

$$E(\mathcal{CCC}(G)) - LE^+(\mathcal{CCC}(G)) = n(m + 1) - 4 - \frac{n^2(m - 1)(m - 3)}{m + 1} = -\frac{f_1(m, n)}{m + 1},$$

where $f_1(m, n) = mn(m - 4)(n - 3) + 2mn(m - 7) + 3n(n - 1) + 4(m + 1)$. For $m \geq 7$ and $n = 2$ we have $f_1(m, n) = 2(m - 1)(m - 7) \geq 0$. Hence, $f_1(7, 2) = 0$ and $f_1(m, 2) > 0$ if $m \geq 9$. Thus, $E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G))$ and $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G))$ according as if $m = 7, n = 2$ and $m \geq 9, n = 2$. For $m \geq 7$ and $n \geq 3$ we have $f_1(m, n) > 0$ and so $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G))$.

If $m \geq 7$ and $n \geq 2$ then, by Theorem 3.3, we also have

$$\begin{aligned} &LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) \\ &= \frac{n^2(m - 1)(m - 3)}{m + 1} - \frac{m^2n^2 - 4mn^2 + m^2n + 3n^2 - 2mn - 2m + 5n - 2}{m + 1} \\ &= -\frac{m^2n - 2mn - 2m + 5n - 2}{m + 1} \\ &= -\frac{(mn - 2)(m - 2) + 5(n - 2) + 4}{m + 1} < 0. \end{aligned}$$

Therefore, $LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$. Thus, if $m = 7$ and $n = 2$ then

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$$

and if $m \geq 7$ and $n \geq 3$ or $m \geq 9$ and $n = 2$ then

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

Case 2. m is even and $n \geq 2$.

If $m = 2$ and $n \geq 2$ then, by Theorem 3.3, we have

$$LE^+(CCC(G)) = E(CCC(G)) = LE(CCC(G)) = 4(n-1).$$

If $m = 4$ and $n \geq 2$ then, by Theorem 3.3, we have

$$LE^+(CCC(G)) = E(CCC(G)) = LE(CCC(G)) = 6(n-1).$$

If $m = 6$ and $n = 2$ then, by Theorem 3.3, we have

$$LE^+(CCC(G)) - E(CCC(G)) = 2(n+2)(n-1) - (n(m+2) - 6) = -4 < 0$$

and

$$\begin{aligned} & E(CCC(G)) - LE(CCC(G)) \\ &= n(m+2) - 6 - \frac{2m^2n^2 - 12mn^2 + m^2n + 16n^2 - 4mn - 2m + 12n - 4}{m+2} = -2 < 0. \end{aligned}$$

Therefore, $LE^+(CCC(G)) < E(CCC(G)) < LE(CCC(G))$.

If $m = 6$ and $n \geq 3$ then by Theorem 3.3

$$E(CCC(G)) - LE^+(CCC(G)) = n(m+2) - 6 - 2(n+2)(n-1) = 2n(3-n) - 2 < 0$$

and

$$\begin{aligned} & LE^+(CCC(G)) - LE(CCC(G)) \\ &= 2(n+2)(n-1) - \frac{2m^2n^2 - 12mn^2 + m^2n + 16n^2 - 4mn - 2m + 12n - 4}{m+2} \\ &= -(n+2) < 0. \end{aligned}$$

Therefore $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$.

If $m \geq 8$ and $n \geq 2$ then, by Theorem 3.3, we have

$$\begin{aligned} E(CCC(G)) - LE^+(CCC(G)) &= n(m+2) - 6 - \frac{2n^2(m-2)(m-4)}{m+2} \\ &= -\frac{2m^2n^2 - 12mn^2 - m^2n + 16n^2 - 4mn + 6m - 4n + 12}{m+2} \\ &= -\frac{f_2(m,n)}{m+2}, \end{aligned}$$

where $f_2(m,n) = mn(2n-1)(m-8) + 2m(2n(n-3) + 3) + 4n(4n-1) + 12$. For $n = 2$ and $m \geq 8$ we have $f_2(m,n) = (6m-2)(m-8) + 52 > 0$. For $n \geq 3$ and $m \geq 8$ we have $f_2(m,n) > 0$. Therefore, if $m \geq 8$ and $n \geq 2$ then $E(CCC(G)) < LE^+(CCC(G))$.

If $m \geq 8$ and $n \geq 2$ then, by Theorem 3.3, we also have

$$\begin{aligned} &LE^+(CCC(G)) - LE(CCC(G)) \\ &= \frac{2n^2(m-2)(m-4)}{m+2} \\ &\quad - \frac{2m^2n^2 - 12mn^2 + m^2n + 16n^2 - 4mn - 2m + 12n - 4}{m+2} \\ &= -\frac{m^2n - 4mn - 2m + 12n - 4}{m+2} \\ &= -\frac{mn(m-8) + 2m(2n-1) + 4(3n-1)}{m+2} < 0. \end{aligned}$$

Therefore, $LE^+(CCC(G)) < LE(CCC(G))$. Thus, if $m \geq 8$ and $n \geq 2$ then

$$E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G)).$$

Hence, the result follows. \square

Theorem 4.4. *If $G = V_{8n}$ then $E(CCC(G)) \leq LE^+(CCC(G)) \leq LE(CCC(G))$. The equality holds if and only if $n = 2$.*

Proof. We shall prove the result by considering the following cases.

Case 1. n is odd.

By Theorem 3.4, we have

$$E(CCC(G)) - LE^+(CCC(G)) = 4n - 4 - \frac{4(2n-1)(2n-2)}{2n+1} = -\frac{4(n-1)(2n-3)}{2n+1}$$

and

$$\begin{aligned} LE^+(CCC(G)) - LE(CCC(G)) &= \frac{4(2n-1)(2n-2)}{2n+1} - \frac{6(2n-1)(2n-2)}{2n+1} \\ &= -\frac{2(2n-1)(2n-2)}{2n+1} < 0. \end{aligned}$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$.

Case 2. n is even.

If $n = 2$ then, by Theorem 3.4, we have

$$E(CCC(G)) = LE^+(CCC(G)) = LE(CCC(G)) = 6.$$

If $n \geq 4$ then, by Theorem 3.4, we have

$$\begin{aligned} E(CCC(G)) - LE^+(CCC(G)) &= 4n - 2 - \frac{16(n-1)(n-2)}{n+1} \\ &= -\frac{2(6n^2 + 25n - 17)}{n+1} = -\frac{2(6n(n-4) + 49n - 7)}{n+1} < 0 \end{aligned}$$

and

$$\begin{aligned} LE^+(CCC(G)) - LE(CCC(G)) &= \frac{16(n-1)(n-2)}{n+1} - \frac{2(2n-3)(5n-7)}{n+1} \\ &= -\frac{2(2n^2-5n+5)}{n+1} = -\frac{2(2n(n-4)+3n+5)}{n+1} < 0. \end{aligned}$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$. Hence, the result follows. \square

Theorem 4.5. *If $G = SD_{8n}$ then $E(CCC(G)) \leq LE^+(CCC(G)) \leq LE(CCC(G))$. The equality holds if and only if $n = 3$.*

Proof. We shall prove the result by considering the following cases.

Case 1. n is odd.

If $n = 3$ then, by Theorem 3.5, we have

$$E(CCC(G)) = LE^+(CCC(G)) = LE(CCC(G)) = 12.$$

If $n = 5$ then, by Theorem 3.5, we have

$$E(CCC(G)) - LE^+(CCC(G)) = 4n - 22 = -2 < 0$$

and

$$LE^+(CCC(G)) - LE(CCC(G)) = 22 - \frac{2(2n-3)(5n-11)}{n+1} = -\frac{32}{3} < 0.$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$.

If $n \geq 7$ then, by Theorem 3.5, we have

$$\begin{aligned} E(CCC(G)) - LE^+(CCC(G)) &= 4n - \frac{16(n-1)(n-3)}{n+1} \\ &= -\frac{4(3n^2-17n+12)}{n+1} = -\frac{4(3n(n-7)+4n+12)}{n+1} < 0 \end{aligned}$$

and

$$\begin{aligned} LE^+(CCC(G)) - LE(CCC(G)) &= \frac{16(n-1)(n-3)}{n+1} - \frac{2(2n-3)(5n-11)}{n+1} \\ &= -\frac{2(2n^2-5n+9)}{n+1} = -\frac{2(2n(n-7)+9n+9)}{n+1} < 0. \end{aligned}$$

Therefore, $E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G))$.

Case 2. n is even.

If $n = 2$ then, by Theorem 3.5, we have

$$E(CCC(G)) - LE^+(CCC(G)) = 4n - 4 - \frac{28}{5} = -\frac{8}{5} < 0$$

and

$$LE^+(CCC(G)) - LE(CCC(G)) = \frac{28}{5} - \frac{6(2n-1)(2n-2)}{2n+1} = -\frac{8}{5} < 0.$$

Therefore, $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.

If $n \geq 4$ then, by Theorem 3.5, we have

$$E(\mathcal{CCC}(G)) - LE^+(\mathcal{CCC}(G)) = 4n - 4 - \frac{4(2n - 1)(2n - 2)}{2n + 1} = -\frac{4(n - 1)(2n - 3)}{2n + 1} < 0$$

and

$$\begin{aligned} LE^+(\mathcal{CCC}(G)) - LE(\mathcal{CCC}(G)) &= \frac{4(2n - 1)(2n - 2)}{2n + 1} - \frac{6(2n - 1)(2n - 2)}{2n + 1} \\ &= -\frac{2(2n - 1)(2n - 2)}{2n + 1} < 0. \end{aligned}$$

Therefore, $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$. Hence, the result follows. \square

Note that Theorems 4.1–4.5 can be summarized in the following way.

Theorem 4.6. *Let G be a finite non-abelian group. Then we have the following.*

(i) *If G is isomorphic to $D_6, D_8, D_{12}, Q_8, Q_{12}, U_{(n,2)}, U_{(n,3)}, U_{(n,4)}$ ($n \geq 2$), V_{16} or SD_{24} then*

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)).$$

(ii) *If G is isomorphic to $D_{20}, Q_{20}, U_{(2,5)}, U_{(3,5)}$ or $U_{(2,6)}$ then*

$$LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

(iii) *If G is isomorphic to $D_{14}, D_{16}, D_{18}, D_{2n}$ ($n \geq 11$), Q_{16}, Q_{24}, Q_{4m} ($m \geq 8$), $U_{(n,5)}$, ($n \geq 4$), $U_{(n,m)}$ ($m \geq 6$ and $n \geq 3$), $U_{(n,m)}$ ($m \geq 8$ and $n \geq 2$), V_{8n} ($n \geq 3$), SD_{16} or SD_{8n} ($n \geq 4$) then*

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

(iv) *If G is isomorphic to Q_{28} or $U_{(2,7)}$ then $E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$.*

(v) *If G is isomorphic to D_{10} then $E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G))$.*

We conclude this section with the following remark regarding Conjecture 2.2 and Question 2.3.

Remark 4.7. By Theorem 4.6, it follows that $E(\mathcal{CCC}(G)) \leq LE(\mathcal{CCC}(G))$ and $LE^+(\mathcal{CCC}(G)) \leq LE(\mathcal{CCC}(G))$ for commuting conjugacy class graph of the groups $D_{2n}, Q_{4m}, U_{(n,m)}, V_{8n}$ and SD_{8n} . Therefore, Conjecture 2.2 holds for commuting conjugacy class graph of these groups whereas the inequality in Question 2.3 does not. However, $LE(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G))$ if $G = D_6, D_8, D_{10}, D_{12}, Q_8, Q_{12}, V_{16}, SD_{24}$ and $U_{(n,m)}$ where $m = 2, 3, 4$ and $n \geq 2$.

5. HYPERENERGETIC AND BORDERENERGETIC GRAPH

It is well-known that

$$(4) \quad E(K_n) = LE(K_n) = LE^+(K_n) = 2(n-1).$$

A graph \mathcal{G} with n vertices is called hyperenergetic, L-hyperenergetic or Q-hyperenergetic according as $E(K_n) < E(\mathcal{G}), LE(K_n) < LE(\mathcal{G})$ or $LE^+(K_n) < LE^+(\mathcal{G})$. Also, \mathcal{G} is called borderenergetic, L-borderenergetic and Q-borderenergetic if $E(K_n) = E(\mathcal{G}), LE(K_n) = LE(\mathcal{G})$ and $LE^+(K_n) = LE^+(\mathcal{G})$ respectively. These graphs are considered in [24, 11, 10, 23, 9]. In this section we consider commuting conjugacy class graph $CCC(G)$ for the groups considered in Section 3 and determine whether they are hyperenergetic, L-hyperenergetic or Q-hyperenergetic. We shall also determine whether they are borderenergetic, L-borderenergetic or Q-borderenergetic.

Theorem 5.1. *Let $G = D_{2n}$.*

- (i) *If n is odd or $n = 4, 6$ then $CCC(G)$ is neither hyperenergetic, borderenergetic, L-hyper-energetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.*
- (ii) *If $n = 8, 10, 12, 14$ then $CCC(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.*
- (iii) *If n is even and $n \geq 16$ then $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.*

Proof. We shall prove the result by considering the following cases.

Case 1. n is odd.

By [20, Proposition 2.1] we have $CCC(G) = K_1 \sqcup K_{\frac{n-1}{2}}$. Therefore, $|V(CCC(G))| = \frac{n+1}{2}$. Using (4), we get

$$(5) \quad E(K_{|V(CCC(G))|}) = LE^+(K_{|V(CCC(G))|}) = LE(K_{|V(CCC(G))|}) = n-1$$

If $n = 3$ then, by Theorem 4.1 and Theorem 3.1, we get

$$(6) \quad E(CCC(G)) = LE^+(CCC(G)) = LE(CCC(G)) = 0 < 2 = E(K_{|V(CCC(G))|}).$$

If $n = 5$ then, by Theorem 4.1 and Theorem 3.1, we get

$$(7) \quad E(CCC(G)) < LE^+(CCC(G)) = LE(CCC(G)) = \frac{8}{3} < 4 = E(K_{|V(CCC(G))|}).$$

Therefore $CCC(G)$ is neither hyperenergetic nor L-hyperenergetic nor Q-hyperenergetic for $n = 5$.

If $n \geq 7$ then, by Theorem 4.1 and Theorem 3.1, we get

$$E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G)) = \frac{(n-3)(n+3)}{n+1}.$$

Again,

$$\frac{(n-3)(n+3)}{n+1} - (n-1) = -\frac{8}{n+1} < 0$$

Therefore,

$$(8) \quad E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) < n-1 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Hence, in view of (5)–(8), it follows that $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

Case 2. n is even.

By [20, Proposition 2.1] we have $\mathcal{CCC}(G) = 2K_1 \sqcup K_{\frac{n}{2}-1}$. Therefore, $|V(\mathcal{CCC}(G))| = \frac{n}{2} + 1$. Using (4), we get

$$(9) \quad E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = n.$$

Subcase 2.1. $\frac{n}{2}$ is even.

If $n = 4$ then, by Theorem 4.1 and Theorem 3.1, we get

$$(10) \quad E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 0 < 4 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, by (9) and by (10), it follows that $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n = 8$ then, by Theorem 4.1 and Theorem 3.1, we get

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = 6 < 8 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Also,

$$LE(\mathcal{CCC}(G)) = 9 > 8 = E(K_{|V(\mathcal{CCC}(G))|}).$$

So, $\mathcal{CCC}(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n \geq 12$ then, by Theorem 3.1, we get

$$LE(\mathcal{CCC}(G)) = \frac{3(n-2)(n-4)}{n+2}.$$

We have

$$n - \frac{3(n-2)(n-4)}{n+2} = -\frac{2(n(n-12) + 2n + 12)}{n+2} < 0.$$

Therefore, $LE(K_{|V(\mathcal{CCC}(G))|}) < LE(\mathcal{CCC}(G))$ and so $\mathcal{CCC}(G)$ is L-hyperenergetic but not L-borderenergetic.

By Theorem 4.1 and Theorem 3.1, we also get

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{2(n-2)(n-4)}{n+2}.$$

We have

$$(11) \quad \frac{2(n-2)(n-4)}{n+2} - n = \frac{n^2 - 14n + 16}{n+2} = \frac{n(n-16) + 2n + 16}{n+2} := f_1(n)$$

Therefore, for $n = 12$, we have $f_1(n) < 0$ and so

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{2(n-2)(n-4)}{n+2} < n = LE^+(K_{|V(\mathcal{CCC}(G))|}).$$

Thus, if $n = 12$ then $\mathcal{CCC}(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n \geq 16$ then, by (11), we have $f_1(n) > 0$ and so $LE^+(\mathcal{CCC}(G)) > n = LE^+(K_{|V(\mathcal{CCC}(G))|})$. Therefore, $\mathcal{CCC}(G)$ is Q-hyperenergetic but not Q-borderenergetic. Also,

$$E(\mathcal{CCC}(G)) = n - 4 < n = E(K_{|V(\mathcal{CCC}(G))|})$$

and so $\mathcal{CCC}(G)$ is neither hyperenergetic nor borderenergetic. Thus, if $n \geq 16$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Subcase 2.2. $\frac{n}{2}$ is odd.

If $n = 6$ then, by Theorem 4.1 and Theorem 3.1, we get

$$E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4 < 6 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n = 10$ then, by Theorem 4.1 and Theorem 3.1, we get

$$LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = 10 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

So, $\mathcal{CCC}(G)$ is L-borderenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n \geq 14$ then, by Theorem 3.1, we get

$$LE(\mathcal{CCC}(G)) = \frac{(n-4)(3n-10)}{n+2}.$$

We have

$$n - \frac{(n-4)(3n-10)}{n+2} = -\frac{2n(n-14) + 4n + 40}{n+2} < 0.$$

So, $LE(K_{|V(\mathcal{CCC}(G))|}) < LE(\mathcal{CCC}(G))$ and so $\mathcal{CCC}(G)$ is L-hyperenergetic but not L-borderenergetic.

By Theorem 4.1 and Theorem 3.1, we also get

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{2(n-2)(n-6)}{n+2}.$$

We have

$$(12) \quad \frac{2(n-2)(n-6)}{n+2} - n = \frac{n^2 - 18n + 24}{n+2} = \frac{n(n-18) + 24}{n+2} := f_2(n).$$

Therefore, for $n = 14$, we have $f_2(n) < 0$ and so

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{2(n-2)(n-4)}{n+2} < n = LE^+(K_{|V(\mathcal{CCC}(G))|}).$$

Thus, if $n = 14$ then $\mathcal{CCC}(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. If $n \geq 18$ then, by (12), we have $f_2(n) > 0$ and so $LE^+(\mathcal{CCC}(G)) > n = LE^+(K_{|V(\mathcal{CCC}(G))|})$. Therefore, $\mathcal{CCC}(G)$ is Q-hyperenergetic but not Q-borderenergetic. Also, $E(\mathcal{CCC}(G)) = n - 2 < n = E(K_{|V(\mathcal{CCC}(G))|})$ and so $\mathcal{CCC}(G)$ is neither hyperenergetic nor borderenergetic. Thus, if $n \geq 18$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic. \square

Theorem 5.2. *Let $G = Q_{4m}$.*

- (i) *If $m = 2, 3, 4$ then $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.*
- (ii) *If $m = 5$ then $\mathcal{CCC}(G)$ is L-borderenergetic but neither hyperenergetic, borderenergetic, L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic.*
- (iii) *If $m = 6, 7$ then $\mathcal{CCC}(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.*
- (iv) *If $m \geq 8$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.*

Proof. We shall prove the result by considering the following cases.

Case 1. m is odd.

By [20, Proposition 2.2] we have $\mathcal{CCC}(G) = K_2 \sqcup K_{m-1}$. Therefore, $|V(\mathcal{CCC}(G))| = m + 1$. Using (4), we get

$$(13) \quad E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = 2m.$$

If $m = 3$ then, by Theorem 4.2 and Theorem 3.2, we get

$$(14) \quad E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4 < 6 = E(K_{|V(\mathcal{CCC}(G))|}).$$

So, by (13) and (14), $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 5$ then, by Theorem 4.2 and Theorem 3.2, we get

$$LE^+(\mathcal{CCC}(G)) < E(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = 10 = E(K_{|V(\mathcal{CCC}(G))|}).$$

So, $CCC(G)$ is L-borderenergetic but neither hyperenergetic, borderenergetic, L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 7$ then, by Theorem 4.2 and Theorem 3.2, we get

$$LE^+(CCC(G)) = E(CCC(G)) = 12 < 14 = E(K_{|V(CCC(G))|}).$$

Also,

$$LE(CCC(G)) = 20 > 14 = LE(K_{|V(CCC(G))|}).$$

So, $CCC(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m \geq 9$ then, by Theorem 4.2 and Theorem 3.2, we get

$$\frac{4(m-1)(m-3)}{m+1} = LE^+(CCC(G)) < LE(CCC(G)).$$

We have

$$2m - \frac{4(m-1)(m-3)}{m+1} = -\frac{2(m(m-9)+6)}{m+1} < 0$$

and so $LE^+(K_{|V(CCC(G))|}) = 2m < \frac{4(m-1)(m-3)}{m+1} = LE^+(CCC(G)) < LE(CCC(G))$. Hence, $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Q-borderenergetic. Also,

$$E(CCC(G)) = 2m - 2 < 2m = E(K_{|V(CCC(G))|}).$$

Therefore, $CCC(G)$ is neither hyperenergetic nor borderenergetic. Thus, if $m \geq 9$ then $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Case 2. m is even.

By [20, Proposition 2.2] we have $CCC(G) = 2K_1 \sqcup K_{m-1}$. Therefore, $|V(CCC(G))| = m + 1$. Using (4), we get

$$(15) \quad E(K_{|V(CCC(G))|}) = LE^+(K_{|V(CCC(G))|}) = LE(K_{|V(CCC(G))|}) = 2m.$$

If $m = 2$ then, by Theorem 4.2 and Theorem 3.2, we get

$$(16) \quad E(CCC(G)) = LE^+(CCC(G)) = LE(CCC(G)) = 0 < 4 = E(K_{|V(CCC(G))|}).$$

So, by (15) and (16), $CCC(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 4$ then, by Theorem 4.2 and Theorem 3.2, we get

$$(17) \quad E(CCC(G)) < LE^+(CCC(G)) < LE(CCC(G)) = \frac{36}{5} < 8 = E(K_{|V(CCC(G))|}).$$

So, by (15) and (17), $CCC(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m \geq 6$ then, by Theorem 3.2, we get

$$LE(\mathcal{CCC}(G)) = \frac{6(m-1)(m-2)}{m+1}.$$

We have

$$2m - \frac{6(m-1)(m-2)}{m+1} = -\frac{4(m^2 - 5m + 3)}{m+1} = -\frac{4(m(m-6) + m + 3)}{m+1} < 0$$

and so

$$LE(K_{|V(\mathcal{CCC}(G))|}) = 2m < \frac{6(m-1)(m-2)}{m+1} = LE(\mathcal{CCC}(G)).$$

Hence, $\mathcal{CCC}(G)$ is L-hyperenergetic but not L-borderenergetic.

By Theorem 4.2 and Theorem 3.2, we also get

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{4(m-1)(m-2)}{m+1}.$$

We have

$$(18) \quad \frac{4(m-1)(m-2)}{m+1} - 2m = \frac{2(m^2 - 7m + 4)}{m+1} = \frac{2(m(m-8) + m + 4)}{m+1} = f(m).$$

Therefore, for $m = 6$, we have $f(m) < 0$ and so

$$E(\mathcal{CCC}(G)) < LE^+(\mathcal{CCC}(G)) = \frac{4(m-1)(m-2)}{m+1} < 2m = LE^+(K_{|V(\mathcal{CCC}(G))|}).$$

Thus, if $m = 6$ then $\mathcal{CCC}(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n \geq 8$ then, by (18), we have $f(m) > 0$ and so $LE^+(\mathcal{CCC}(G)) > 2m = LE^+(K_{|V(\mathcal{CCC}(G))|})$. Therefore, $\mathcal{CCC}(G)$ is Q-hyperenergetic but not Q-borderenergetic. Also, $E(\mathcal{CCC}(G)) = 2m - 4 < 2m = E(K_{|V(\mathcal{CCC}(G))|})$ and so $\mathcal{CCC}(G)$ is neither hyperenergetic nor borderenergetic. Thus, if $n \geq 8$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic. \square

Theorem 5.3. *Let $G = U_{(n,m)}$.*

- (i) *If $m = 2, 3, 4$ and $n \geq 2$ or $m = 6$ and $n = 2$ then $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.*
- (ii) *If $m = 5$ and $n = 2$ then $\mathcal{CCC}(G)$ is L-borderenergetic but neither hyperenergetic, borderenergetic, L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic.*
- (iii) *If $m = 5$ and $n = 3$, $m = 6$ and $n = 3$ or $m = 7$ and $n = 2$ then $\mathcal{CCC}(G)$ is L-hyper-energetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.*
- (iv) *If $m = 5, 6$ and $n \geq 4$; $m = 7$ and $n \geq 3$ or $m \geq 8$ and $n \geq 2$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-border-energetic nor Q-borderenergetic.*

Proof. We shall prove the result by considering the following cases.

Case 1. m is odd and $n \geq 2$.

By [20, Proposition 2.3] we have $\mathcal{CCC}(G) = K_{\frac{n(m-1)}{2}} \sqcup K_n$. Therefore, $|V(\mathcal{CCC}(G))| = \frac{n(m+1)}{2}$. Using (4), we get

$$(19) \quad E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = mn + n - 2.$$

By Theorem 3.3 we get

$$E(\mathcal{CCC}(G)) = mn + n - 4 < mn + n - 2.$$

Therefore, $\mathcal{CCC}(G)$ is neither hyperenergetic nor borderenergetic.

If $m = 3$ and $n \geq 2$ then, by Theorem 3.3, we get

$$LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4n - 4 < 4n - 2 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if $m = 3$ and $n \geq 2$ then $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 5$ and $n = 2$ then, by Theorem 4.3 and Theorem 3.3, we get

$$LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = 10 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is L-borderenergetic but neither L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if $m = 5$ and $n = 2$ then $\mathcal{CCC}(G)$ is L-borderenergetic but neither hyperenergetic, borderenergetic, L-hyperenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 5$ and $n = 3$ then, by Theorem 3.3, we get

$$LE(\mathcal{CCC}(G)) = 20 > 16 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is L-hyperenergetic but not L-borderenergetic. Also,

$$LE^+(\mathcal{CCC}(G)) = 12 < 16 = LE^+(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is neither Q-hyperenergetic nor Q-borderenergetic. Thus, if $m = 5$ and $n = 3$ then $\mathcal{CCC}(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 5$ and $n \geq 4$ then, by Theorem 4.3 and Theorem 3.3, we get

$$\frac{2(2n+3)(n-1)}{3} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

We have

$$6n - 2 - \frac{2(2n+3)(n-1)}{3} = -\frac{4(n^2 - n - 1)}{3} = -\frac{4(n(n-4) + 3n - 1)}{3} < 0$$

and so $LE^+(K_{|V(\mathcal{CCC}(G))|}) = 6n - 2 < \frac{2(2n+3)(n-1)}{3} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$. Therefore, $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Q-borderenergetic. Thus, if $m = 5$ and $n \geq 4$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

If $m \geq 7$ and $n \geq 2$ then, by Theorem 3.3, we get

$$LE(\mathcal{CCC}(G)) = \frac{m^2n^2 - 4mn^2 + m^2n + 3n^2 - 2mn - 2m + 5n - 2}{m + 1}.$$

We have

$$\begin{aligned} mn + n - 2 - LE(\mathcal{CCC}(G)) &= -\frac{m^2n^2 - 4mn^2 - 4mn + 3n^2 + 4n}{m + 1} \\ &= -\frac{mn^2(m - 7) + 2mn(n - 2) + mn^2 + 3n^2 + 4n}{m + 1} < 0 \end{aligned}$$

and so $LE(K_{|V(\mathcal{CCC}(G))|}) = mn + n - 2 < LE(\mathcal{CCC}(G))$. Therefore, $\mathcal{CCC}(G)$ is L-hyperenergetic but not L-borderenergetic. By Theorem 4.3 and Theorem 3.3, we also get

$$\frac{n^2(m - 1)(m - 3)}{m + 1} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

Let $f_1(m, n) = \frac{n^2(m-1)(m-3)}{m+1} - (mn + n - 2)$. Then

$$\begin{aligned} f_1(m, n) &= \frac{2 + 2m - 2mn - m^2n - n + 3n^2 - 4mn^2 + m^2n^2}{m + 1} \\ &= \frac{mn^2(m - 11) + mn^2 + m^2n(n - 2) + 2mn(n - 2) + 2n(3n - 1) + 2(m + 1)}{2(m + 1)}. \end{aligned}$$

For $m = 7$ and $n = 2$ we have $f_1(m, n) = -2 < 0$ and so

$$LE^+(\mathcal{CCC}(G)) = \frac{n^2(m - 1)(m - 3)}{m + 1} < mn + n - 2 = LE^+(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is neither Q-hyperenergetic nor Q-borderenergetic. Thus, if $m = 7$ and $n = 2$ then $\mathcal{CCC}(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 7$ and $n \geq 3$ then $f_1(m, n) = \frac{n(3n-8)+16}{8} > 0$. Therefore,

$$LE^+(K_{|V(\mathcal{CCC}(G))|}) = mn + n - 2 < \frac{n^2(m - 1)(m - 3)}{m + 1} = LE^+(\mathcal{CCC}(G))$$

and so $\mathcal{CCC}(G)$ is Q-hyperenergetic but not Q-borderenergetic. Thus, if $m = 7$ and $n \geq 3$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Now, for $m = 9$ and $n = 2$ we have $f_1(m, n) = \frac{6}{5} > 0$. For $m = 9$ and $n \geq 3$ we have $f_1(m, n) = \frac{2n(12n-25)+10}{5} > 0$. For $m \geq 11$ and $n \geq 2$ we have $f_1(m, n) > 0$. Therefore, for $m \geq 9$ and $n \geq 2$ we have

$$LE^+(K_{|V(\mathcal{CCC}(G))|}) = mn + n - 2 < \frac{n^2(m - 1)(m - 3)}{m + 1} = LE^+(\mathcal{CCC}(G))$$

and so $\mathcal{CCC}(G)$ is Q-hyperenergetic but not Q-borderenergetic. Thus, if $m \geq 9$ and $n \geq 2$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Case 2. m is even and $n \geq 2$.

By [20, Proposition 2.3] we have $\mathcal{CCC}(G) = K_{\frac{n(m-2)}{2}} \sqcup 2K_n$. Therefore, $|V(\mathcal{CCC}(G))| = \frac{n(m+2)}{2}$. Using (4), we get

$$(20) \quad E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = mn + 2n - 2.$$

By Theorem 3.3 we get

$$E(\mathcal{CCC}(G)) = 4n - 4 < 4n - 2 = E(K_{|V(\mathcal{CCC}(G))|}),$$

if $m = 2$. If $m \geq 4$ then

$$E(\mathcal{CCC}(G)) = mn + 2n - 6 < mn + 2n - 2 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is neither hyperenergetic nor borderenergetic.

If $m = 2$ and $n \geq 2$ then, by Theorem 3.3, we get

$$LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 4n - 4 < 4n - 2.$$

Therefore, $\mathcal{CCC}(G)$ is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if $m = 2$ and $n \leq 2$ then $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 4$ and $n \geq 2$ then, by Theorem 3.3, we get

$$LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 6n - 6 < 6n - 2.$$

Therefore, $\mathcal{CCC}(G)$ is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if $m = 4$ and $n \leq 2$ then $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 6$ and $n = 2$ then, Theorem 4.3 and Theorem 3.3, we get

$$LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)) = 12 < 14 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if $m = 6$ and $n = 2$ then $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $m = 6$ and $n \geq 3$ then, by Theorem 3.3, we get

$$LE(\mathcal{CCC}(G)) = 2n^2 + 3n - 2.$$

We have

$$8n - 2 - (2n^2 + 3n - 2) = -n(2n - 5) < 0.$$

Therefore,

$$LE(K_{|V(\mathcal{CCC}(G))|}) = 8n - 2 < 2n^2 + 3n - 2 < LE(\mathcal{CCC}(G))$$

and so $\mathcal{CCC}(G)$ is L-hyperenergetic but not L-borderenergetic. By Theorem 3.3, we also get

$$LE^+(\mathcal{CCC}(G)) = 2(n + 2)(n - 1).$$

Let $g(n) = 2(n + 2)(n - 1) - (8n - 2)$. Then $g(n) = 2(n(n - 4) + n - 1)$. Therefore, if $n = 3$ then $g(n) = -2 < 0$ and so

$$LE^+(\mathcal{CCC}(G)) = 2(n + 2)(n - 1) < 8n - 2 = LE^+(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is neither Q-hyperenergetic nor Q-borderenergetic. Thus, if $m = 6$ and $n = 3$ then $\mathcal{CCC}(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n \geq 4$ then $g(n) > 0$ and so

$$LE^+(K_{|V(\mathcal{CCC}(G))|}) = 8n - 2 < 2(n + 2)(n - 1) = LE^+(\mathcal{CCC}(G)).$$

Therefore, $\mathcal{CCC}(G)$ is Q-hyperenergetic but not Q-borderenergetic. Thus, if $m = 6$ and $n \geq 4$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

If $m \geq 8$ and $n \geq 2$ then, by Theorem 4.3 and Theorem 3.3, we get

$$\frac{2n^2(m - 2)(m - 4)}{m + 2} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

We have

$$\begin{aligned} mn + 2n - 2 - \frac{2n^2(m - 2)(m - 4)}{m + 2} &= -\frac{4 + 2m - m^2n - 4n - 4mn + 16n^2 - 12mn^2 + 2m^2n^2}{m + 2} \\ &= -f_2(m, n), \end{aligned}$$

where $f_2(m, n) = \frac{mn^2(m-12)+m^2n(n-2)+mn(m-6)+2n(m-2)+16n^2+2m+4}{m+2}$.

For $m = 8$ and $n = 2$ we have $f_2(m, n) = \frac{6}{5} > 0$. For $m = 8$ and $n \geq 3$ we have $f_2(m, n) = \frac{2}{5}(12n^2 - 25n + 5) = \frac{2}{5}(12n(n - 3) + 11n + 5) > 0$. For $m = 10$ and $n \geq 2$ we have $f_2(m, n) = 2(4n^2 - 6n + 1) = 2(4n(n - 2) + 2n + 1) > 0$. For $m \geq 12$ and $n \geq 2$ we have $f_2(m, n) > 0$. Therefore,

$$LE^+(K_{|V(\mathcal{CCC}(G))|}) = mn + 2n - 2 < \frac{2n^2(m - 2)(m - 4)}{m + 2} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G))$$

and so $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Q-borderenergetic. Thus, if $m \geq 8$ and $n \geq 2$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic. \square

Theorem 5.4. *Let $G = V_{8n}$.*

- (i) *If $n = 2$ then $CCC(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-border-energetic, Q-hyperenergetic nor Q-borderenergetic.*
- (ii) *If $n = 3, 4$ then $CCC(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.*
- (iii) *If $n \geq 5$ then $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.*

Proof. We shall prove the result by considering the following cases.

Case 1. n is odd.

By [20, Proposition 2.4] we have $CCC(G) = 2K_1 \sqcup K_{2n-1}$. Therefore, $|V(CCC(G))| = 2n + 1$. Using (4), we get

$$(21) \quad E(K_{|V(CCC(G))|}) = LE^+(K_{|V(CCC(G))|}) = LE(K_{|V(CCC(G))|}) = 4n.$$

By Theorem 3.4 we get

$$LE(CCC(G)) = \frac{6(2n-1)(2n-2)}{2n+1}.$$

We have

$$4n - \frac{6(2n-1)(2n-2)}{2n+1} = -\frac{4(4n^2 - 10n + 3)}{2n+1} = -\frac{4(4n(n-3) + 2n + 3)}{2n+1} < 0$$

and so $LE(K_{|V(CCC(G))|}) = 4n < \frac{6(2n-1)(2n-2)}{2n+1} = LE(CCC(G))$. Hence, $CCC(G)$ is L-hyperenergetic but not L-borderenergetic.

By Theorem 4.4 and Theorem 3.4, we also get

$$E(CCC(G)) < LE^+(CCC(G)) = \frac{4(2n-1)(2n-2)}{2n+1}.$$

We have

$$(22) \quad \frac{4(2n-1)(2n-2)}{2n+1} - 4n = \frac{4(2n^2 - 7n + 2)}{2n+1} = \frac{4(2n(n-5) + 3n + 2)}{2n+1} := g_1(n).$$

Therefore, for $n = 3$, we have $g_1(n) < 0$ and so

$$E(CCC(G)) < LE^+(CCC(G)) = \frac{4(2n-1)(2n-2)}{2n+1} < 4n = LE^+(K_{|V(CCC(G))|}).$$

Thus, if $n = 3$ then $CCC(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. If $n \geq 5$ then, by (22), we have $g_1(n) > 0$ and so $LE^+(CCC(G)) > 4n = LE^+(K_{|V(CCC(G))|})$. Therefore, $CCC(G)$ is Q-hyperenergetic but not Q-borderenergetic. Also, $E(CCC(G)) = 4n - 4 < 4n = E(K_{|V(CCC(G))|})$ and so $CCC(G)$ is neither hyperenergetic nor borderenergetic. Thus, if $n \geq 5$ then $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Case 2. n is even.

By [20, Proposition 2.4] we have $\mathcal{CCC}(G) = 2K_2 \sqcup K_{2n-2}$. Therefore $|V(\mathcal{CCC}(G))| = 2n + 2$. Using (4), we get

$$(23) \quad E(K_{|V(\mathcal{CCC}(G))|}) = LE^+(K_{|V(\mathcal{CCC}(G))|}) = LE(K_{|V(\mathcal{CCC}(G))|}) = 4n + 2.$$

If $n = 2$ then, by Theorem 4.4 and Theorem 3.4, we get

$$(24) \quad E(\mathcal{CCC}(G)) = LE^+(\mathcal{CCC}(G)) = LE(\mathcal{CCC}(G)) = 6 < 10 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, by (23) and (24), we have $\mathcal{CCC}(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n \geq 4$ then, Theorem 3.4, we get

$$E(\mathcal{CCC}(G)) = 4n - 2 < 4n + 2 = E(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is neither hyperenergetic nor borderenergetic.

By Theorem 4.4 and Theorem 3.4, we also get

$$\frac{16(n-1)(n-2)}{n+1} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

We have

$$(25) \quad \frac{16(n-1)(n-2)}{n+1} - (4n+2) = \frac{6(2n^2 - 9n + 5)}{n+1} = \frac{6(2n(n-6) + 3n + 5)}{n+1} := g_2(n).$$

Therefore, for $n = 4$ we have $g_2(n) < 0$ and so

$$LE^+(\mathcal{CCC}(G)) = \frac{16(n-1)(n-2)}{n+1} < 4n + 2 = LE^+(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is neither Q-hyperenergetic nor Q-borderenergetic. Also,

$$LE(\mathcal{CCC}(G)) = \frac{130}{5} = 26 > 18 = LE(K_{|V(\mathcal{CCC}(G))|}).$$

Therefore, $\mathcal{CCC}(G)$ is L-hyperenergetic but not L-borderenergetic. Thus, if $n = 4$ then $\mathcal{CCC}(G)$ is L-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n \geq 6$ then, by (25), we have $g_2(n) > 0$ and so

$$LE^+(K_{|V(\mathcal{CCC}(G))|}) = 4n + 2 < \frac{16(n-1)(n-2)}{n+1} = LE^+(\mathcal{CCC}(G)) < LE(\mathcal{CCC}(G)).$$

Therefore, $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Q-borderenergetic. Thus, if $n \geq 6$ then $\mathcal{CCC}(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic. \square

Theorem 5.5. *Let $G = SD_{8n}$.*

(i) *If $n = 2, 3$ then $CCC(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.*

(ii) *If $n = 5$ then $CCC(G)$ is L-hyperenergetic and Q-borderenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-hyperenergetic.*

(iii) *If $n = 4$ or $n \geq 6$ then $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.*

Proof. We shall prove the result by considering the following cases.

Case 1. n is odd.

By [20, Proposition 2.5] we have $CCC(G) = K_4 \sqcup K_{2n-2}$. Therefore, $|V(CCC(G))| = 2n + 2$. Using (4), we get

$$(26) \quad E(K_{|V(CCC(G))|}) = LE^+(K_{|V(CCC(G))|}) = LE(K_{|V(CCC(G))|}) = 4n + 2.$$

By Theorem 3.5 we get

$$E(CCC(G)) = 4n < 4n + 2.$$

Therefore, $CCC(G)$ is neither hyperenergetic nor borderenergetic.

If $n = 3$ then, by Theorem 3.5, we get

$$LE^+(CCC(G)) = LE(CCC(G)) = 12 < 14 = LE(K_{|V(CCC(G))|}).$$

Therefore, $CCC(G)$ is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if $n = 3$ then $CCC(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic.

If $n = 5$ then, by Theorem 4.5 and Theorem 3.5, we get

$$LE^+(K_{|V(CCC(G))|}) = 22 = LE^+(CCC(G)) < LE(CCC(G)).$$

Therefore, $CCC(G)$ is L-hyperenergetic and Q-borderenergetic but neither L-borderenergetic nor Q-hyperenergetic. Thus, if $n = 5$ then $CCC(G)$ is L-hyperenergetic and Q-borderenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-hyperenergetic.

If $n \geq 7$ then, by Theorem 4.5 and Theorem 3.5, we get

$$\frac{16(n-1)(n-3)}{n+1} = LE^+(CCC(G)) < LE(CCC(G)).$$

We have

$$4n + 2 - \frac{16(n-1)(n-3)}{n+1} = -\frac{2(6n^2 - 35n + 23)}{n+1} = -\frac{2(6n(n-7) + 7n + 23)}{n+1} < 0.$$

So, $LE^+(K_{|V(CCC(G))|}) = 4n+2 < \frac{16(n-1)(n-3)}{n+1} = LE^+(CCC(G)) < LE(CCC(G))$ and so $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Q-borderenergetic.

Thus, if $n \geq 7$ then $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic.

Case 2. n is even.

By [20, Proposition 2.5] we have $CCC(G) = 2K_1 \sqcup K_{2n-1}$. Therefore, $|V(CCC(G))| = 2n + 1$. Using (4), we get

$$(27) \quad E(K_{|V(CCC(G))}) = LE^+(K_{|V(CCC(G))}) = LE(K_{|V(CCC(G))}) = 4n.$$

By Theorem 3.5 we get

$$E(CCC(G)) = 4n - 4 < 4n.$$

Therefore, $CCC(G)$ is neither hyperenergetic nor borderenergetic.

If $n = 2$ then, by Theorem 4.5 and Theorem 3.5, we get

$$LE^+(CCC(G)) < LE(CCC(G)) = \frac{36}{5} < 8 = LE(K_{|V(CCC(G))}).$$

Therefore, $CCC(G)$ is neither L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic. Thus, if $n = 2$ then $CCC(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-border-energetic, Q-hyperenergetic nor Q-borderenergetic.

If $n \geq 4$ then, by Theorem 4.5 and Theorem 3.5, we get

$$\frac{4(2n - 1)(2n - 2)}{2n + 1} = LE^+(CCC(G)) < LE(CCC(G)).$$

We have

$$4n - \frac{4(2n - 1)(2n - 2)}{2n + 1} = -\frac{4(2n^2 - 7n + 2)}{2n + 1} = -\frac{4(2n(n - 4) + n + 2)}{2n + 1} < 0.$$

Therefore, $LE^+(K_{|V(CCC(G))}) = 4n < \frac{4(2n-1)(2n-2)}{2n+1} = LE^+(CCC(G)) < LE(CCC(G))$ and so $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither L-borderenergetic nor Q-borderenergetic. Thus, if $n \geq 4$ then $CCC(G)$ is L-hyperenergetic and Q-hyperenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic nor Q-borderenergetic. \square

We conclude this paper with the following characterization of commuting conjugacy class graph.

Theorem 5.6. *Let G be a finite non-abelian group. Then*

- (i) $CCC(G)$ is neither hyperenergetic, borderenergetic, L-hyperenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic if G is isomorphic to $D_8, D_{12}, D_{2n}(n \text{ is odd}), Q_8, Q_{12}, Q_{16}, U_{(2,6)}, U_{(n,2)}, U_{(n,3)}, U_{(n,4)}(n \geq 2), V_{16}, SD_{16}$ or SD_{24} .
- (ii) $CCC(G)$ is L-borderenergetic but neither hyperenergetic, borderenergetic, L-borderenergetic, Q-hyperenergetic nor Q-borderenergetic if G is isomorphic to Q_{20} or $U_{(2,5)}$.

- (iii) $CCC(G)$ is L -hyperenergetic but neither hyperenergetic, borderenergetic, L -borderenergetic, Q -hyperenergetic nor Q -borderenergetic if G is isomorphic to D_{16} , D_{20} , D_{24} , D_{28} , Q_{24} , Q_{28} , $U_{(3,5)}$, $U_{(3,6)}$, $U_{(2,7)}$, V_{24} or V_{32} .
- (iv) $CCC(G)$ is L -hyperenergetic and Q -borderenergetic but neither hyperenergetic, borderenergetic, L -borderenergetic nor Q -hyperenergetic if G is isomorphic to SD_{40} .
- (v) $CCC(G)$ is L -hyperenergetic and Q -hyperenergetic but neither hyperenergetic, borderenergetic, L -borderenergetic nor Q -borderenergetic if G is isomorphic to D_{2n} (n is even, $n \geq 16$), Q_{4m} ($m \geq 8$), $U_{(n,5)}$ ($n \geq 4$), $U_{(n,6)}$ ($n \geq 4$), $U_{(n,7)}$ ($n \geq 3$), $U_{(n,m)}$ ($n \geq 2$ and $m \geq 8$), V_{8n} ($n \geq 5$), SD_{32} or SD_{8n} ($n \geq 6$).

Theorem 5.7. *Let G be a finite non-abelian group. Then*

- (i) *If G is isomorphic to D_{2n} , Q_{4m} , $U_{(n,m)}$, V_{8n} or SD_{8n} then $CCC(G)$ is neither hyperenergetic nor borderenergetic.*
- (ii) *If G is isomorphic to D_{2n} (n is even, $n \geq 8$), Q_{4m} ($m \geq 6$), $U_{(n,5)}$ ($n \geq 3$), $U_{(n,6)}$ ($n \geq 3$), $U_{(n,m)}$ ($n \geq 2$ and $m \geq 7$), V_{8n} ($n \geq 3$) or SD_{8n} ($n \geq 4$) then $CCC(G)$ is L -hyperenergetic*
- (iii) *If G is isomorphic to Q_{20} or $U_{(2,5)}$ then $CCC(G)$ is L -borderenergetic.*
- (iv) *If G is isomorphic to D_{2n} (n is even, $n \geq 16$), Q_{4m} ($m \geq 8$), $U_{(n,5)}$ ($n \geq 4$), $U_{(n,6)}$ ($n \geq 4$), $U_{(n,7)}$ ($n \geq 3$), $U_{(n,m)}$ ($n \geq 2$ and $m \geq 8$), V_{8n} ($n \geq 5$), SD_{32} or SD_{8n} ($n \geq 6$) then $CCC(G)$ is Q -hyperenergetic.*
- (v) *If G is isomorphic to SD_{40} then $CCC(G)$ is Q -borderenergetic.*

6. ACKNOWLEDGMENT

The authors would like to thank the referee for his/her valuable comments. The first author is thankful to Council of Scientific and Industrial Research for the fellowship (File No. 09/796(0094)/2019-EMR-I).

REFERENCES

- [1] N. M. M. Abreu, C. T. M. Vinagre, A. S. Bonifácio and I. Gutman, *The Laplacian energy of some Laplacian integral graph*, MATCH Commun. Math. Comput. Chem., **60** (2008) 447-460.
- [2] K. Balińska, D. Cvetković, Z. Radosavljević, S. Simić and D. Stevanović, *A survey on integral graphs*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat., **13** (2003) 42-65.
- [3] R. Brauer and K. A. Fowler, *On groups of even order*, Ann. of Math., **62** No. 2 (1955) 565-583.
- [4] D. Cvetković, P. Rowlinson, S. Simić, *Signless Laplacian of finite graphs*, Linear Algebra Appl., **423** (2007) 155-171.
- [5] P. Dutta, B. Bagchi and R. K. Nath, *Various energies of commuting graphs of finite nonabelian groups*, Khayyam J. Math., **6** No. 1 (2020) 27-45.
- [6] J. Dutta and R. K. Nath, *Spectrum of commuting graphs of some classes of finite groups*, Matematika, **33** No. 1 (2017) 87-95.

- [7] J. Dutta and R. K. Nath, *Finite groups whose commuting graphs are integral*, Mat. Vesnik, **69** No. 3 (2017) 226-230.
- [8] J. Dutta and R. K. Nath, *Laplacian and signless Laplacian spectrum of commuting graphs of finite groups*, Khayyam J. Math., **4** No. 1 (2018) 77-87.
- [9] W. N. T. Fasfous, R. K. Nath and R. Sharafadini, *Various spectra and energies of commuting graphs of finite rings*, Hacet. J. Math. Stat., **49** No. 6 (2020) 1915-1925.
- [10] S.C. Gong, X. Li, G.H. Xu, I. Gutman and B. Furtula, *Borderenergetic graphs*, MATCH Commun. Math. Comput. Chem., **74** (2015) 321-332.
- [11] I. Gutman, *Hyperenergetic molecular graphs*, J. Serb. Chem. Soc., **64** (1999) 199-205.
- [12] I. Gutman, N. M. M. Abreu, C. T. M. Vinagre, A. S. Bonifácio and S. Radenković, *Relation between energy and Laplacian energy*, MATCH Commun. Math. Comput. Chem., **59** (2008) 343-354.
- [13] F. Harary and A. J. Schwenk, *Which graphs have integral spectra?*, Graphs and Combin., Lect. Notes Math., **406** (1974), Springer-Verlag, Berlin, 45-51.
- [14] M. Herzog, M. Longobardi and M. Maj, *On a commuting graph on conjugacy classes of groups*. Comm. Algebra, **37** No. 10 (2009) 3369-3387.
- [15] S. Kirkland, *Constructably Laplacian integral graphs*, Linear Algebra Appl., **423** (2007) 3-21.
- [16] J. Liu, and B. Liu, *On the relation between energy and Laplacian energy*, MATCH Commun. Math. Comput. Chem., **61** (2009) 403-406.
- [17] R. Merris, *Degree maximal graphs are Laplacian integral*, Linear Algebra Appl., **199** (1994) 381-389.
- [18] A. Mohammadian, A. Erfanian, D. G. M. Farrokhi and B. Wilkens, *Triangle-free commuting conjugacy class graphs*. J. Group Theory, **19** (2016) 1049-1061.
- [19] R. K. Nath, *Various spectra of commuting graphs n -centralizer finite groups*, International Journal of Engineering Science and Technology, accepted for publication.
- [20] M. A. Salahshour and A. R. Ashrafi, *Commuting conjugacy class graph of finite CA-groups*, Khayyam J. Math., **6** No. 1 (2020) 108-118.
- [21] S. K. Simić and Z. Stanić, *Q -integral graphs with edge-degrees at most five*, Discrete Math., **308** (2008) 4625-4634.
- [22] D. Stevanović, I. Stanković, and M. Milošević, *More on the relation between energy and Laplacian energy*, MATCH Commun. Math. Comput. Chem., **61** (2008) 395-401.
- [23] F. Tura, *L -borderenergetic graphs*, MATCH Commun. Math. Comput. Chem., **77** (2017) 37-44.
- [24] H.B. Walikar, H.S. Ramane and P.R. Hampiholi, *On the energy of a graph*, Graph Connections, Allied Publishers, New Delhi, 1999, pp. 120-123, Eds. R. Balakrishnan, H. M. Mulder, A. Vijayakumar.

Parthajit Bhowal

Department of Mathematical Sciences,

Tezpur university, Napaam

Assam, India.

bhowal.parthajit8@gmail.com

Rajat Kanti Nath

Department of Mathematical Sciences,

Tezpur university, Napaam

Assam, India.

rajatkantinath@yahoo.com