



Research Paper

A VARIATION OF δ -LIFTING AND δ -SUPPLEMENTED MODULES WITH RESPECT TO AN EQUIVALENCE RELATION

ESRA ÖZTÜRK SÖZEN*

ABSTRACT. In this paper we introduce Goldie* δ -supplemented modules as follows. A module M is called Goldie* δ -supplemented (briefly, G_δ^* -supplemented) if there exists a δ -supplement T of M for every submodule A of M such that $A\beta_\delta^*T$. We say that a module M is called Goldie* δ -lifting (briefly, G_δ^* -lifting) if there exists a direct summand D of M for every submodule A of M such that $A\beta_\delta^*D$. Note that the last concept given in [4] as a δ - H -supplemented module. We present fundamental properties of these modules. We indicate that these modules lie between δ -lifting and δ -supplemented modules. Also we prove that our modules coincide with some variations of δ -supplemented modules for δ -semiperfect modules.

1. INTRODUCTION

Throughout this study, R denotes an associative ring with identity and M denotes a unitary left R -module. The notations $A \leq M$ and $A \leq_\oplus M$ point that A is a *submodule* of M and

DOI: 10.29252/as.2021.2334

MSC(2010): 16D10, 16D70, 16D80

Keywords: Goldie* δ -lifting module, Goldie* δ -supplemented module.

Received: 09 March 2021, Accepted: 24 September 2021.

*Corresponding author

A is a *direct summand* of M , respectively. A submodule A is called *small* in M (denoted by $A \ll M$), if $A + X \neq M$ for any proper submodule X of M . A *supplement* submodule T of A in M provides that $A + T = M$ and $A \cap T \ll T$. A module M is called *supplemented* if every submodule of M has a supplement in M . M is called an *amply supplemented module* if for every submodule A of M with $M = A + B$ there exists a supplement submodule T of A contained in B . A module M is called *lifting* if, for any submodule A of M there exists a decomposition $M = X \oplus Y$ such that $X \leq A$ and $A \cap Y \ll Y$ [3]. And, M is called a \oplus -*supplemented module* if every submodule of M has a supplement which is a direct summand in M [10]. Also, in [10], a module M is called *H-supplemented* if for any $A \leq M$, there exists a submodule $D \leq_{\oplus} M$ such that $M = A + X$ if and only if $M = D + X$ for any $X \leq M$.

An *essential submodule* A of M provides that $\{0\}$ is the only submodule of M whose intersection with A is zero. A module M is called *singular (non-singular)* $Z(M) = M$ ($Z(M) = 0$) where $Z(M) = \{m \in M \mid \text{Ann}(m) \trianglelefteq R\}$. In [22] and in [9] new generalizations of small submodules, lifting modules and supplemented modules are introduced via singularity as follows. A δ -*small submodule* A of M is a submodule satisfying $A + X \neq M$ for every proper submodule X of M where $\frac{M}{X}$ is singular. We use the notation $\delta(M)$ for the sum of all δ -small submodules of M . Let φ be the class of all singular simple modules. For a module M , $\delta(M) = \cap\{N \leq M \mid \frac{M}{N} \in \varphi\}$. A submodule T of M is called a δ -*supplement* of A in M if $A + T = M$ and $A \cap T \ll_{\delta} T$. M is called δ -*supplemented* if every submodule of M has a δ -supplement in M . A module M is called δ -*lifting*, if for any submodule A of M there exists a decomposition $M = X + Y$ such that $X \leq A$ and $A \cap Y \ll_{\delta} Y$. If every submodule of M has a δ -supplement which is a direct summand of M , then M is called a \oplus - δ -*supplemented module* [18]. In [6], a module M is called δ -*H-supplemented*, if for any $A \leq M$ there exists a submodule $D \leq_{\oplus} M$ such that $M = A + X$ if and only if $M = D + X$ for every $X \leq M$ with $\frac{M}{X}$ is singular.

In [2], the authors defined an equivalence relation β^* and defined G^* -supplemented and G^* -lifting modules via this relation. Therefore, two new algebraic structures are contributed between lifting and supplemented modules. Owing to this fact, the open problem given as 'Is every H -supplemented module supplemented?' in [10] is handled. Thus, the following implications are obtained between some variations of supplemented modules and theirs such that:

$$\textit{lifting} \implies G^* - \textit{lifting} \implies H\textit{-supplemented} \implies G^*\textit{-supplemented} \implies \textit{supplemented}$$

and also we have the relation

$$\textit{lifting} \implies \textit{amply supplemented} \implies G^*\textit{-supplemented}$$

In [7], motivated by the equivalence relation β^* , the authors defined the relation β_δ^* as an extended alternative to β^* . For $A, B \leq M$, it is said that A *equivalents to B with respect to β_δ^** if and only if $\frac{A+B}{A} \ll_\delta \frac{M}{A}$ and $\frac{A+B}{B} \ll_\delta \frac{M}{B}$. A module M is called *principally Goldie*-\delta-lifting* (*principally Goldie*-\delta-supplemented*) if for any cyclic submodule Rm of M , there exists a direct summand D (δ -supplement T) in M such that $Rm\beta_\delta^*D$ ($Rm\beta_\delta^*T$). In [6], H -supplemented modules are designed according to the singularity. A module M is called δ - H -supplemented if every $A \leq M$ there exists a submodule $D \leq_\oplus M$ such that $M = A + X$ if and only if $M = D + X$ for any $X \leq M$ with $\frac{M}{X}$ is singular. Also we recommend [16] and [17] as a source to get more information about fundamental concepts used in this study.

In this study, inspired from [6] and from the equivalence relation β_δ^* given in [7], we generalize G^* -lifting and G^* -supplemented modules using singularity. We say that a module M is G_δ^* -lifting (G_δ^* -supplemented), if for any submodule A of M there exists a direct summand D (δ -supplement T) in M such that $A\beta_\delta^*D$ ($A\beta_\delta^*T$). By means of these concepts we obtain two new algebraic structures between δ -lifting and δ -supplemented modules. We indicate that δ - H -supplemented modules coincide with G_δ^* -supplemented modules. Also, we prove that our modules coincide with some variations of δ -supplemented modules for δ -semiperfect modules.

2. Preliminaries

Definition 2.1. Given submodules $A \leq B \leq M$, the inclusion $A \leq B$ is called δ -cosmall in M if $\frac{B}{A} \ll_\delta \frac{M}{A}$, denoted by $A \hookrightarrow_{\delta\text{-cs}} B$ [15].

Definition 2.2. Let M be a module and $A, B \leq M$. The submodule A is called β_δ^* equivalent to B (denoted by $A\beta_\delta^*B$) if $\frac{A+B}{A} \ll_\delta \frac{M}{A}$ and $\frac{A+B}{B} \ll_\delta \frac{M}{B}$.

It can be seen from [7, Lemma 3.2] that the relation given above is an equivalence relation.

Theorem 2.3. *Let $A, B \leq M$. Then the following statements are equivalent:*

- i. $A\beta_\delta^*B$
- ii. $A \hookrightarrow_{\delta\text{-cs}} A + B$ and $B \hookrightarrow_{\delta\text{-cs}} A + B$.
- iii. For every $X \leq M$ with $\frac{M}{X}$ is singular, if $A + B + X = M$ then $A + X = M$ then $B + X = M$.
- iv. If $X \leq M$ with $\frac{M}{X}$ is singular and $A + X = M$ then $B + X = M$ and, if $X \leq M$ with $\frac{M}{X}$ is singular and $B + X = M$ then $A + X = M$.

Proof. (i) \implies (ii) Let $A\beta_\delta^*B$. Therefore, we have $\frac{A+B}{A} \ll_\delta \frac{M}{A}$ and $\frac{A+B}{B} \ll_\delta \frac{M}{B}$, that is, $A \hookrightarrow_{\delta\text{-cs}} A + B$ and $B \hookrightarrow_{\delta\text{-cs}} A + B$.

(ii) \implies (iii) By assumption, it can be written that $\frac{A+B}{B} + \frac{X+B}{B} = \frac{M}{B}$. As $\frac{\frac{M}{X}}{\frac{B+X}{X}} \cong \frac{M}{B+X}$ is singular, we have $B + X = M$ is obtained. By the same way $A + X = M$ can be verified.

(iii) \iff (iv) Let $A + X = M$ for $X \leq M$ with $\frac{M}{X}$ is singular. By hypothesis, we get $B + X = B$ because $A + B + X = M$. Similarly, $A + X = M$ can be shown whenever $B + X = M$ for $X \leq M$ with $\frac{M}{X}$ is singular. Conversely, let $A + B + X = M$ such that $\frac{M}{X}$ is singular. Since $A + (B + X) = M$. and $\frac{M}{B+X}$ is singular, then $B + (B + X) = B + X = M$ is obtained from the assumption. Similarly, $A + X = M$ is shown. \square

Corollary 2.4. *Let $A, B \leq M$ such that $A \leq X + B$ and $B \leq Y + A$, where $X, Y \ll_{\delta} M$. Then $A\beta_{\delta}^*B$.*

Proof. Let $A + B + T = M$ for $T \leq M$ with $\frac{M}{T}$ is singular. Since $A \leq X + B$, then we have $(X + B) + B + T = X + B + T = M$. It follows that $B + T = M$ as $X \ll_{\delta} M$ and $\frac{M}{B+T}$ is singular as a factor module of a singular module $\frac{M}{T}$. Moreover, using the fact $Y \ll_{\delta} M$, it can be shown that $A + T = M$ and so, $A\beta_{\delta}^*B$ is obtained from Theorem 2.3. \square

Proposition 2.5. *If $A, B, X \leq M$ such that $M = A + X = B + X$, $B \cap X \leq A \cap X$ and $B \hookrightarrow_{\delta\text{-cs}} A + B$, then $A \hookrightarrow_{\delta\text{-cs}} A + B$, so $A\beta_{\delta}^*B$.*

Proof. It can be proved similar to that of [2, Proposition 2.5] using [22, Lemma 1.2]. \square

Proposition 2.6. *Let $P, T \leq M$ where P is maximal such that $\frac{M}{P} \in \varphi$.*

- i. *Let $A, B \leq M$ such that $A + B = M$, B is proper in M with $\frac{M}{B}$ is singular and $T\beta_{\delta}^*A$. Then T is not contained in B .*
- ii. *If $T\beta_{\delta}^*K$ and $T \leq P$, then $K \leq P$.*
- iii. *If $T\beta_{\delta}^*P$, then $T \leq P$. And, if $T\beta_{\delta}^*K$ then $T \leq \delta(M)$ if and only if $Y \leq \delta(M)$.*

Proof. i. Assume that $T \leq B$. By assumption, we have $A + B + T = M$. Then, $B + T = M$ is obtained from Theorem 2.3 since $\frac{M}{B}$ is singular. Hence, we get the contradiction $B = M$.

ii. Suppose that K is not contained in P . By maximality of P we have $K + P = M$ and so, $T + K + P = M$. As $\frac{M}{P} \in \varphi$ and $T\beta_{\delta}^*K$, we have $T + P = M$ from Theorem 2.3. Thus, $P = M$ is obtained which is a contradiction.

iii. By (ii), taking P instead of K , we get $T \leq P$ as $T\beta_{\delta}^*P$ and $P \leq P$. \square

Proposition 2.7. *Let $A, B, C, D \leq M$ such that $A\beta_{\delta}^*C$ and $C\beta_{\delta}^*D$. Then, $(A+C)\beta_{\delta}^*(B+D)$ and $(A+D)\beta_{\delta}^*(B+C)$.*

Proof. Let $X \leq M$ with $\frac{M}{X}$ is singular and $(A + C) + (B + D) + X = M$. Then we have $C + B + D + X = M$ and $A + C + D + X = M$ as $\frac{\frac{M}{X}}{C+D+X} \cong \frac{M}{C+D+X}$ is singular and $A\beta_\delta^*B$. Following, $B + D + X = M$ and $A + C + X = M$ as $\frac{\frac{M}{X}}{B+X}, \frac{M}{A+X}$ is singular and $C\beta_\delta^*D$. Hence, $(A + C)\beta_\delta^*(B + D)$ is obtained. Similarly, $(A + D)\beta_\delta^*(B + C)$ can be shown from the symmetry of β_δ^* . \square

Corollary 2.8. *Let $A, B_1, B_2, \dots, B_n \leq M$. If $A\beta_\delta^*B_i$ for each $i = 1, 2, \dots, n$, then $A\beta_\delta^*(B_1 + B_2 + \dots + B_n)$.*

Remark 2.9. The result given in Proposition 2.7 can not be extended to infinite sums. Let us consider that the \mathbb{Z} -module Q . It is a known fact that $\delta(\mathbb{Q}) = \mathbb{Q} = \sum_{n \in \mathbb{Z}^+} \frac{1}{n}\mathbb{Z}$ where $\frac{1}{n}\mathbb{Z} \ll_\delta \mathbb{Q}$ for each integer n . Clearly, $\frac{1}{n}\mathbb{Z} \beta_\delta^*0$ for each integer n . If the contrast of the claim would be true, then $\sum_{n \in \mathbb{Z}^+} \frac{1}{n}\mathbb{Z} \beta_\delta^*0 = \mathbb{Q} \beta_\delta^*0$ and so, $\mathbb{Q} \ll_\delta \mathbb{Q}$ is a contradiction.

Definition 2.10. Let $A \leq M$. Then $\beta_\delta^*(A) = \Sigma\{N \leq M \mid A\beta_\delta^*N\}$.

Note that $\beta_\delta^*(0) = \delta(M)$. On the other hand, let $A \leq P$ where $\frac{M}{P} \in \varphi$ which is the set of all singular simple modules. If $A\beta_\delta^*N$, then $N \leq P$ from Proposition 2.6. Hence, $\beta_\delta^*(A) \leq P$. Also, if $A\beta_\delta^*B$, then $\beta_\delta^*(A) = \beta_\delta^*(B)$.

3. GOLDIE $_\delta^*$ -LIFTING MODULES AND GOLDIE $_\delta^*$ -SUPPLEMENTED MODULES

Definition 3.1. A module M is called Goldie $_\delta^*$ -lifting (briefly, G_δ^* -lifting) if and only if for each $A \leq M$ there exists a direct summand D of M such that $A\beta_\delta^*D$.

Recall from [6] that a module M is called δ - H -supplemented if for every submodule A of M there exists a direct summand D of M such that $M = A + X$ if and only if $M = D + X$ for any $X \leq M$ with $\frac{M}{X}$ is singular. Let us indicate that this concept is the same with the definition given above. In view of brevity, we will use the term of G_δ^* -lifting for a this type of module. As it is possible to see the other fundamental properties of them in [6], we will omit them and give another ones.

Definition 3.2. A module M is called Goldie $_\delta^*$ -supplemented (briefly, G_δ^* -supplemented) if and only if for each $A \leq M$ there exists a δ -supplement D of M such that $A\beta_\delta^*D$.

Note that if M is a singular module or M has no projective submodule, then the concepts of being G^* -supplemented (G^* -lifting) and G_δ^* -supplemented (G_δ^* -lifting) coincide. In particular, a \mathbb{Z} -module M is G^* -supplemented (G^* -lifting) if and only if M is G_δ^* -supplemented (G_δ^* -lifting).

Proposition 3.3. *Let M be a δ -hollow module. Then, M is G_δ^* -lifting.*

Proof. Let X be any submodule of M . From the assumption, $X \ll_\delta M$. Therefore, $X\beta_\delta^*0$ is obtained where $\{0\}$ is a direct summand of M . Hence, M is G_δ^* -lifting. \square

Proposition 3.4. *Every semisimple module is G_δ^* -lifting.*

Proof. Let $A \leq M$. As M is semisimple, there exists a submodule B of M such that $M = A \oplus B$. From the symmetry of β_δ^* we have $A\beta_\delta^*B$. Hence, M is G_δ^* -lifting. \square

Proposition 3.5. *Let M be a G_δ^* -lifting module and $A \leq M$. If $\frac{A+D}{A} \leq_\oplus \frac{M}{A}$ for any $D \leq_\oplus M$, then $\frac{M}{A}$ is G_δ^* -lifting.*

Proof. Let $\frac{X}{A} \leq \frac{M}{A}$. Since M is G_δ^* -lifting, then there exists a decomposition $M = D \oplus D'$ such that $X\beta_\delta^*D$. Let $\pi : M \rightarrow \frac{M}{A}$ be the canonical epimorphism. Then, $\pi(X)\beta_\delta^*\pi(D)$ and so, $\frac{X}{A}\beta_\delta^*\frac{D+A}{A}$ is obtained. Hence, M is G_δ^* -lifting from the hypothesis. \square

If the sum of any two direct summands of M is a direct summand, then M has the summand sum property.

Proposition 3.6. *Let M be a G_δ^* -lifting module. If M has the summand sum property, then any direct summand of M is G_δ^* -lifting module.*

Proof. Let $D \leq_\oplus M$. Then $M = D \oplus D'$ for some $D' \leq M$. We will verify that $\frac{M}{D'}$ is G_δ^* -lifting. With this aim, we will show that for any $X \leq_\oplus M$, $\frac{X+D'}{D'} \leq_\oplus \frac{M}{D'}$. From the assumption, as D' and X are direct summands of M , then $X + D' \leq_\oplus M$. Therefore, there is a submodule T of M such that $M = (X + D') \oplus T$. It follows that, $\frac{M}{D'} = \frac{X+D'}{D'} + \frac{T+D'}{D'}$. Moreover, we get $(X + D') \cap (T + D') = [(X + D') \cap T] + D' = 0 + D' = D'$ from modularity. Thus, $\frac{M}{D'} = \frac{X+D'}{D'} \oplus \frac{T+D'}{D'}$. Hence, $D \cong \frac{M}{D'}$ is G_δ^* -lifting from Proposition 3.5. \square

Proposition 3.7. *Let M be a π -projective module and let us consider the following statements.*

- (1) M is \oplus - δ -supplemented.
- (2) M is G_δ^* -lifting.

Then (1) \implies (2) holds. In particular, if M is singular, then the converse is provided.

Proof. (1) \implies (2) : Let A be any submodule of M . By (1), there exists a δ -supplement T of M which is a direct summand of M such that

$$M = A + T, \quad A \cap T \ll_\delta T \quad \text{and} \quad M = T \oplus T'$$

As M is π -projective, there is a submodule $X \leq A$ provided that $M = X \oplus T$. Clearly, $\frac{X+A}{A} \ll_{\delta} \frac{M}{A}$. Moreover, as $A = X \oplus (A \cap T)$ from modularity and $A \cap T \ll_{\delta} T$, we obtain $\frac{A}{X} = \frac{X+A}{X} \ll_{\delta} \frac{M}{X}$, that is, $A\beta_{\delta}^* X$ where $X \leq_{\oplus} M$. Hence, M is G_{δ}^* -lifting.

Let M be a singular G_{δ}^* -lifting module and $A \leq M$. Then, there exists a direct summand D of M such that $M = D \oplus D'$ for some $D' \leq M$ and $A\beta_{\delta}^* D$. Therefore, $\frac{D+A}{D} \ll_{\delta} \frac{M}{D}$ and $\frac{D+A}{A} \ll_{\delta} \frac{M}{A}$. As $\frac{M}{A} = \frac{D+A}{A} + \frac{D'+A}{A}$ and $\frac{M}{D+A} \cong \frac{\frac{M}{A}}{\frac{D+A}{A}}$ is singular, then $M = A + D'$ is obtained. Now, it remains to show that $A \cap D' \ll_{\delta} D'$. Let $(A \cap D') + B = D'$ where $\frac{D'}{B}$ is singular. Thus, $M = (A \cap D') + B + D = A + B + D$ and so $\frac{M}{D} = \frac{A+D}{D} + \frac{B+D}{D}$ is obtained. As $\frac{D+A}{D} \ll_{\delta} \frac{M}{D}$ and M is singular, we have $M = B + D$ and so $D' = M \cap D' = (B + D) \cap D' = B + (D \cap D') = B$ from modularity. This completes the proof. Hence, M is a \oplus - δ -supplemented module. \square

Theorem 3.8. *Let M be a singular G_{δ}^* -lifting module. Then M is δ -supplemented.*

Proof. Let $A \leq M$. By hypothesis, there exists a direct summand D of M such that $M = D \oplus D'$ and $A\beta_{\delta}^* D$. Then, $\frac{A+D}{D} \ll_{\delta} \frac{M}{D}$, $\frac{A+D}{A} \ll_{\delta} \frac{M}{A}$ and we have $\frac{M}{A} = \frac{A+D}{A} + \frac{A+D'}{A}$. As M is singular, $M = A + D'$ is obtained. To complete the proof it must be shown that $A \cap D' \ll_{\delta} D'$. Let $A \cap D' + B = D'$ with $\frac{D'}{B}$ is singular. Then, $M = D' + D = [A \cap D' + B] + D = A + B + D$ and so, $\frac{M}{D} = \frac{A+D}{D} + \frac{B+D}{D}$ is obtained. By hypothesis, we get $M = B + D$ and so, $D' = B$ by modularity. Hence, M is δ -supplemented. \square

Theorem 3.9. *Let M be a π -projective module. If M is G_{δ}^* -supplemented, then it is G_{δ}^* -lifting.*

Proof. Let $A \leq M$. Then, there exists a δ -supplement T of M such that $A\beta_{\delta}^* T$. Assume that T is a δ -supplement of W in M , that is $W + T = M$ and $W \cap T \ll_{\delta} T$. As M is π -projective, there exists a direct summand T' of M contained in T such that $M = T' \oplus W$. Now, we aim to verify that $A\beta_{\delta}^* T'$. It is clear that, $\frac{A+T'}{A} \leq \frac{A+T}{A} \ll_{\delta} \frac{M}{A}$ by [22, Lemma 1.3]. In the remaining part of the proof we will show that $\frac{A+T'}{T'} \ll_{\delta} \frac{M}{T'}$. Suppose that, $\frac{A+T'}{T'} + \frac{B}{T'} = \frac{M}{T'}$ with $\frac{M}{B}$ is singular. Then $M = A + T' + B = A + T + B$ and so $\frac{M}{T} = \frac{A+T}{T} + \frac{B+T}{T}$. As $A\beta_{\delta}^* T$ and $\frac{\frac{M}{B}}{\frac{B+T}{B}} \cong \frac{M}{B+T}$ is singular, then $M = B + T$. Moreover, we have $T = T' \oplus (W \cap T)$ by modularity. Thus, $M = B + T = B + [T' + (W \cap T)] = (B + T') + W \cap T$. As $W \cap T \ll_{\delta} M$ and $\frac{M}{B+T}$ is singular, $M = B + T'$ and so, $M = B$ is obtained due to the fact that $T' \leq B$. Hence the proof is completed. \square

Proposition 3.10. *Let M be a Noetherian module which has the summand sum property. Then M is principally G_{δ}^* -lifting if and only if M is G_{δ}^* -lifting.*

Proof. The sufficiency is clear. For the necessity, let $A \leq M$. As M is Noetherian, A is finitely generated and so $A = Rx_1 + Rx_2 + \dots + Rx_n$ for some $x_1, x_2, \dots, x_n \in M$. Since M is principally G_δ^* -lifting, there exists direct summands D_1, D_2, \dots, D_n of M such that $Rx_1\beta_\delta^*D_1, Rx_2\beta_\delta^*D_2, \dots, Rx_n\beta_\delta^*D_n$. Then, $A\beta_\delta^*D = D_1 + D_2 + \dots + D_n$ where $D \leq_\oplus M$ since M has the summand sum property. Hence, M is G_δ^* -lifting. \square

Proposition 3.11. *Let M be a module and $A \leq M$ such that $A = C + S$ where C is cyclic in M and $S \ll_\delta M$ for any $A \leq M$. Then M is principally G_δ^* -lifting and G_δ^* -lifting.*

Proof. Let $A = C + S$ for a cyclic submodule C of M and $S \ll_\delta M$. As M is principally G_δ^* -lifting, then a direct summand D of M corresponds to C such that $C\beta_\delta^*D$. Therefore, $A = (C + S)\beta_\delta^*D$ by [7, Lemma 3.6] which implies M is G_δ^* -lifting. The sufficiency is clear from implications. \square

Proposition 3.12. *Let M be a module and A be any submodule of M . If there exists a δ -supplement (direct summand) T and a δ -small submodule S of M such that $A + S = T + S$, then M is a G_δ^* -supplemented (G_δ^* -lifting) module.*

Proof. From assumption, it remains to show that $A\beta_\delta^*T$. Since $A \leq A + S = T + S$, $T \leq T + S = A + S$ and $S \ll_\delta M$, then we have $A\beta_\delta^*T$ from Corollary 2.4. \square

Corollary 3.13. *Let M be a module and A be any submodule of M . If there exists a δ -supplement T and a δ -small submodule S of M such that $A = T + S$, then M is a G_δ^* -supplemented module.*

Theorem 3.14. *Let M be a module and consider the statements given below,*

- a. M is δ -lifting.
- b. M is G_δ^* -lifting.
- c. M is δ -H-supplemented.
- d. M is G_δ^* -supplemented.

Then, (a) \implies (b) \iff (c) \implies (d).

Proof. (a) \implies (b) Let M be a δ -lifting module. Then, there exists a direct summand D for any submodule A of M satisfying $\frac{A}{D} \ll_\delta \frac{M}{D}$. Therefore, it can be written that $\frac{A+D}{D} \ll_\delta \frac{M}{D}$ and $\frac{A+D}{A} = 0 \ll_\delta \frac{M}{A}$ which implies $A\beta_\delta^*D$. Hence, we obtain the existence of a direct summand D for every submodule A of M such that $A\beta_\delta^*D$, that is, M is G_δ^* -lifting.

(b) \iff (c) This fact is clear from [6, Lemma 2.2].

(c) \implies (d) is clear because every direct summand is a δ -supplement. \square

Proposition 3.15. *Let M be a module whose submodules are of δ -supplements which are relatively projective direct summands of M . Then, M is G_δ^* -lifting.*

Proof. Let $A \leq M$. Then, there is a δ -supplement T of M such that $M = A + T$, $A \cap T' \ll_\delta T'$ and $M = T \oplus T'$ where T, T' are relatively projective. It follows that $M = B \oplus T$ for some $B \leq A$ since T' is T -projective from [10, Lemma 4.47]. Therefore, M is δ -lifting. Hence, M is G_δ^* -lifting from Theorem 3.14. \square

Proposition 3.16. *Let M be a π -projective and singular module. Then the following statements hold equivalently.*

- (1) M is δ -lifting.
- (2) M is G_δ^* -lifting.
- (3) M is \oplus - δ -supplemented module.

Proof. (1) \implies (2) : is clear from Theorem 3.14.

(2) \iff (3) : is clear from Proposition 3.7.

(3) \implies (1) : Let $A \leq M$. From assumption, there exists a direct summand D of M such that $M = D \oplus D'$, $A + D = M$ and $A \cap D \ll_\delta D$. On the other hand, as M is π -projective and $D \leq_\oplus M$, then there exists a direct summand A' of M contained A such that $M = A' \oplus D$ from [3, 4.14(1)]. Thus, for every $A \leq M$, there exists a decomposition $M = A' \oplus D$ such that $A' \leq A$ and $A \cap D \ll_\delta D$. Hence M is δ -lifting. \square

Proposition 3.17. *Let M be a singular π -projective module. Then, M is G_δ^* -lifting if and only if every submodule of M is a direct sum of a direct summand of M and a δ -small submodule of M .*

Proof. (\implies) Let M be a G_δ^* -lifting module, then M is a δ -lifting module from Proposition 3.16. Then for any $A \leq M$, there exists a decomposition $M = D \oplus D'$ such that $D \leq A$ and $A \cap D' \ll_\delta M$. It follows that $A = D \oplus (A \cap D')$ where $D \leq_\oplus M$ and $S = A \cap D' \ll_\delta M$.

(\impliedby) For the necessity, it can be said that M is δ -lifting from [9, Lemma 2.3(b)]. Hence, M is G_δ^* -lifting by Theorem 3.14. \square

Proposition 3.18. *Let R be a left non-singular ring, M be a left G_δ^* -supplemented R -module and P be a maximal submodule of M with $\frac{M}{P}$ is singular. If T is a δ -supplement of P with $\frac{M}{T}$ is singular, then $P = S + (P \cap T)$, where S is a δ -supplement of T and T is δ -local.*

Proof. Let M be a G_δ^* -supplemented module. Then, there exists a δ -supplement submodule S corresponding to P satisfying $P\beta_\delta^*S$. By hypothesis, T is a δ -supplement of S . Therefore, we have $P = S + (P \cap T)$ from [7, Theorem 3.7]. Moreover, since T is a δ -supplement submodule of the maximal submodule P , then T is δ -local or semisimple projective from [19, Lemma 2.22]. If T is semisimple projective, then $T \ll_\delta T \leq M$. On the other hand, as T is a δ -supplement of P in M , $P + T = M$ and $P \cap T \ll_\delta T$. Since $T \ll_\delta M$ and $\frac{M}{P}$ is singular, then $P = M$ is got which contradicts with maximality of P in M . Hence, it forces T to be δ -local. \square

Example 3.19. Let $R = \frac{\mathbb{Z}}{8\mathbb{Z}}$ and $M = \frac{\mathbb{Z}}{2\mathbb{Z}} \oplus \frac{\mathbb{Z}}{4\mathbb{Z}} \oplus \frac{\mathbb{Z}}{8\mathbb{Z}}$. It is a known fact from [9, Example 2.2(2)] that M is not a δ -lifting module. On the other hand, M is a G_δ^* -lifting module as it is G^* -lifting [2, Example 3.9(i)].

Example 3.20. Let $M = \mathbb{F} \oplus \mathbb{F}$ where \mathbb{F} is a quotient field of a DVR R which is not complete. Then it can be seen that clearly M is a δ -supplemented module which is not G_δ^* -supplemented from [3, Example 23.7] and [2, Example 3.9(iii)].

Definition 3.21. A δ -supplemented module M is called *strongly \oplus - δ -supplemented* if every δ -supplement submodule of M is a direct summand of M .

Clearly, every δ -lifting module is strongly \oplus - δ -supplemented.

Proposition 3.22. *Let M be a module.*

- i. M is δ -lifting if and only if M is amply δ -supplemented and strongly \oplus - δ -supplemented.
- ii. If M is G_δ^* -supplemented and strongly \oplus - δ -supplemented, then M is G_δ^* -lifting.

Proof. i. The implication is clear from [1, Proposition 4.2] and [12, Lemma 2.3].

ii. Let A be any submodule of M . By assumption, there is a δ -supplement submodule X of M provided that $A\beta_\delta^*X$. As M is strongly \oplus - δ -supplemented X is a direct summand of M . Hence, M is G_δ^* -lifting. \square

Proposition 3.23. *Let M be a module.*

- i. M is amply δ -supplemented.
- ii. For each $A \leq M$ there there is a δ -supplement T and a submodule X of M such that $M = T + X = A + X$, $T + X \leq A + X$ and $T \leftrightarrow_{\delta\text{-cs}} A + T$.
- iii. M is G_δ^* -supplemented.

Then, the condition given above implies that (i) \implies (ii) \implies (iii).

Proof. (i) \implies (ii) It is clear from [15, Theorem 3.7].

(ii) \implies (iii) By hypothesis, we have $A\beta_\delta^*T$ from Proposition 2.5. Hence, M is G_δ^* -supplemented. \square

Proposition 3.24. *Let M be a projective module. Then the following statements are equivalent:*

- i. M is δ -semiperfect.
- ii. M is δ -lifting.
- iii. M is \oplus - δ -supplemented.
- iv. M is amply δ -supplemented.
- v. M is δ -supplemented.
- vi. M is G_δ^* -supplemented.
- vii. M is G_δ^* -lifting.

Proof. It can be seen clearly via Theorem 3.14, Proposition 3.23 and [12, Lemma 2.4]. \square

The following hierarchy is valid for given modules below.

$$\delta\text{-lifting} \implies G_\delta^*\text{-lifting} \implies \text{principally } G_\delta^*\text{-lifting}$$

Now we will verify the converse implications given above are not provided.

Example 3.25. Let us consider \mathbb{Z} -module \mathbb{Q} . Since every finitely generated submodule of \mathbb{Z} -module \mathbb{Q} is δ -small in \mathbb{Q} , then ${}_Z\mathbb{Q}$ is a principally G_δ^* -lifting module. On the other hand, it is not G_δ^* -lifting as it is not δ -supplemented.

Example 3.26. Let $R = \mathbb{Z}$ and $M = \mathbb{Z}_2 \oplus \mathbb{Z}_8$. M is a G_δ^* -lifting module which is not δ -lifting [8].

4. ACKNOWLEDGMENTS

The authors wish to sincerely thank the referees for several useful comments.

REFERENCES

[1] C. Abdioğlu and S. Şahinkaya, *Some Results On δ -Semiperfect Rings and δ -Supplemented Modules*, Kyungpook Math. J., **55** No. 2 (2015) 289-300.

[2] G.F. Birkenmeier, F.T. Mutlu, C. Nebiyev, N. Sokmez and A. Tercan, *Goldie* Supplemented Modules*, Glas. Math. J., **52** No. 2 (2010) 41-52.

[3] J. Clark, C. Lomp, N. Vanaja and R. Wisbauer, *Lifting Modules: Supplements and Projectivity in Module Theory*, Birkhauser Verlag-Basel, Boston-Berlin, 2006.

- [4] F. Goodearl, *Ring Theory: Nonsingular Rings and Modules*, Crc Press, Marcel Dekker Inc., 1976.
- [5] A.T. Guroğlu and E.T. Meric, *Principally Goldie* Lifting Modules*, Ukr. Math. J., **70** No. 3 (2018) 1042-1051.
- [6] A.R.M. Hamzekolae, *H-supplemented Modules and Singularity*, J. Alg. Str. App., **7** No. 1 (2020) 49-57.
- [7] M. Hosseinpour, B. Ungor, Y. Talebi and A. Harmancı, *A Generalization of the Class of Principally Lifting Modules*, Rocky Mount. J. Math., **47** No. 5 (2017) 1539-1563.
- [8] H. İnankıl, S. Halcioğlu and A. Harmancı, *On A Class of Lifting Modules*, Viet. J. Math., **38** (2010) 189-201.
- [9] M.T. Koşan, *δ -Lifting and δ -Supplemented Modules*, Alg. Coll., **14** No. 1 (2007) 53-60.
- [10] S.H. Mohamed and B.J. Müller, *Continuous and Discrete Modules*, London Math. Soc. Lecture Note Ser. 147, Cambridge University Press, 1990.
- [11] M.J. Nematollahi, *Quasi δ -Discrete Modules*, Alg., Gr. and Geo., **28** No. 3 (2011) 259-274.
- [12] X. Nguyen, M. Koşan and Y. Zhou, *On δ -Semiperfect Modules*, Comm. Alg., **46** No. 11 (2018) 4965-4977.
- [13] A.C. Ozcan, *The Torsion Theory Cogenerated By δ -M-small Modules and GCO-modules*, Comm. Alg., **35** (2009) 623-633.
- [14] Y. Talebi and A.R.M. Hamzekolae, *Closed Weak δ -Supplemented Modules*, JP J. Alg., Num. Th. and App., **13** No. 2 (2009) 193-208.
- [15] Y. Talebi and B. Talae, *On δ -Coclosed Submodules*, Far East J. Math. Sci., **35** No. 1 (2009) 19-31.
- [16] Y. Talebi, R. Tribak and A.R.M. Hamzekolae, *On H-cofinitely Supplemented Modules.*, Bull. Iranian. Math. Soc., **39** No. 2 (2013) 325-346.
- [17] R. Tribak, Y. Talebi, A.R.M. Hamzekolae and S. Asgari, *\oplus -Supplemented Modules Relative to An Ideal*, Hacettepe J. Math. Stat., **45** No. 1 (2016) 107-120.
- [18] Y. Talebi and M.H. Pour, *On \oplus - δ -supplemented Modules*, J. Alg. Num. Th. Adv. and App., **1** No. 2 (2009) 89-97.
- [19] R. Tribak, *When Finitely Generated δ -Supplemented Modules are Supplemented*, Alg. Coll., **22** No. 1 (2015) 119-130.
- [20] D.K. Tutuncu, M.J. Nematollahi and Y. Talebi, *On H-Supplemented Modules*, Alg. Coll., **18** No. 1 (2011) 915-924.
- [21] R. Wisbauer, *Foundations of Module and Ring Theory: A handbook for study and research*, Gordon and Breach Science Publishing, London-Routledge, 1991.
- [22] Y. Zhou, *Generalizations of Perfect, Semiperfect and Semiregular Rings*, Alg. Coll., **7** No. 3 (2000) 305-318.

Esra Öztürk Sözen

Department of mathematics, Faculty of science and arts

Sinop university

Sinop, Turkey.

esozen@sinop.edu.tr