



Algebraic Structures and Their Applications Vol. 1 No. 1 (2014), pp 35-48.

STABILIZER TOPOLOGY OF HOOPS

R. A. BORZOOEI*, M. AALY KOLOGANI

Communicated by S. Alikhani

ABSTRACT. In this paper, we introduce the concepts of right, left and product stabilizers on hoops and study some properties and the relation between them. And we try to find that how they can be equal and investigate that under what condition they can be filter, implicative filter, fantastic and positive implicative filter. Also, we prove that right and product stabilizers are filters and if they are proper, then they are prime filters. Then by using the right stabilizers produce a basis for a topology on hoops. We show that the generated topology by this basis is Baire, connected, locally connected and separable and we investigate the other properties of this topology. Also, by the similar way, we introduce the right, left and product stabilizers on quotient hoops and introduce the quotient topology that is generated by them and investigate that under what condition this topology is Hausdorff space, T_0 or T_1 spaces.

MSC(2010): Primary:03G25 Secondary:06B99

Keywords: Hoop algebra, stabilizer topology, Baire space, connected, locally connected, separable topology.

Received: 12 July 2014, Accepted: 02 Nov 2014.

*Corresponding author

1. INTRODUCTION

Hoop-algebras are naturally ordered commutative residuated integral monoids, introduced by B. Bosbach in [2, 3] then study by J. R. Büchi and T. M. Owens in [6], a paper never published. In the last years, hoops theory was enriched with deep structure theorems(see [1, 2, 3]). Many of these results have a strong impact with fuzzy logic. Particularly, from the structure theorem of finite basic hoops([1, Corollary 2.10]) one obtains an elegant short proof of the completeness theorem for propositional basic logic(see [1, Theorem 3.8]), introduced by Hájek in [8]. The algebraic structures corresponding to Hájek's propositional (fuzzy) basic logic, BL-algebras, are particular cases of hoops. The main example of BL-algebras in interval $[0, 1]$ endowed with the structure induced by a t-norm. MV-algebras, product algebras and Gödel algebras are the most known classes of BL-algebras. Recent investigations are concerned with non-commutative generalizations for these structures.

A stabilizer is a part of a monoid acting on a set. Specifically, let A be a monoid operating on a set A and let X be a subset of S . The *stabilizer* of X , sometimes denoted $St(X)$ is the set of elements of a of A for which $a(X) \subseteq X$. The *strict stabilizer* is the set of $a \in A$ for which $a(X) = X$. In the other words, the stabilizer of X is the transporter of X to itself.

In this paper, we introduce the concept of right, left and product stabilizers on hoops. Then we use the right stabilizer of a hoop A and produce a basis for topology on A . We show that the generated topology by this basis is Baire, connected, locally connected and separable and we investigate the other properties of this topology.

2. PRELIMINARIES

In this section, we gather some basic notions relevant to hoop which will need in the next sections.

Definition 2.1. [1] A *hoop* is an algebra $(A, \odot, \rightarrow, 1)$ of type $(2, 2, 0)$ such that, for all $x, y, z \in A$:

(HP1) $(A, \odot, 1)$ is a commutative monoid,

(HP2) $x \rightarrow x = 1$,

(HP3) $(x \odot y) \rightarrow z = x \rightarrow (y \rightarrow z)$,

(HP4) $x \odot (x \rightarrow y) = y \odot (y \rightarrow x)$.

On hoop A we define $x \leq y$ if and only if $x \rightarrow y = 1$. It is easy to see that " \leq " is a partial order relation on A . A hoop A is *bounded* if there is an element $0 \in A$ such that $0 \leq x$, for all $x \in A$. We let $x^0 = 1$, $x^n = x^{n-1} \odot x$, for any $n \in \mathbb{N}$. Let A be a bounded hoop. We define a negation " $'$ " on A by, $x' = x \rightarrow 0$, for all $x \in A$. If $(x')' = x$, for all $x \in A$, then the bounded hoop A is said to have the *double negation property*, or (DNP) for short.

The following proposition provides some properties of hoop.

Proposition 2.2. [2, 3] *Let $(A, \odot, \rightarrow, 1)$ be a hoop. Then the following condition hold, for all $x, y, z, a \in A$:*

- (i) (A, \leq) is a \wedge -semilattice with $x \wedge y = x \odot (x \rightarrow y)$,
- (ii) $x \odot y \leq z$ if and only if $x \leq y \rightarrow z$,
- (iii) $x \odot y \leq x, y$,
- (iv) $x \leq y \rightarrow x$,
- (v) $x \rightarrow 1 = 1$,
- (vi) $1 \rightarrow x = x$,
- (vii) $x \odot (x \rightarrow y) \leq y$,
- (viii) $x \leq y$ implies $x \odot a \leq y \odot a$,
- (ix) $x \leq y$ implies $z \rightarrow x \leq z \rightarrow y$,
- (x) $x \leq y$ implies $y \rightarrow z \leq x \rightarrow z$,
- (xi) $((x \rightarrow y) \rightarrow y) \rightarrow y = x \rightarrow y$,
- (xii) $(x \rightarrow y) \odot (y \rightarrow z) \leq (x \rightarrow z)$.

Proposition 2.3. [7] *Let A be a bounded hoop. Then the following conditions hold, for all $x, y \in A$:*

- (i) $x \leq x''$,
- (ii) $x \odot x' = 0$,
- (iii) if $x = x''$, then $x \rightarrow y = y' \rightarrow x'$,
- (iv) $x = x''$ if and only if $(x \rightarrow y) \rightarrow y = (y \rightarrow x) \rightarrow x$.

Proposition 2.4. [7] *Let A be a hoop and for any $x, y \in A$, we define, $x \sqcup y = ((x \rightarrow y) \rightarrow y) \wedge ((y \rightarrow x) \rightarrow x)$. Then the following conditions are equivalent:*

- (i) \sqcup is associative operation on A ,
- (ii) $x \leq y$ implies $x \sqcup z \leq y \sqcup z$, for all $x, y, z \in A$,
- (iii) $x \sqcup (y \wedge z) \leq (x \sqcup y) \wedge (x \sqcup z)$, for all $x, y, z \in A$,
- (iv) \sqcup is the join operation on A .

Definition 2.5. A hoop A is called a \sqcup -hoop, if \sqcup is a join operation on A .

Remark 2.6. [7, Remark 2.4] \sqcup -hoop (A, \sqcup, \wedge) is a distributive lattice.

Proposition 2.7. [7] *Let A be a \sqcup -hoop. Then, for all $x, y, z \in A$,*

- (i) $(x \sqcup y) \rightarrow z = (x \rightarrow z) \wedge (y \rightarrow z)$,
- (ii) $x \odot (y \sqcup z) = (x \odot y) \sqcup (x \odot z)$.

Definition 2.8. [7] Let A be a hoop. A non-empty subset F of A is called a *filter* of A if,

- (F1) $x, y \in F$ implies $x \odot y \in F$,
- (F2) $x \leq y$ and $x \in F$ imply $y \in F$, for any $x, y \in A$.

We use $\mathcal{F}(A)$ to denote the set of all filters of A . Clearly, $1 \in F$, for all $F \in \mathcal{F}(A)$. $F \in \mathcal{F}(A)$ is called a *proper filter* if $F \neq A$. It can be easily seen that, if A is a bounded hoop, then a filter is proper if and only if it is not containing 0.

Proposition 2.9. [7] *Let A be a hoop and F be a non-empty subset of A . Then $F \in \mathcal{F}(A)$ if and only if $1 \in F$ and if $x, x \rightarrow y \in F$, then $y \in F$, for any $x, y \in A$.*

Let A be a hoop and $F \in \mathcal{F}(A)$. We define a binary relation \sim_F on A by $x \sim_F y$ if and only if $x \rightarrow y, y \rightarrow x \in F$, for any $x, y \in A$. Then \sim_F is a congruence relation on A . Let $A/F = \{\bar{x} \mid x \in A\}$, where $\bar{x} = \{y \in A \mid x \sim_F y\}$. Then the binary relation \leq on A/F which is defined by:

$$\bar{x} \leq \bar{y} \text{ if and only if } x \rightarrow y \in F.$$

is an order relation on A/F (see [15]). Now, let A be a hoop and $F \in \mathcal{F}(A)$. Then $(A/F, \otimes, \rightsquigarrow, 1_{A/F})$ is a hoop, where for any $x, y \in A$:

$$1_{A/F} = \bar{1}, \bar{x} \otimes \bar{y} = \overline{x \odot y}, \bar{x} \rightsquigarrow \bar{y} = \overline{x \rightarrow y}.$$

Definition 2.10. [9, 13] Consider X as a non-empty set. A mapping $\varphi : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ is called a *closure operator* on X , if the following holds, for all $Y, Z \in \mathcal{P}(X)$:

- (C1) $Y \subseteq \varphi(Y)$,
- (C2) $\varphi^2(Y) = \varphi(Y)$,
- (C3) $Y \subseteq Z$ implies $\varphi(Y) \subseteq \varphi(Z)$.

Definition 2.11. [9] A topological space (X, \mathcal{T}) is called *Baire space*, if for each countable collection of open dense sets U_n , their intersection $\bigcap U_n$ is dense.

Note: From now on, in this paper, we let A be a hoop, unless otherwise state.

3. STABILIZER OF HOOPS

In this section, we introduce the concept of right, left and product stabilizers on hoops and investigate some properties of them.

Definition 3.1. Let $X \subseteq A$. Then

- $St_r(X) = \{a \in A \mid a \rightarrow x = x, \forall x \in X\}$ is called the *right stabilizer* of X .
- $St_l(X) = \{a \in A \mid x \rightarrow a = a, \forall x \in X\}$ is called the *left stabilizer* of X .
- $St_{\odot}(X) = \{a \in A \mid a \odot x = x, \forall x \in X\}$ is called the *product stabilizer* of X .

Example 3.2. (i) Let $A = \{0, a, b, 1\}$ and operations \odot and \rightarrow on A are defined as follows:

\odot	0	a	b	1	\rightarrow	0	a	b	1
0	0	0	0	0	0	1	1	1	1
a	0	0	0	a	a	b	1	1	1
b	0	0	a	b	b	a	b	1	1
1	0	a	b	1	1	0	a	b	1

Routine calculations show that A with these operations is a bounded hoop and $St_r(\{a\}) = St_\odot(\{b\}) = St_l(\{b\}) = \{1\}$.

(ii) Let $A = \{0, a, b, 1\}$. Define \odot and \rightarrow on A as follows:

\odot	0	a	b	1	\rightarrow	0	a	b	1
0	0	0	0	0	0	1	1	1	1
a	0	0	a	a	a	a	1	1	1
b	0	a	b	b	b	0	a	1	1
1	0	a	b	1	1	0	a	b	1

Then $(A, \odot, \rightarrow, 1)$ is a bounded hoop. Let $X_1 = \{b\}$ and $X_2 = \{0, a, 1\}$. Then $St_l(X_1) = \{0, a, 1\}$, $St_\odot(X_1) = \{b, 1\}$ and $St_r(X_2) = \{b, 1\}$.

Note: $St_r(St_l(X))$ is called *right-left stabilizer of X* and we show it by $(St_l(X))_r$, for short.

The following proposition provides some properties of stabilizers of hoops.

Proposition 3.3. *For all $x, y \in A$ and $X, Y \subseteq A$, the following statements hold:*

- (i) $1 \in St_r(X) \cap St_l(X) \cap St_\odot(X)$,
- (ii) if $X \subseteq Y$, then $St_r(Y) \subseteq St_r(X)$, $St_l(Y) \subseteq St_l(X)$ and $St_\odot(Y) \subseteq St_\odot(X)$,
- (iii) $X \subseteq (St_r(X))_l \cap (St_l(X))_r$,
- (iv) $St_r(X) = ((St_r(X))_l)_r$ and $St_l(X) = ((St_l(X))_r)_l$,
- (v) $St_r(\bigcup_{i \in I} X_i) = \bigcap_{i \in I} (St_r(X_i))$, $St_l(\bigcup_{i \in I} X_i) = \bigcap_{i \in I} (St_l(X_i))$ and $St_\odot(\bigcup_{i \in I} X_i) = \bigcap_{i \in I} (St_\odot(X_i))$,
- (vi) $St_r(A) = St_l(A) = St_\odot(A) = \{1\}$,
- (vii) $St_r(\{1\}) = St_l(\{1\}) = A$,
- (viii) $\{x\} \subseteq St_\odot(\{x\})$ if and only if $x^2 = x$, for all $x \in A$,
- (ix) if $F \in \mathcal{F}(A)$, then $St_\odot(F) \subseteq F$,
- (x) if $h : A \rightarrow A$ is a homomorphism and $x \in A$, then

$$h(St_r(\{x\})) \subseteq St_r(\{h(x)\}), h(St_l(\{x\})) \subseteq St_l(\{h(x)\}), h(St_\odot(\{x\})) \subseteq St_\odot(\{h(x)\}).$$

Proof. The proofs of (i), (ii), (iii), (vii) and (viii) are clear by Proposition 2.2 and some modifications.

(iv) Since by (iii), $X \subseteq (St_l(X))_r$ and by (ii), $((St_l(X))_r)_l \subseteq St_l(X)$. Moreover, by (iii), $St_l(X) \subseteq ((St_l(X))_r)_l$. So, $St_l(X) = ((St_l(X))_r)_l$. By the similar way, we have, $St_r(X) = ((St_r(X))_l)_r$.

(v) For all $i \in I$, since $X_i \subseteq \bigcup_{i \in I} X_i$, by (ii), $St_l(\bigcup_{i \in I} X_i) \subseteq St_l(X_i)$, then $St_l(\bigcup_{i \in I} X_i) \subseteq \bigcap_{i \in I} (St_l(X_i))$. Now, suppose that $a \in \bigcap_{i \in I} (St_l(X_i))$. Then $a \in St_l(X_i)$, for all $i \in I$. Thus, $x_i \rightarrow a = a$, for all $x_i \in X_i$, and so $x \rightarrow a = a$, for all $x \in \bigcup_{i \in I} X_i$. Then $a \in St_l(\bigcup_{i \in I} X_i)$. Therefore, $St_l(\bigcup_{i \in I} X_i) = \bigcap_{i \in I} (St_l(X_i))$. By the similar way, we can prove the other statements.

(vi) Let $a \in St_r(A)$. Then $a \rightarrow x = x$, for all $x \in A$. By considering, $x = a$, and by (HP2),

$1 = a \rightarrow a = a$. Hence, $St_r(A) = \{1\}$. By the similar way, the other statements are clear.

(ix) Let $F \in \mathcal{F}(A)$ and $a \in St_{\odot}(F)$. Then, for all $x \in F$, $a \odot x = x$, and so $a \odot x \in F$. Since by Proposition 2.2(iii), $a \odot x \leq a$, and $F \in \mathcal{F}(A)$, we have $a \in F$. Hence, $St_{\odot}(F) \subseteq F$.

(x) Let $x \in A$, $h : A \rightarrow A$ be a homomorphism and $y \in h(St_r(\{x\}))$. Then there exists $a \in St_r(\{x\})$ such that $y = h(a)$. Since $a \in St_r(\{x\})$, $a \rightarrow x = x$. Moreover, since h is a homomorphism, we have $h(a) \rightarrow h(x) = h(a \rightarrow x) = h(x)$, and so $y \rightarrow h(x) = h(x)$. Hence, $y \in St_r(\{h(x)\})$.

By the similar way, we can prove the other inclusion. \square

Example 3.4. In Example 3.2(ii), let $F = \{b, 1\}$. It is clear that $F \in \mathcal{F}(A)$. By routine calculations, $St_{\odot}(F) = \{1\}$. Thus, $F \not\subseteq St_{\odot}(F)$. So, if $F \in \mathcal{F}(A)$ and $F \neq \{1\}$, then $St_{\odot}(F) \neq F$.

Proposition 3.5. Let A be a \sqcup -hoop, $X \subseteq A$ and $a \in A$ such that $a \sqcup x = 1$, for all $x \in X$. Then $a \in St_r(X) \cap St_l(X)$.

Proof. Let $a \in A$ such that $a \sqcup x = 1$, for all $x \in X$. Then by Proposition 2.4, $1 = a \sqcup x = ((x \rightarrow a) \rightarrow a) \wedge ((a \rightarrow x) \rightarrow x)$, and so $(x \rightarrow a) \rightarrow a = 1$ and $(a \rightarrow x) \rightarrow x = 1$. Suppose that $(x \rightarrow a) \rightarrow a = 1$. Then by Proposition 2.2(iv), $a \leq x \rightarrow a$, and so $a \rightarrow (x \rightarrow a) = 1$. Hence, $x \rightarrow a = a$, for all $x \in X$. Thus, $a \in St_l(X)$. By the similar way, we can see that $a \rightarrow x = x$, for all $x \in X$, and so $a \in St_r(X)$. Therefore, $a \in St_r(X) \cap St_l(X)$. \square

Theorem 3.6. Let $X \subseteq A$. Then $St_r(X)$ and $St_{\odot}(X)$ are filters of A .

Proof. Let $X \subseteq A$. Then by Proposition 3.3(i), $1 \in St_r(X)$, and so $St_r(X) \neq \emptyset$. Now, let $a \leq b$ and $a \in St_r(X)$, for some $a, b \in A$. Then by Proposition 2.2(x), $b \rightarrow x \leq a \rightarrow x$, for all $x \in X$. Since $a \in St_r(X)$, we have $a \rightarrow x = x$, and so $b \rightarrow x \leq x$, for all $x \in X$. Moreover, since by Proposition 2.2(iv), $x \leq b \rightarrow x$ and so $b \rightarrow x = x$, for all $x \in X$. Hence, $b \in St_r(X)$. Suppose that $a, b \in St_r(X)$, for some $a, b \in A$. Since $a \rightarrow x = b \rightarrow x = x$, for all $x \in X$, by (HP3) we have, $(a \odot b) \rightarrow x = a \rightarrow (b \rightarrow x) = a \rightarrow x = x$, for all $x \in X$. So, $a \odot b \in St_r(X)$. Therefore, $St_r(X) \in \mathcal{F}(A)$. Now, we prove that $St_{\odot}(X) \in \mathcal{F}(A)$, too. Let $a, b \in St_{\odot}(X)$, for some $a, b \in A$. Then $a \odot x = b \odot x = x$, for all $x \in X$. Hence, by (HP1), $(a \odot b) \odot x = a \odot (b \odot x) = a \odot x = x$, for all $x \in X$. So, $a \odot b \in St_{\odot}(X)$. Now, suppose that $a \leq b$ and $a \in St_{\odot}(X)$. Then $a \odot x = x$, for all $x \in X$. Hence, by Proposition 2.2(iii) and (viii), $x = a \odot x \leq b \odot x \leq x$, and so $b \odot x = x$, for all $x \in X$, which implies that $b \in St_{\odot}(X)$. Therefore, $St_{\odot}(X) \in \mathcal{F}(A)$. \square

Example 3.7. In Example 3.2(ii), let $X = \{b\}$. Since $St_l(X) = \{0, a, 1\}$, it is clear that $St_l(X)$ is not a filter.

Definition 3.8. [12] Let $F \subseteq A$ such that $1 \in F$. Then

- F is called an *implicative filter* of A if $x \rightarrow (y \rightarrow z) \in F$ and $x \rightarrow y \in F$, then $x \rightarrow z \in F$, for all $x, y, z \in A$.
- F is called a *positive implicative filter* of A if $((x \rightarrow y) \rightarrow x) \rightarrow x \in F$, for all $x, y \in A$.
- F is called a *fantastic filter* of A if $z \rightarrow (y \rightarrow x) \in F$ and $z \in F$, then $((x \rightarrow y) \rightarrow y) \rightarrow x \in F$, for all $x, y, z \in A$.

Theorem 3.9. *Let $X \subseteq A$. Then the following statements hold.*

- (i) *If $a^2 = a$, for all $a \in A$, then $St_r(X)$ and $St_{\odot}(X)$ are implicative filters.*
- (ii) *If A is a bounded hoop with (DNP), then $St_r(X)$ and $St_{\odot}(X)$ are fantastic filters.*
- (iii) *If A is a bounded hoop with (DNP) and $a^2 = a$, for all $a \in A$, then $St_r(X)$ and $St_{\odot}(X)$ are positive implicative filters.*

Proof. (i) Let $X \subseteq A$ such that $a^2 = a$, for all $a \in A$. Then by Theorem 3.6, $St_r(X) \in \mathcal{F}(A)$, thus, $1 \in St_r(X)$. Now, suppose that $a \rightarrow (b \rightarrow c)$ and $a \rightarrow b \in St_r(X)$, for some $a, b, c \in A$. Then, for all $x \in X$, $(a \rightarrow (b \rightarrow c)) \rightarrow x = x$ and $(a \rightarrow b) \rightarrow x = x$. Hence, $x = (a \rightarrow (b \rightarrow c)) \rightarrow x = (a \rightarrow (b \rightarrow c)) \rightarrow ((a \rightarrow b) \rightarrow x)$. By (HP1) and (HP3), we have

$$x = (a \rightarrow (b \rightarrow c)) \rightarrow ((a \rightarrow b) \rightarrow x) = (b \rightarrow (a \rightarrow c)) \rightarrow ((a \rightarrow b) \rightarrow x) = [(a \rightarrow b) \odot (b \rightarrow (a \rightarrow c))] \rightarrow x.$$

Then by Proposition 2.2(xii) and (x),

$$x = [(a \rightarrow b) \odot (b \rightarrow (a \rightarrow c))] \rightarrow x \geq (a \rightarrow (a \rightarrow c)) \rightarrow x = (a^2 \rightarrow c) \rightarrow x.$$

Moreover, since $a^2 = a$, for all $a \in A$, we have $x \geq (a \rightarrow c) \rightarrow x$. Also, by Proposition 2.2(iv), $x \leq (a \rightarrow c) \rightarrow x$. Hence, $(a \rightarrow c) \rightarrow x = x$, for all $x \in X$. Then $a \rightarrow c \in St_r(X)$. Therefore, $St_r(X)$ is an implicative filter. Now, we prove that $St_{\odot}(X)$ is an implicative filter. By Theorem 3.6, $St_{\odot}(X) \in \mathcal{F}(A)$, and so $1 \in St_{\odot}(X)$. Let $a \rightarrow (b \rightarrow c)$ and $a \rightarrow b \in St_{\odot}(X)$, for some $a, b, c \in A$. Then, for all $x \in X$, $(a \rightarrow (b \rightarrow c)) \odot x = x$ and $(a \rightarrow b) \odot x = x$. Thus, by (HP1), $((a \rightarrow b) \odot (a \rightarrow (b \rightarrow c))) \odot x = x$. Hence, by (HP3) and Proposition 2.2(xii), (viii) and (iii),

$$x = ((a \rightarrow b) \odot (b \rightarrow (a \rightarrow c))) \odot x \leq (a \rightarrow (a \rightarrow c)) \odot x = (a^2 \rightarrow c) \odot x \leq x.$$

Then $(a^2 \rightarrow c) \odot x = x$. Since $a^2 = a$, for all $a \in A$, we have $(a \rightarrow c) \odot x = x$, for all $x \in X$, and so $a \rightarrow c \in St_{\odot}(X)$. Therefore, $St_{\odot}(X)$ is an implicative filter.

- (ii) Let A be a bounded hoop with (DNP) and $X \subseteq A$. Then by Theorem 3.6, $St_r(X) \in \mathcal{F}(A)$, and so $1 \in St_r(X)$. Now, suppose that $a \rightarrow (b \rightarrow c)$ and $a \in St_r(X)$, for some $a, b, c \in A$. Since $St_r(X) \in \mathcal{F}(A)$, by Proposition 2.9, $b \rightarrow c \in St_r(X)$. Hence, $(b \rightarrow c) \rightarrow x = x$, for all $x \in X$. Now, we prove that $((c \rightarrow b) \rightarrow b) \rightarrow c \in St_r(X)$. Since A has (DNP), by Proposition 2.3(iv), for all $x \in X$, $[((c \rightarrow b) \rightarrow b) \rightarrow c] \rightarrow x = [((b \rightarrow c) \rightarrow c) \rightarrow c] \rightarrow x$. Thus, by Proposition 2.2(xi), $[((c \rightarrow b) \rightarrow b) \rightarrow c] \rightarrow x = (b \rightarrow c) \rightarrow x$. Since $b \rightarrow c \in St_r(X)$, we have $(b \rightarrow c) \rightarrow x = x$, for all $x \in X$, and so $[((c \rightarrow b) \rightarrow b) \rightarrow c] \rightarrow x = x$. Hence, $((c \rightarrow b) \rightarrow b) \rightarrow c \in St_r(X)$.

Therefore, $St_r(X)$ is a fantastic filter. Now, we prove that $St_{\odot}(X)$ is a fantastic filter. By Theorem 3.6, $St_{\odot}(X) \in \mathcal{F}(A)$, and so $1 \in St_{\odot}(X)$. Suppose that $a \rightarrow (b \rightarrow c)$ and $a \in St_{\odot}(X)$, for some $a, b, c \in A$. Since $St_{\odot}(X) \in \mathcal{F}(A)$, by Proposition 2.9, $b \rightarrow c \in St_{\odot}(X)$. Hence, $(b \rightarrow c) \odot x = x$, for all $x \in X$. Now, we prove that $((c \rightarrow b) \rightarrow b) \rightarrow c \in St_{\odot}(X)$. Since A has (DNP), by Proposition 2.3(iv), for all $x \in X$, $[((c \rightarrow b) \rightarrow b) \rightarrow c] \odot x = [((b \rightarrow c) \rightarrow c) \rightarrow c] \odot x$. Thus, by Proposition 2.2(xi), $[((c \rightarrow b) \rightarrow b) \rightarrow c] \odot x = (b \rightarrow c) \odot x$. Since $b \rightarrow c \in St_{\odot}(X)$, we have $(b \rightarrow c) \odot x = x$, for all $x \in X$. So, $[((c \rightarrow b) \rightarrow b) \rightarrow c] \odot x = x$. Hence, $St_{\odot}(X)$ is a fantastic filter.

(iii) By Theorem 3.6, $St_r(X) \in \mathcal{F}(A)$. Then $1 \in St_r(X)$. Now, we prove that $((a \rightarrow b) \rightarrow a) \rightarrow a \in St_r(X)$. Since A has (DNP), by Proposition 2.3(iv) and (HP3), $((a \rightarrow b) \rightarrow a) \rightarrow a = (a \rightarrow (a \rightarrow b)) \rightarrow (a \rightarrow b) = (a^2 \rightarrow b) \rightarrow (a \rightarrow b)$. Moreover, since $a^2 = a$, for all $a \in A$, we have $[((a \rightarrow b) \rightarrow a) \rightarrow a] = (a \rightarrow b) \rightarrow (a \rightarrow b) = 1$. Thus, $[((a \rightarrow b) \rightarrow a) \rightarrow a] \rightarrow x = 1 \rightarrow x = x$. Hence, $St_r(X)$ is a positive implicative filter of A . By the similar way, we can prove that $St_{\odot}(X)$ is a positive implicative filter of A . \square

Proposition 3.10. *Let A be bounded with (DNP) and $X \subseteq A$. Then $St_r(X) = St_l(X)$.*

Proof. Let A be bounded with (DNP), $X \subseteq A$ and $a \in St_r(X)$. Then $a \rightarrow x = x$, for all $x \in X$. Hence, $(a \rightarrow x) \rightarrow x = 1$. Since A has (DNP), by Proposition 2.3(iv), $(a \rightarrow x) \rightarrow x = (x \rightarrow a) \rightarrow a$, and so $(x \rightarrow a) \rightarrow a = 1$. Also, by Proposition 2.2(iv), $a \rightarrow (x \rightarrow a) = 1$, then $x \rightarrow a = a$. Thus, $a \in St_l(X)$. Hence, $St_r(X) \subseteq St_l(X)$. By the similar way, we have $St_l(X) \subseteq St_r(X)$. Therefore, $St_r(X) = St_l(X)$. \square

Definition 3.11. [4] Let A be a \sqcup -hoop and F be a proper filter of A . Then F is called a *prime filter* of A , if $x \sqcup y \in F$, for some $x, y \in A$, then $x \in F$ or $y \in F$.

Theorem 3.12. *Let A be a \sqcup -hoop and $a \in A$. If $St_r(\{a\})(St_{\odot}(\{a\}))$ is a proper subset of A , then it is a prime filter.*

Proof. Let $a \in A$. Then by Theorem 3.6, $St_r(\{a\}) \in \mathcal{F}(A)$. Thus, $St_r(\{a\})$ is a proper filter. Now, let, for $x, y \in A$, $x \sqcup y \in St_r(\{a\})$, but $x, y \notin St_r(\{a\})$. Then $x \rightarrow a \neq a$ and $y \rightarrow a \neq a$, and so by Proposition 2.2(iv), $a < x \rightarrow a$ and $a < y \rightarrow a$. Since $x \sqcup y \in St_r(\{a\})$, we have $(x \sqcup y) \rightarrow a = a$. Hence, by Proposition 2.7(i), $a < (x \rightarrow a) \wedge (y \rightarrow a) = (x \sqcup y) \rightarrow a = a$, which is a contradiction. Therefore, $St_r(\{a\})$ is a prime filter. Moreover, by considering $a \in A$ and Theorem 3.6, $St_{\odot}(\{a\}) \in \mathcal{F}(A)$, and so $St_r(\{a\})$ is a proper filter. Now, let $x, y \in A$, $x \sqcup y \in St_{\odot}(\{a\})$ but $x, y \notin St_{\odot}(\{a\})$. Then $x \odot a \neq a$ and $y \odot a \neq a$, and so by Proposition 2.2(iii), $x \odot a < a$ and $y \odot a < a$. Since $x \sqcup y \in St_{\odot}(\{a\})$, we have $(x \sqcup y) \odot a = a$. Hence, by Proposition 2.7(ii), $a = (x \sqcup y) \odot a = (x \odot a) \sqcup (y \odot a) < a$, which is a contradiction. Therefore, $St_{\odot}(\{a\})$ is a prime filter. \square

Corollary 3.13. *Let A be a \sqcup -hoop and $X \subseteq A$. If $St_r(X)(St_\odot(X))$ is a proper subset of A , then it is a prime filter.*

Proof. The proof is similar to the proof of Proposition 3.12, by some modification. \square

Theorem 3.14. *Let A be a \sqcup -hoop, $X \subseteq A$ and $F_1, F_2 \in \mathcal{F}(A)$. If $St_r(X)$ is a proper subset of A such that $St_r(X) = F_1 \cap F_2$, then $St_r(X) = F_1$ or $St_r(X) = F_2$.*

Proof. Let $St_r(X)$ be a proper subset of A such that $St_r(X) = F_1 \cap F_2$, for some $F_1, F_2 \in \mathcal{F}(A)$. Since $St_r(X) = F_1 \cap F_2$, we have $St_r(X) \subseteq F_1, F_2$. Now, suppose that $St_r(X) = F_1 \cap F_2$ such that $F_1 \not\subseteq St_r(X)$ and $F_2 \not\subseteq St_r(X)$. Then there exist $a \in F_1$ and $b \in F_2$ such that $a, b \notin St_r(X)$. Since $F_1, F_2 \in \mathcal{F}(A)$ and $a, b \leq a \sqcup b$, $a \sqcup b \in F_1 \cap F_2 = St_r(X)$. Moreover, since $St_r(X)$ is a proper subset of A , by Corollary 3.13, $St_r(X)$ is a prime filter. Hence, $a \in St_r(X)$ or $b \in St_r(X)$, which is a contradiction. Thus, $F_1 \subseteq St_r(X)$ or $F_2 \subseteq St_r(X)$. Therefore, $St_r(X) = F_1$ or $St_r(X) = F_2$. \square

By the similar way, we can prove that the Theorem 3.14 holds for $St_\odot(X)$, for $X \subseteq A$.

Proposition 3.15. *Let $X \subseteq A$. Then*

- (i) $A/St_r(X) = \{((a \rightarrow b) \odot (b \rightarrow a)) \rightarrow x = x, \text{ for all } x \in X\}$.
- (ii) $A/St_\odot(X) = \{((a \rightarrow b) \odot (b \rightarrow a)) \odot x = x, \text{ for all } x \in X\}$.

Proof. (i) Since by Theorem 3.6, $St_r(X) \in \mathcal{F}(A)$, we have $A/St_r(X)$ is well-defined. Hence,

$$A/St_r(X) = \{\bar{b} \mid b \in A\} = \{a \in A \mid a \rightarrow b \text{ and } b \rightarrow a \in St_r(X)\}.$$

Thus, for all $x \in X$, $(a \rightarrow b) \rightarrow x = x$ and $(b \rightarrow a) \rightarrow x = x$. By (HP3), for all $x \in X$,

$$((a \rightarrow b) \odot (b \rightarrow a)) \rightarrow x = (a \rightarrow b) \rightarrow ((b \rightarrow a) \rightarrow x) = (a \rightarrow b) \rightarrow x = x.$$

Therefore, $A/St_r(X) = \{((a \rightarrow b) \odot (b \rightarrow a)) \rightarrow x = x, \text{ for all } x \in X\}$.

(ii) The proof is similar to the proof of (i). \square

Proposition 3.16. *Let $X \subseteq A$ and $F \in \mathcal{F}(A)$. Then the following statements hold.*

- (i) *If $F \subseteq St_r(X)$ and $\bar{a} \in St_r(X)/F$, then $(a \rightarrow x) \rightarrow x \in F$, for all $x \in X$.*
- (ii) *If $F \subseteq St_\odot(X)$ and $\bar{a} \in St_\odot(X)/F$, then $x \rightarrow (a \odot x) \in F$, for all $x \in X$.*

Proof. (i) Let $F \subseteq St_r(X)$ and $\bar{a} \in St_r(X)/F$. Then there exists $b \in St_r(X)$ such that $\bar{a} = \bar{b}$. By definition of quotient, $a \rightarrow b, b \rightarrow a \in F$. Since $F \subseteq St_r(X)$, we have $a \rightarrow b, b \rightarrow a \in St_r(X)$. By Theorem 3.6, $St_r(X) \in \mathcal{F}(A)$ and since $b \rightarrow a, b \in St_r(X)$, by Proposition 2.9, $a \in St_r(X)$. Hence, for all $x \in X$, $a \rightarrow x = x$, and so $(a \rightarrow x) \rightarrow x = 1 \in F$. Therefore, $(a \rightarrow x) \rightarrow x \in F$.

(ii) Let $F \subseteq St_\odot(X)$ and $\bar{a} \in St_\odot(X)/F$. Then there exists $b \in St_\odot(X)$ such that $\bar{a} = \bar{b}$. By definition

of quotient, $a \rightarrow b$, $b \rightarrow a \in F$. Since $F \subseteq St_{\odot}(X)$, we have $a \rightarrow b$, $b \rightarrow a \in St_{\odot}(X)$. By Theorem 3.6, $St_{\odot}(X) \in \mathcal{F}(A)$ and since $b \rightarrow a$, $b \in St_{\odot}(X)$, by Proposition 2.9, $a \in St_{\odot}(X)$. Then, for all $x \in X$, $a \odot x = x$, and so $x \leq a \odot x$. Hence, $x \rightarrow (a \odot x) = 1 \in F$. Therefore, $x \rightarrow (a \odot x) \in F$. \square

Proposition 3.17. *Let $X \subseteq A$ and $F \in \mathcal{F}(A)$ such that $F \subseteq St_r(X)$ ($F \subseteq St_{\odot}(X)$). Then $St_r(X)/F$ ($St_{\odot}(X)/F$) is a filter of A/F .*

Proof. Let $X \subseteq A$ and $F \in \mathcal{F}(A)$ such that $F \subseteq St_r(X)$. Since $F \in \mathcal{F}(A)$, we have $St_r(X)/F$ and A/F are well-defined. Now, let $\bar{a}, \bar{b} \in St_r(X)/F$, for $a, b \in St_r(X)$. Then by Theorem 3.6, $St_r(X) \in \mathcal{F}(A)$, and so $a \odot b \in St_r(X)$. Hence, $\overline{a \odot b} = \bar{a} \odot \bar{b} = a \odot b/F \in St_r(X)/F$. Now, suppose that $\bar{a} \leq \bar{b}$ and $\bar{a} \in St_r(X)/F$. Since $\bar{a} \leq \bar{b}$, we have $a \rightarrow b \in F$. Moreover, since $F \subseteq St_r(X)$, $a \rightarrow b \in St_r(X)$. Also, $a \in St_r(X)$ and by Theorem 3.6, $St_r(X) \in \mathcal{F}(A)$, then by Proposition 2.9, $b \in St_r(X)$. Hence, $\bar{b} \in St_r(X)/F$. Therefore, $St_r(X)/F$ is a filter of A/F . \square

Proposition 3.18. (i) $St_r(A/St_r(A)) = \bar{1}$.

(ii) $St_{\odot}(A/St_{\odot}(A)) = \bar{1}$.

Proof. (i) Suppose that $\bar{a} \in St_r(A/St_r(A))$. Then, for all $\bar{x} \in A/St_r(A)$, $\bar{a} \rightarrow \bar{x} = \bar{x}$. By definition of quotient, $(a \rightarrow x) \rightarrow x \in St_r(A)$. So, for all $x \in A$, $((a \rightarrow x) \rightarrow x) \rightarrow x = x$. Since by Proposition 2.2(xi), $((a \rightarrow x) \rightarrow x) \rightarrow x = a \rightarrow x$, we have $a \rightarrow x = x$, and so $a \in St_r(A)$. Moreover, suppose that $x = 1$. Since $a \rightarrow 1 = 1 \in St_r(A)$ and $1 \rightarrow a = a \in St_r(A)$, $a \sim_{St_r(A)} 1$. This means that $\bar{a} = \bar{1}$. Therefore, $St_r(A/St_r(A)) = \bar{1}$.

(ii) Suppose that $\bar{a} \in St_{\odot}(A/St_{\odot}(A))$. Then, for all $\bar{x} \in A/St_{\odot}(A)$, $\bar{a} \odot \bar{x} = \bar{x}$. By definition of quotient, $x \rightarrow (a \odot x) \in St_{\odot}(A)$. So, for all $x \in A$, $(x \rightarrow (a \odot x)) \odot x = x$. By Proposition 2.2(vii) and (iii), $x = x \odot (x \rightarrow (a \odot x)) \leq a \odot x \leq x$. Then $a \odot x = x$, and so $a \in St_{\odot}(A)$. Hence, similar to the proof of (i), $\bar{a} = \bar{1}$. Therefore, $St_{\odot}(A/St_{\odot}(A)) = \bar{1}$. \square

4. STABILIZER TOPOLOGY

In this section, we use of the right and left stabilizers of a hoop and produce a basis for a topology on it. Then we show that the generated topology by this basis is Baire, connected, locally connected and separable and investigate the other properties of this topology.

Theorem 4.1. *Let $X \subseteq A$. Define $\alpha : \mathcal{P}(A) \rightarrow \mathcal{P}(A)$ such that $\alpha(X) = (St_l(X))_r$, for all $X \in \mathcal{P}(A)$.*

Then

(i) α is a closure map.

(ii) $X \subseteq \alpha(Y)$ if and only if $\alpha(X) \subseteq \alpha(Y)$, for all $Y \subseteq A$.

(iii) $\beta_{\alpha} = \{X \in \mathcal{P}(A) \mid \alpha(X) = X\}$ is a basis for a topology on A .

Proof. (i) By Proposition 3.3(iii), (iv) and (ii), the proof is clear.

(ii) It is straightforward.

(iii) Let $\beta_\alpha = \{X \in \mathcal{P}(A) \mid \alpha(X) = X\}$. It is clear that $\emptyset \in \beta_\alpha$. Also, by Proposition 3.3(vi) and (vii), $\alpha(A) = (St_l(A))_r = St_r(\{1\}) = A$. Thus, $\alpha(A) = A$, and so $A \in \beta_\alpha$. Now, suppose that $X, Y \in \beta_\alpha$. Then $\alpha(X) = X$ and $\alpha(Y) = Y$. We prove that $X \cap Y \in \beta_\alpha$. Since $X \cap Y \subseteq X, Y$, by (i), $\alpha(X \cap Y) \subseteq \alpha(X)$ and $\alpha(Y)$. Thus, $\alpha(X \cap Y) \subseteq \alpha(X) \cap \alpha(Y)$. Also, since $X, Y \in \beta_\alpha$, we have $\alpha(X \cap Y) \subseteq X \cap Y$. Moreover, by Proposition 3.3(iii), $X \cap Y \subseteq \alpha(X \cap Y)$. Then $\alpha(X \cap Y) = X \cap Y$, and so $X \cap Y \in \beta_\alpha$. Therefore, β_α is a basis. \square

Definition 4.2. According to Theorem 4.1, the topological space, (A, \mathcal{T}_α) is called a *stabilizer topology*.

Note: Since $St_r(X) \in \mathcal{F}(A)$, for any $X \subseteq A$, every element of β_α is a filter of A .

Example 4.3. In Example 3.2(ii), $(A, \odot, \rightarrow, 1)$ is a bounded hoop. By Proposition 3.3(i) and (vi), we see that $\{1\} \in \alpha(X)$, for all non-empty subset X of A . So, if $1 \notin X \subseteq A$, then $X \notin \beta_\alpha$. By some manipulations, we get that $\beta_\alpha = \{\emptyset, A, \{1, b\}, \{1\}\}$. Thus, $\mathcal{T}_\alpha = \{\emptyset, A, \{1\}, \{1, b\}\}$. By routine calculation, we can see that in this example, $\{1, a\} \notin \beta_\alpha$, because $St_l(\{1, a\}) = \{1\}$ and $St_r(\{1\}) = A$, then $\alpha(\{1, a\}) = A$, and so $\alpha(\{1, a\}) \neq \{1, a\}$.

Theorem 4.4. *The stabilizer topology (A, \mathcal{T}_α) is*

- (i) *connected.*
- (ii) *locally connected.*
- (ii) *Hausdorff space if and only if $A = \{1\}$.*

Proof. It is straightforward. \square

Theorem 4.5. *Let (A, \mathcal{T}_α) be a stabilizer topology. If $\emptyset \neq X \subseteq A$ such that $1 \in X$, then $\overline{X} = A$.*

Proof. Let $\emptyset \neq X \subseteq A$ such that $1 \in X$. It is enough to prove that $A \subseteq \overline{X}$. Let $x \in A$. If $x = 1$, then $x \in \overline{X}$. Hence, $\overline{X} = A$. Now, suppose that $1 \neq x \in A$. Then there exists an open subset $U \in \beta_\alpha$ such that $x \in U$. Since $U \in \mathcal{F}(A)$ and $1 \in U$, we have $U \cap (X - \{x\}) \neq \emptyset$. Hence, $x \in \overline{X}$, and so $\overline{X} = A$. \square

Corollary 4.6. *(A, \mathcal{T}_α) is separable.*

Proof. Since $\{1\} \in \beta_\alpha$, by Theorem 4.5, $\overline{\{1\}} = A$. Hence, (A, \mathcal{T}_α) is separable. \square

Theorem 4.7. *(A, \mathcal{T}_α) is Baire space.*

Proof. Let $U \in \mathcal{T}_\alpha$. Since $U \in \mathcal{F}(A)$, we have $1 \in U$. Then by Theorem 4.5, $\overline{U} = A$. Thus, every open set of (A, \mathcal{T}_α) is dense. On the other side, for each collection of open set $U_n, \bigcap U_n \in \mathcal{F}(A)$, and so $1 \in \bigcap U_n$. Thus, by Theorem 4.5, $\bigcap U_n$ is dense. Therefore, (A, \mathcal{T}_α) is Baire space. \square

In the following example, we show that (A, \mathcal{T}_α) is not a T_0 -space or T_1 -space.

Example 4.8. In Example 3.2(i) and (ii), $\beta_\alpha = \{\emptyset, A, \{1\}\}$. Since $a \neq b$, for $a, b \in A$, there is not $U \in \beta_\alpha$ such that $a \in U$ and $b \notin U$. Therefore, (A, \mathcal{T}_α) is not a T_0 -space. It is clear that (A, \mathcal{T}_α) is not a T_1 -space.

Theorem 4.9. *Let A be bounded. If A has a cover of $U_i \in \beta_\alpha$, for $i \in I$, then there exists $i \in I$ such that $U_i = A$.*

Proof. Let A be bounded and $\{U_i\}_{i \in I}$ be a cover of A such that, for all $i \in I$, $U_i \in \beta_\alpha$ and $A \subseteq \bigcup_{i \in I} U_i$. Since, for all $i \in I$, $U_i \in \beta_\alpha$, we have $U_i \in \mathcal{F}(A)$. On the other side, A is bounded, then $0 \in A$, and so $0 \in \bigcup_{i \in I} U_i$. Thus, there exists $i \in I$ such that $0 \in U_i$. Since $U_i \in \mathcal{F}(A)$ and $0 \in U_i$, then $U_i = A$. Hence, there exists a finite family of $\{U_i\}_{i \in I}$ such that $A \subseteq \bigcup_{i=1}^n U_i$. \square

Theorem 4.10. *Let $F \in \mathcal{F}(A)$ and $\bar{\alpha} : \mathcal{P}(A/F) \rightarrow \mathcal{P}(A/F)$ is defined by $\bar{\alpha}(X/F) = (St_l(X/F))_r$, for all $X/F \in \mathcal{P}(A/F)$. Then*

- (i) $\bar{\alpha}$ is a closure map.
- (ii) $\beta_{\bar{\alpha}} = \{X/F \mid \bar{\alpha}(X/F) = X/F, F \subseteq X \subseteq A\}$ is a basis for a topology on A/F .

Proof. The proof is similar to the proof of Theorem 4.1. \square

Note: If $\mathcal{T}_{\bar{\alpha}}$ is a topology induced by $\beta_{\bar{\alpha}}$, then $(A/F, \mathcal{T}_{\bar{\alpha}})$ is called the *quotient stabilizer topology* or *QS-topology*, for short. It is clear that $\mathcal{T}_{\bar{\alpha}} = \{O \subseteq A/F \mid \pi^{-1}(O) \in \mathcal{T}_\alpha\}$, where $\pi : A \rightarrow A/F$ is the canonical epimorphism.

Theorem 4.11. *Let $F \in \mathcal{F}(A)$, $F \subseteq X \subseteq A$ and $\pi : A \rightarrow A/F$ be an open canonical epimorphism. Then X/F is an open set in the QS-topology, $\mathcal{T}_{\bar{\alpha}}$.*

Proof. Let $F \in \mathcal{F}(A)$ and $F \subseteq X \subseteq A$. Then $X/F \subseteq A/F$. Since π is an open epimorphism, there exists an open subset U of A such that $\pi(U) = X/F$. Thus, $U = \pi^{-1}(X/F)$. Since U is open, there is a $V \in \beta_\alpha$ such that $V \subseteq U = \pi^{-1}(X/F)$. Hence, $\pi(V) \subseteq X/F$, and so $V/F \subseteq X/F$. Therefore, X/F is an open subset in the QS-topology, $\mathcal{T}_{\bar{\alpha}}$. \square

Theorem 4.12. *Let S be a topological subalgebra of A and $F \in \mathcal{F}(A)$ such that $F \subseteq S$. Then the QS-topology on S/F is stronger than the topology induced on the subalgebra S/F of A/F by the QS-topology on A/F .*

Proof. Let $\pi_S : S \rightarrow S/F$ be the canonical map and U be a set in the subspace topology of S/F . Then $U = W \cap S/F$, where W is an open set in A/F . We obtain that $\pi_S^{-1}(U) = \pi_S^{-1}(W) \cap S$. Since W and S/F are open sets and π_S is a continuous epimorphism, we have $\pi_S^{-1}(W) \cap S$ is an open set. Hence, $\pi^{-1}(U)$ is open in S . Therefore, U is an open set in the quotient topology of S/F . \square

Theorem 4.13. *Let $F \in \mathcal{F}(A)$ and $(A/F, \mathcal{T}_{\bar{\alpha}})$ be the QS-topology. Then $(A/F, \mathcal{T}_{\bar{\alpha}})$ is connected.*

Proof. The proof is similar to the proof of Theorems 4.4(i). \square

Theorem 4.14. *Let $F \in \mathcal{F}(A)$, $(A/F, \mathcal{T}_{\bar{\alpha}})$ be the QS-topology and $\pi : A \rightarrow A/F$ be an open continuous epimorphism. Then $(A/F, \mathcal{T}_{\bar{\alpha}})$ is Hausdorff if and only if F is closed.*

Proof. (\Rightarrow) We prove that F^c is open. Let $x \in F^c$, for some $x \in A$. Since $x \notin F$, then $\bar{x} \neq \bar{1}$. Since A/F is Hausdorff, there exist two open neighborhoods U and V of \bar{x} and $\bar{1}$, respectively such that $U \cap V = \emptyset$. So, $\pi^{-1}(U)$ and $\pi^{-1}(V)$ are two open neighborhoods of x and 1 , respectively. Let $z \in F$. Then $\pi(z) = \bar{z} = \bar{1} \in V$. Hence, $z \in \pi^{-1}(V)$ and then $F \subseteq \pi^{-1}(V)$. We show that $\pi^{-1}(U) \subseteq F^c$. Let $z \in \pi^{-1}(U)$. Then $\bar{z} \in U$. Since $U \cap V = \emptyset$, we have $z \notin \pi^{-1}(V)$. Moreover, since $F \subseteq \pi^{-1}(V)$, we have $z \notin F$. So, $z \in F^c$. Hence, $\pi^{-1}(U) \subseteq F^c$. Since $\pi^{-1}(U)$ is an open set, we get that F^c is open. Therefore, F is closed.

(\Leftarrow) Suppose that F is closed and $\bar{x} \neq \bar{y}$ in A/F . Then $\bar{x} \rightarrow \bar{y} \neq \bar{1}$ or $\bar{y} \rightarrow \bar{x} \neq \bar{1}$. Let $\bar{x} \rightarrow \bar{y} \neq \bar{1}$. Then $x \rightarrow y \notin F$. So, $x \rightarrow y \in F^c$. Moreover, since F^c is an open, there exists an open subset U such that $x \rightarrow y \in U \subseteq F^c$ and $U \cap F = \emptyset$. Since π is an open continuous epimorphism, we have $\pi(U)$ is open neighborhood of $\bar{x} \neq \bar{y}$. If $\bar{1} \in \pi(U)$, then there exists $a \in U$ such that $\bar{a} = \bar{1}$, and so $a \rightarrow 1, 1 \rightarrow a \in F$. Hence, $a \in F$, which is a contradiction. Therefore, $\bar{1} \notin \pi(U)$. Now, since $\mathcal{T}_{\bar{\alpha}}$ is the QS-topology on A/F , there exist open neighborhoods \bar{P} and \bar{Q} of \bar{x} and \bar{y} such that

$$\bar{x} \rightarrow \bar{y} \in \bar{P} \rightarrow \bar{Q} \subseteq \overline{\bar{P} \rightarrow \bar{Q}} \subseteq \pi(P \rightarrow Q) \subseteq \pi(P) \rightarrow \pi(Q) \subseteq \pi(U).$$

Now, we prove that $\pi(P) \cap \pi(Q) = \emptyset$. Let $\bar{z} \in \pi(P) \cap \pi(Q)$. Then there are $a \in P$ and $b \in Q$ such that $\bar{a} = \bar{z} = \bar{b}$. Since $\bar{z} \rightarrow \bar{z} = \bar{a} \rightarrow \bar{b} \in \pi(P) \rightarrow \pi(Q) \subseteq \pi(U)$. Hence, $\bar{1} \in \pi(U)$, which is a contradiction. Therefore, $(A/F, \mathcal{T}_{\bar{\alpha}})$ is Hausdorff. \square

Acknowledgments. The authors wish to express their appreciation for several excellent suggestions for improvements in this paper made by the referee.

5. CONCLUSIONS AND FUTURE WORKS

In this paper, we introduce the concepts of right, left and product stabilizers on hoops. Then by using the right stabilizers produce a basis for a topology on hoops. We show that the generated topology by this basis is Baire, connected, locally connected and separable. Finally, we investigate the other properties of this topology.

REFERENCES

- [1] P. Aglianó, I. M. A. Ferreirim, F. Montagna, Basic hoops: an algebraic study of continuous t-norm, draft, (2000).
- [2] B. Bosbach, Komplementäre Halbgruppen. Axiomatik und Arithmetik, *Fundamenta Mathematicae*, Vol. 64 (1969), 257-287.
- [3] B. Bosbach, Komplementäre Halbgruppen. Kongruenzen and Quotienten, *Fundamenta Mathematicae*, Vol. 69 (1970), 1-14.
- [4] M. Botur, A. Dvurečenskij, T. Kowalski, On normal-valued basic pseudo-hoop, *Soft Comput*, Vol. 16, (2012), 635-644.
- [5] N. Bourbaki, Topologie Générale, *Springer Berlin Heidelberg*, (2007).
- [6] J. R. Büchi, T. M. Owens, Complemented monoids and hoops, unpublished manuscript, (1975).
- [7] G. Georgescu, L. Leustean, V. Preoteasa, Pseudo-hoops, *Journal of Multiple-Valued logic and Soft Computing*, Vol. 11. No 1-2, (2005), 153-184.
- [8] P. Hájek, Metamathematics of fuzzy logic, *Springer*, Vol. 4. (1998).
- [9] K. D. Joshi, Introduction to general topology, *New Age International Publisher, India*, (1983).
- [10] Y. B. Jun, H. S. Kim, Uniform structures in positive implication algebras, *Intern. Math. J.* Vol. 2, No 2, (2002), 215-219.
- [11] Y. B. Jun, E. H. Roh, On uniformities of BCK-algebras, *Commun. Korean Math. Soc.* Vol. 10, No 1, (1995), 11-14.
- [12] M. Kondo, Some types of filters in hoops, *Multiple-Valued Logic (ISMVL)*, (2011), 41st IEEE International Symposium on. IEEE, 50-53.
- [13] J. R. Munkres, Topology a first course, Prentice-Hall, (1975).
- [14] B. T. Sims, Fundamentals of Topology, *Macmillan Publishing Co., Inc., New York*, (1976).
- [15] D. S. Yoon, H. S. Kim, Uniform structures in BCI-algebras, *Commun. Korean Math. Soc.* Vol. 17, No 3, (2002), 403-408.

R. A. Borzooei

Department of Mathematics, Shahid Beheshti University, Tehran, Iran

Email: borzooei@sbu.ac.ir

M. Aaly Kologani

Department of Mathematics, Payamenour University, Tehran, Iran

Email: alykologani@phd.pnu.ac.ir