



## ON NEW CLASSES OF MULTICONE GRAPHS DETERMINED BY THEIR SPECTRUMS

ALI ZEYDI ABDIAN AND S. MORTEZA MIRAFZAL

Communicated by M.A. Iranmanesh

**ABSTRACT.** A multicone graph is defined to be join of a clique and a regular graph. A graph  $G$  is cospectral with graph  $H$  if their adjacency matrices have the same eigenvalues. A graph  $G$  is said to be determined by its spectrum or DS for short, if for any graph  $H$  with  $Spec(G) = Spec(H)$ , we conclude that  $G$  is isomorphic to  $H$ . In this paper, we present new classes of multicone graphs that are DS with respect to their spectrums. Also, we show that complement of these graphs are DS with respect to their adjacency spectrums. In addition, we show that graphs cospectral with these graphs are perfect. Finally, we find automorphism group of these graphs and one conjecture for further researches is proposed.

### 1. INTRODUCTION

In this paper, we are concerned only with finite undirected simple graph (loops and multiple edge are not allowed). All terminology and notation on graphs not defined here can be found

---

MSC(2010): Primary:05C50

Keywords: Adjacency spectrum, Laplacian spectrum, Multicone graph, DS graph, Automorphism group.

Received: 07 May 2015, Accepted: 12 Oct 2015.

\*Corresponding author

in [2, 5, 9]. Let  $\Gamma = (V, E)$  be a simple graph, where  $V$  is the set of vertices and  $E$  is the set of edges of  $\Gamma$ . An edge joining the vertices  $u$  and  $v$  is denoted by  $\{u, v\}$ . Permutation of a set  $\Omega$  is a bijection from  $\Omega$  to itself.  $\text{Sym}(\Omega)$  denotes the set of all permutations of  $\Omega$ .  $S_n$  denotes the symmetric group  $\text{sym}(\{1, 2, \dots, n\})$ . Automorphism of a simple graph  $\Gamma = (V, E)$  is a permutation of the vertices of  $\Gamma$  which preserves the relation of adjacency; that is, a bijection  $\pi : V \rightarrow V$  such that  $\{u, v\} \in E$  if and only if  $\{\pi(u), \pi(v)\} \in E$ . Under composition, the automorphisms form a group, called the automorphism group (or symmetry group) of  $\Gamma$ , and this is denoted by  $\text{Aut}(\Gamma)$ . In most situations it is very difficult to determine the automorphism group of a graph and this has been the subject of many research papers. Some of recent works appear in references [7, 8, 9]. The complement of a graph  $G$ , denoted by  $\overline{G}$ , is the graph on the vertex set of  $G$  such that two vertex of  $\overline{G}$ , are adjacent if and only if they are not adjacent in  $G$ . A graph and its complement have the same automorphisms. The automorphism group of the complete graph  $K_n$  and the empty graph  $\overline{K_n}$  is the symmetric group  $S_n$ . The union of two vertex disjoint graphs  $G_1$  and  $G_2$  denoted by  $G_1 \cup G_2$ , is the graph whose vertex (respectively, edge) set is the union of vertex (respectively, edge) sets of  $G_1$  and  $G_2$ . The join of two vertex disjoint graphs  $G_1$  and  $G_2$  is the graph obtained from  $G_1 \cup G_2$  by joining each vertex in  $G_1$  with every vertex in  $G_2$ . It is denoted by  $G_1 \nabla G_2$ . The group  $G$  is called a semidirect product of  $N$  by  $Q$ , denoted by  $G = N \rtimes Q$ , if  $G$  contains subgroups  $N$  and  $Q$  such that (i)  $N \trianglelefteq G$  ( $N$  is a subgroup normal of  $G$ ); (ii)  $NQ = G$ ; (iii)  $N \cap Q = 1$ . If  $\Gamma$  and  $\Delta$  are nonempty sets then we write  $\text{Fun}(\Gamma, \Delta)$  to denote the set of all functions  $\Gamma$  into  $\Delta$ . In the case when  $K$  is a group, we can turn  $\text{Fun}(\Gamma, \Delta)$  into a group by defining a product "pointwise":  $fg(\gamma) = f(\gamma)g(\gamma)$ , for all  $f, g \in \text{Fun}(\Gamma, \Delta)$  and  $\gamma \in \Gamma$ . and the product on the right is in  $K$ . In the case that  $\Gamma$  is finite of size  $m$ , say  $\Gamma = \{\gamma_1, \gamma_2, \dots, \gamma_m\}$  then the group  $\text{Fun}(\Gamma, \Delta)$  is isomorphic to  $K^m$  ( a direct product of  $m$  copies of  $K$  ) via the isomorphism. Let  $K$  and  $H$  be groups and suppose  $H$  acts on the nonempty set  $\Gamma$ . Then the wreath product of  $K$  by  $H$  with respect to this action is defined to be the semidirect  $\text{Fun}(\Gamma, K) \rtimes H$  where  $H$  acts on the group  $\text{Fun}(\Gamma, \Delta)$  via  $f^x(\gamma) = f(\gamma^{x^{-1}})$  for all  $f \in \text{Fun}(\Gamma, \Delta)$ ,  $\gamma \in K$  and  $x \in H$ . We denote this group by  $K \wr_{\Gamma} H$ .

Let  $A(G)$  denotes the  $(0, 1)$ -adjacency matrix of graph  $G$ . The characteristic polynomial of  $G$  is  $\det(\lambda I - A(G))$ , and is denoted by  $P_G(\lambda)$ . The roots of  $P_G(\lambda)$  are called the adjacency eigenvalues of  $G$  and since  $A(G)$  is real and symmetric, the eigenvalues are real numbers. If  $G$  has  $n$  vertices, then it has  $n$  eigenvalues in descending order as  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ . Let  $\lambda_1, \lambda_2, \dots, \lambda_s$  be the distinct eigenvalues of  $G$  with multiplicity  $m_1, m_2, \dots, m_s$ , respectively. The multi-set  $\text{Spec}(G) = \{\lambda_1^{m_1}, \lambda_2^{m_2}, \dots, \lambda_s^{m_s}\}$  of eigenvalues of  $A(G)$  is called the adjacency spectrum of  $G$ . For two graphs  $G$  and  $H$ , if  $\text{Spec}(G) = \text{Spec}(H)$ , we say  $G$  and  $H$  are cospectral with respect to adjacency matrix. A graph  $H$  is said to be determined by its spectrum or DS for short, if for a graph  $H$  with  $\text{Spec}(G) = \text{Spec}(H)$ , one has  $G$  isomorphic to  $H$ . Let  $G$

be a graph with adjacency matrix  $A$  and  $D$  be the diagonal matrix of vertex degrees for  $G$ . The matrix  $SL(G) = D + A$  and  $L(G) = D - A$  are known as signless Laplacian matrix and Laplacian matrix for  $G$ , respectively.

## 2. Preliminaries

**Lemma 2.1.** [1, 15] *Let  $G$  be a graph. For the adjacency matrix and Laplacian matrix, the following can be obtained from the spectrum:*

- (i) *The number of vertices,*
- (ii) *The number of edges.*

*For the adjacency matrix, the following follows from the spectrum*

- (iii) *The number of closed walks of any length.*
- (iv) *Being regular or not and the degree of regularity.*
- (v) *Being bipartite or not.*

*For the Laplacian matrix, the following follows from the spectrum*

- (vi) *The number of spanning tree.*
- (vii) *The number of component.*
- (viii) *The sum of squares of degrees of vertices.*

**Theorem 2.2.** [4] *If  $G_1$  is  $r_1$ -regular with  $n_1$  vertices, and  $G_2$  is  $r_2$ -regular with  $n_2$  vertices, then the characteristic polynomial of the join  $G_1 \nabla G_2$  is given by:*

$$P_{G_1 \nabla G_2}(x) = \frac{P_{G_1} P_{G_2}(x)}{(x-r_1)(x-r_2)} ((x-r_1)(x-r_2) - n_1 n_2).$$

**Proposition 2.3.** [17] *Let  $G$  be a disconnected graph that is determined by the Laplacian spectrum. Then the cone over  $G$ , the graph  $H$ ; that is, obtained from  $G$  by adding one vertex that is adjacent to all vertices of  $G$ , is also determined by its Laplacian spectrum.*

**Theorem 2.4.** [19] *A graph has exactly one positive eigenvalue if and only if its non-isolated vertices form a complete multipartite graph.*

**Theorem 2.5.** [1] *Let  $G$  be a simple graph with  $n$  vertices and  $m$  edges. Let  $\delta = \delta(G)$  be the minimum degree of vertices of  $G$  and  $\varrho(G)$  be the spectral radius of the adjacency matrix of  $G$ . Then*

$$\varrho(G) \leq \frac{\delta-1}{2} + \sqrt{2m - n\delta + \frac{(\delta+1)^2}{4}}$$

*Equality holds if and only if  $G$  is either a regular graph or a bidegreed graph in which each vertex is of degree either  $\delta$  or  $n - 1$ .*

**Lemma 2.6.** [3] *Let  $\Gamma$  be a connected non-regular graph with three distinct eigenvalues  $\theta_0 > \theta_1 > \theta_2$ . Then the following hold:*

- (i)  $\Gamma$  has diameter two.
- (ii) If  $\theta_0$  is not an integer, then  $\Gamma$  is complete bipartite.
- (iii)  $\theta_1 \geq 0$  with equality if and only if  $\Gamma$  is complete bipartite.
- (iv)  $\theta_2 \leq -\sqrt{2}$  with equality if and only if  $\Gamma$  is the path of length 2.

**Theorem 2.7.** [13, 14] Let  $G$  and  $G$  be two graphs with Laplacian spectrum  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$  and  $\mu_1 \geq \mu_2 \geq \dots \geq \mu_m$ , respectively. Then Laplacian spectra of  $\overline{G}$  and  $G \nabla H$  are  $n - \lambda_1, n - \lambda_2, \dots, n - \lambda_{n-1}, 0$  and  $n + m, m + \lambda_1, \dots, m + \lambda_{n-1}, n + \mu_1, \dots, m + \mu_{m-1}, 0$ , respectively.

**Lemma 2.8.** [14] Let  $G$  be a graph on  $n$  vertices. Then  $n$  is Laplacian eigenvalue of  $G$  if and only if  $G$  is the join of two graphs.

**Proposition 2.9.** [11] For a graph  $G$ , the following statements are equivalent:

- (i)  $G$  is  $d$ -regular.
- (ii)  $\rho(G) = d_G$ , the average vertex degree.
- (iii)  $G$  has  $v = (1, 1, \dots, 1)^t$  as an eigenvector for  $\rho(G)$ .

**Proposition 2.10.** [4] Let  $G - j$  be the graph obtained from  $G$  by deleting the vertex  $j$  and all edges containing  $j$ . Then  $P_{G-j} = P_G(x) \sum_{i=1}^m \frac{\alpha_{ij}^2}{x - \mu_i}$ , where  $m$  is the number of distinct eigenvalues of graph  $G$ .

**Theorem 2.11.** [6] Suppose  $G_1$  and  $G_2$  are two graphs. If  $H_1 \leq \text{Aut}(G_1)$  and  $H_2 \leq \text{Aut}(G_2)$ , then  $H_1 \times H_2 \leq \text{Aut}(G_1 \times G_2)$ . Also, if  $|d(x_i) - d(x_j)| \neq |n_1 - n_2|$ ,  $i = 1, 2, \dots, n_1$  and  $j = 1, 2, \dots, n_2$  then  $\text{Aut}(G_1 \nabla G_2) = \text{Aut}(G_1) \times \text{Aut}(G_2)$ .

**Lemma 2.12.** [11] Let  $G \neq K_1$  be connected with  $P(G) = \sum_{i=0}^n a_i t^{n-i}$  and  $\lambda = \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n = \Lambda$ , where  $P_G$  is characteristic polynomial of graph  $G$  and  $\lambda_i$  ( $1 \leq i \leq n$ ) is eigenvalue of  $G$ . The following are equivalent:

- (i)  $G$  is bipartite.
- (ii)  $a_{2i-1} = 0$  for all  $1 \leq i \leq \lceil \frac{n}{2} \rceil$ .
- (iii)  $\lambda_i = -\lambda_{n+1-i}$  for  $1 \leq i \leq n$ .
- (iv)  $\Lambda = -\lambda$ .

Moreover,  $m(\lambda_i) = m(-\lambda_i)$ , where  $m(\lambda_i)$  denotes the multiplicity of  $\lambda_i$ .

**Theorem 2.13.** [2] The automorphism group of  $n$  disjoint copies of a graph  $\Gamma$  is  $\text{Aut}(n\Gamma) = S_n[\text{Aut}(\Gamma)] = \text{Aut}(\Gamma) \wr_I S_n$ , where  $I = \{1, 2, \dots, n\}$ .

In the following, we always suppose that  $w$  and  $n$  are natural numbers. Also,  $CP(n)$  denotes cocktail party graph; that is,  $CP(n) = \underbrace{K_{2, 2, \dots, 2}}_{n \text{ times}}$ .

### 3. Main Results

The purpose of this section is to show multicone graphs  $K_w \nabla CP(n)$  are DS with respect to their adjacency spectrums.

#### 3.0.1. Connected graphs cospectral with multicone graphs $K_w \nabla CP(n)$ .

**Proposition 3.1.** *Let  $G$  be a graph. If  $Spec(G) = spec(K_w \nabla CP(n))$ , then  $Spec(G) = \left\{ \left[ \frac{\Omega + \sqrt{\Omega^2 - 4\Gamma}}{2} \right]^1, [-1]^{w-1}, [0]^n, [-2]^{n-1}, \left[ \frac{\Omega - \sqrt{\Omega^2 - 4\Gamma}}{2} \right]^1 \right\}$ , where  $\Omega = 2n + w - 3$  and  $\Gamma = (w - 1)(2n - 2) - 2nw$ .*

*Proof.* Theorem 2.2 together with  $Spec(CP(n)) = \{[2n - 2]^1, [0]^n, [-2]^{n-1}\}$  complete the proof.  $\square$

In the next lemma we show that if  $n \neq 1$ , then any graph cospectral with multicone graph  $K_w \nabla CP(n)$  must be bidegreed.

**Lemma 3.2.** *If  $G$  be a graph and  $Spec(G) = Spec(K_w \nabla CP(n))$ , then  $G$  is either regular or bidegreed, in which case any vertex of  $G$  is of degree  $2n + w - 2$  or  $2n + w - 1$ .*

*Proof.* If  $n = 1$ , then  $G$  is regular. Hence we suppose that  $n \neq 1$ . It is obvious that  $G$  cannot be regular, since regularity of a graph can be determined by its spectrum. By contrary, we suppose that the sequence of degrees of vertices of graph  $G$  consists of at least three number. Hence the equality in Theorem 2.5 cannot happen for any  $\delta$ . But, if we put  $\delta = w + 2n - 2$ , then the equality in Theorem 2.5 holds. So,  $G$  must be bidegreed. Now, we show that  $\Delta = \Delta(G) = w + 2n - 1$ . By contrary, we suppose that  $\Delta < w + 2n - 1$ . Therefore, the equality in theorem 2.5 cannot hold for any  $\delta$ . But, if we put  $\delta = w + 2n - 2$ , then this equality holds. This is a contradiction and so  $\Delta = w - 1 + 2n$ . Now,  $\delta = w + 2n - 2$ , since  $G$  is bidegreed and  $G$  has  $w + 2n$  vertices,  $\Delta = w - 1 + 2n$  and  $w(w - 1 + 2n) + 2n(w + 2n - 2) = w\Delta + 2n(w + 2n - 2) = \sum_{i=1}^{2n+w} \deg v_i$ . This completes the proof.  $\square$

In the next lemma we show that multicone graphs  $K_1 \nabla CP(n)$  are DS.

**Lemma 3.3.** *Any graph cospectral with multicone graph  $K_1 \nabla CP(n)$  is DS.*

*Proof.* By Lemma 3.2,  $G$  has one vertex of degree  $2n$ , say  $j$ . Now,  $P_G = (x - \mu_4)^n(x + 2)^{n-1}(x^2 - (2n - 2)x - 2n)$ . On the other hand, from Theorem 2.10, it follows that  $P_{G-j} = (x - \mu_4)^{n-1}(x - \mu_3)^{n-2}[\alpha_{1j}^2 A + \alpha_{2j}^2 B + \alpha_{3j}^2 C + \alpha_{4j}^2 D]$ , where  
 $A = (x - \mu_2)(x - \mu_3)(x - \mu_4)$ ,  
 $B = (x - \mu_1)(x - \mu_3)(x - \mu_4)$ ,




FIGURE 1. Multicone graph  $\Gamma = K_{20} \nabla CP(2)$ .

$$C = (x - \mu_1)(x - \mu_2)(x - \mu_3),$$

$$D = (x - \mu_1)(x - \mu_2)(x - \mu_3),$$

where  $\mu_1 = \frac{n - 1 + \sqrt{n^2 + 1}}{2}$ ,  $\mu_2 = \frac{n - 1 - \sqrt{n^2 + 1}}{2}$ ,  $\mu_3 = -2$  and  $\mu_4 = 0$ .

We know that  $P_{G-j}$  has  $2n$  roots. Hence we have:

$$x + y + 2n - 2 = (2 - n)\mu_3,$$

$$x^2 + y^2 + (2n - 2)^2 = 4n^2 - 4n - [(n - 2) \cdot \mu_2^2],$$

$$x^3 + y^3 + (2n - 2)^3 = 48 \binom{n}{3} - [(n - 2) \cdot \mu_2^3],$$

where  $x$  and  $y$  are eigenvalues of  $P_{G-j}$ . If we solve the above equations, we obtain  $x = 0$  and  $y = -2$ . So,  $Spec(G - j) = \{[2n - 2]^1, [0]^n, [-2]^{n-1}\}$ . We know that graph  $G - j$  is a regular graph such that degree of its regularity is  $2n - 2$ . Theorem 2.4 implies that  $G$  is a complete multipartite graph. Hence  $G - j \cong \underbrace{K_{2, 2, \dots, 2}}_{n \text{ times}} = CP(n)$  and so  $G \cong K_1 \nabla CP(n)$ . This completes the proof.  $\square$

Hitherto, we have shown multicone graphs  $K_1 \nabla CP(n)$  are DS. The natural questions is; what happens for multicone graphs  $K_w \nabla CP(n)$ ? We answer this question in the next theorem.




FIGURE 2. Multicone graph  $\pi = K_8 \nabla C_4$ .

**Theorem 3.4.** *Multicone graphs  $K_w \nabla CP(n)$  are DS with respect to their adjacency spectrums.*

*Proof.* We solve the problem by induction on  $w$ . If  $w = 1$ , there is nothing for proof. Let the claim be true for  $w$ ; that is, if  $Spec(G_1) = Spec(K_w \nabla CP(n))$ , then  $G_1 \cong K_w \nabla CP(n)$ , where  $G_1$  is a graph. We show that  $Spec(G) = Spec(K_{w+1} \nabla CP(n))$  follows that  $K_w \nabla CP(n)$ , where  $G$  is a graph. By lemma 3.2,  $G = K_1 \nabla G_1$ . This completes the proof.  $\square$

**Corollary 3.5.** *Multicone graphs  $K_w \nabla C_4 = K_w \nabla CP(2)$  are DS with respect to their adjacency spectrums.*

In the following section we show that multicone graphs  $K_w \nabla CP(n)$  are DS with respect to their Laplacian spectrums.

#### 4. CONNECTED GRAPHS COSPECTRAL WITH MULTICONE GRAPH $K_w \nabla CP(n)$ WITH RESPECT TO LAPLACIAN SPECTRUMS

**Theorem 4.1.** *Multicone graphs  $K_w \nabla CP(n)$  are DS with respect to their Laplacian spectra.*

*Proof.* We solve the problem by induction on  $w$ . If  $w = 1$ , there is nothing to prove. Let the claim be true for  $w$ ; that is, if  $Spec(L(G_1)) = Spec(L(K_w \nabla CP(n))) = \{[w + 2n]^{w-1+n}, [w + 2n - 2]^n, [0]^1\}$ , then  $G_1 \cong K_w \nabla CP(n)$ , where  $G_1$  is a graph. We




FIGURE 3. Multicone graph  $\beta = K_{10} \nabla CP(3)$ .

show that from  $Spec(K_{w+1} \nabla CP(n)) = \{[w + 2n + 1]^{w+n}, [w + 2n - 1]^n, [0]^1\}$  follows that  $G \cong K_{w+1} \nabla CP(n)$ . We conclude from Lemma 2.8 that  $G_1$  and  $G$  are join of two graphs. On the other hand  $G$  has one vertex and  $2n + w$  edges more than  $G_1$ . Hence  $G$  has one vertex of degree  $2n + w$ , say  $j$ . So,  $Spec(L(G - j)) = Spec(K_w \nabla CP(n))$ . Hence  $G - j \cong K_w \nabla CP(n)$  and so  $G \cong K_1 \nabla (K_w \nabla CP(n))$ . This completes the proof.  $\square$

**Corollary 4.2.** *Multicone graphs  $K_w \nabla C_4$  are DS with respect to their Laplacian spectrums.*

In the next section we show that split graphs  $\overline{K_w \nabla CP(n)}$  are DS with respect to their adjacency spectrums.

5. SPLIT GRAPHS WHOSE ARE COSPECTRAL WITH COMPLEMENT OF MULTICONE GRAPHS

$$K_w \nabla CP(n).$$

**Proposition 5.1.** *Let  $G$  be cospectral with adjacency spectrum of complement of multicone graph  $K_w \nabla CP(n)$ . Then  $Spec(G) = \{[1]^n, [0]^w, [-1]^n\}$ .*

*Proof.* Straightforward.  $\square$

**Theorem 5.2.** *If  $Spec(G) = Spec(\overline{K_w \nabla CP(n)})$ , then  $G \cong \overline{K_w \nabla CP(n)}$ .*






FIGURE 4. Multicone graph  $\Theta = K_{15} \nabla CP(2)$

*Proof.* If  $n = 1$ , there is nothing to prove. Hence we suppose that  $n \neq 1$ . It is obvious that  $G$  cannot be regular, since regularity of a graph can be determined by its spectrum. Firstly, we show that  $G$  is disconnected. By contrary, we suppose that  $G$  is connected. So, by Lemma 2.6,  $G$  must be a complete bipartite graph and so  $G \cong K_2$ . This is a contradiction. Hence  $G = G_1 \cup G_2 \cup \dots \cup G_k$ , where  $G_i$  is a connected component of  $G$  and  $1 \leq i \leq k$ . Now, we show that  $G_i$  cannot have three distinct eigenvalues. By contrary, we suppose that  $G_i$  has three distinct eigenvalues. We show that  $G_i$  must be regular. By contrary, we assume that  $G_i$  is non-regular. Hence Theorem 2.6 implies that  $G_i$  is a complete bipartite graph. So,  $G_i \cong K_2$ . This is a contradiction. Therefore,  $G_i$  is a regular graph and also multiplicity of eigenvalue 1, is 1, since  $G_i$  is connected. Now, from Lemma 2.6, it follows that  $G_i$  is a complete multipartite and so  $G_i = mK_2$ , where  $m \geq 1$ . This implies that  $G_i \cong K_2$ . This is a contradiction. Therefore  $G_i$  has one or two eigenvalue(s). Hence, any of connected components of  $G$  is either  $K_1$  or a complete graph. Hence  $G \cong wK_1 \cup nK_2$ . This completes the proof.  $\square$

In the following, we show that any graph cospectral with multicone graph  $K_w \nabla CP(n)$  must be perfect. Also, we show that any graph cospectral with multicone graph  $K_w \nabla CP(n)$  with respect to Laplacian spectrum is perfect. In addition, we prove that any graph cospectral

with split graph  $\overline{K_w \nabla CP(n)}$  must be perfect. Finally, we obtain automorphism group of multicone graphs  $K_w \nabla CP(n)$ .

#### 6. SOME OF ALGEBRAIC PROPERTIES ABOUT MULTICONE GRAPHS $K_w \nabla CP(n)$ .

Suppose  $\chi(G)$  and  $\omega(G)$  are chromatic number and clique number of graph  $G$ , respectively. A graph is **perfect** if  $\chi(H) = \omega(H)$  for every induced subgraph  $H$  of  $G$ . It is proved that a graph  $G$  is perfect if and only if  $G$  is Berge; that is, it contains no odd hole or antihole as induced subgraph, where odd hole and antihole are odd cycle,  $C_m$  for  $m \geq 5$ , and its complement, respectively. Also, in 1972 Lovász proved that, A graph is perfect if and only if its complement is perfect [1].

**Theorem 6.1.** *Let graph  $G$  be cospectral with multicone graph  $K_w \nabla CP(n)$ . Then  $G$  and  $\overline{G}$  are perfect.*

*Proof.* Straightforward.  $\square$

**Theorem 6.2.** *Let  $\text{Spec}(L(G)) = \text{Spec}(L(K_w \nabla CP(n)))$ . Then  $G$  and  $\overline{G}$  are perfect.*

*Proof.* Straightforward.  $\square$

**Theorem 6.3.** *Let graph  $G$  be cospectral with graph  $\overline{K_w \nabla CP(n)}$ . Then  $G$  and  $\overline{G}$  are perfect.*

*Proof.* Straightforward.  $\square$

**Proposition 6.4.**  *$\text{Aut}(K_w \nabla CP(n)) \cong S_w \times [S_2 \text{ wr } S_n]$ , where  $I = \{1, 2, \dots, n\}$ .*

*Proof.* It is clear that  $\text{Aut}(CP(n)) \cong \text{Aut}(nK_2)$ . Now, Theorem 2.11 together with Theorem 2.13 complete the proof.  $\square$

In the following, we pose one conjecture.

## 7. COROLLARIES AND CONJECTURE

**Corollary 7.1.** *Let  $G$  be a graph. The following statements are equivalent:*

- (i)  $G \cong K_w \nabla CP(n)$ .
- (ii)  $\text{Spec}(G) = \text{Spec}(K_w \nabla CP(n))$ .
- (iii)  $\text{Spec}(L(G)) = \text{Spec}(L(K_w \nabla CP(n)))$ .

**Corollary 7.2.** *Let  $G$  be a graph. The following statements are equivalent:*

- (i)  $G \cong \overline{K_w \nabla CP(n)}$ .
- (ii)  $\text{Spec}(G) = \text{Spec}(\overline{K_w \nabla CP(n)})$ .
- (iii)  $\text{Spec}(L(G)) = \text{Spec}(L(\overline{K_w \nabla CP(n)}))$ .

**Conjecture 7.3.** *Multicone graphs  $K_w \nabla CP(n)$  are DS with respect to their signless Laplacian spectrums.*

## 8. ACKNOWLEDGMENTS

The authors wish to sincerely thank the referees for several useful comments.

## REFERENCES

- [1] A. Abdollahi, S. Janbaz and M. Oubodi, *Graphs Cospectral with A Friendship Graph Or its Complement*, Trans. Combin. Vol. 2 No. 4 (2013), pp. 37–52.
- [2] N. L. Biggs, *Algebraic Graph Theory*, (second edition), Cambridge University press, cambridge, (1933).
- [3] X. M. Cheng, G. R. W. Greaves, J. H. Koolen, *Graphs with three eigenvalues and second largest eigenvalue at most 1*, <http://de.arxiv.org/abs/1506.02435v1>.
- [4] D. Cvetković, P. Rowlinson and S. Simić, *An Introduction to the theory of graph spectra*, London Mathematical Society Student Texts, **75**, Cambridge University Press, Cambridge, 2010.
- [5] J. D. Dixon and B. Mortimer, *Permutation Groups*, Math. Proc. Cambridge Phil. Soc. (1998).
- [6] G. H. Fath-Tabar, *The Automorphism Group Of Finite Graphs*, Iran. J. Math. Sci. Inf., (2007), 29–33.
- [7] A. Ganasem, *Automorphism groups of graphs*, ArXiv: 1206.6279v1 (2012).
- [8] A. Ganasem, *Automorphism of Cayley graph generated by transposition sets*, arXiv 1303.5974v2.

- [9] C. D. Godsil, *On the full automorphism group of a graph*, *Combinatorica*, **1** (1981) 243–256.
- [10] C. Godsil and G. Royle, *Algebraic Graph Theory*, Springer-Verlag, New York, 2001.
- [11] U. Kanauer, *Algebraic Graph Theory, Morphism, Monoids and Matrices*, (2011).
- [12] B. Liu, *On an upper bound of the spectral radius of graphs*, *Discrete Mathematics* **308** (2008) 5317–5324.
- [13] R. Merris, *Laplacian graph eigenvectors*, *Linear Algebra Appl.* **278** (1998), 221–236.
- [14] A. Mohammadian, B. Tayfeh Rezaie, *Graphs with four distinct Laplacian eigenvalues*, *J. Algebraic Combin.* **34** (2011), 671–682.
- [15] G. R. Omid, *On graphs with largest Laplacian eigenvalues at most 4*, *Australas. J. Combin.*, **44** (2009) 163–170.
- [16] E. R. van Dam and W. H. Haemers, *Which Graphs are determined by their spectrum?*, *Linear Alg. Appl.* **373** (2003) 241–272.
- [17] E. R. van Dam, Willem H. Haemers, *Developments on spectral characterizations of graphs*, *Discrete Math.*, **309** (2009) 576–586.
- [18] J. F. Wang, H. Zhao, Q. Haung, *Spectral Characterization of Multicone Graphs*. *Czec. Math. J.*, **62** **137** (2012), 117–126.
- [19] J. Wang, Q. Huang, *Spectral Characterization of Generalized Cocktail-Party Graphs*. *J. Math. Research Appl.*, (2012), Vol. **32**, No. 6, pp. 666–672.
- [20] D.B. West, *Introduction to Graph Theory* (Second Edition), University of Illinois—Urbana (2001).

**Ali Zeydi Abdian**

Department of mathematics, College of science

Lorestan university, Lorestan

Khoramabad, Iran.

azeydiabdi@gmail.com

**S. Morteza Mirafzal**

Department of mathematics, College of science

Lorestan university, Lorestan

Khoramabad, Iran.

mirafzal.m@lu.ac.ir