



Research Paper

CONNECTIONS BETWEEN GE ALGEBRAS AND PRE-HILBERT ALGEBRAS

ANDRZEJ WALENDZIAK*

ABSTRACT. GE algebras (generalized exchange algebras) and pre-Hilbert algebras are a generalization of well-known Hilbert algebras. In the paper, connections between these algebras are studied. In particular, it is proven that π -BE algebras are equivalent with GE algebras satisfying the exchange property. Some characterizations of transitive GE algebras and exchange pre-Hilbert algebras are given. It is shown that the intersection of classes of GE algebras and pre-Hilbert algebras is the class of transitive GE algebras. Moreover, GE, BE and pre-Hilbert algebras with the antisymmetry property are investigated. It is proven that transitive GE algebras satisfying the property of antisymmetry coincide with Hilbert algebras. Finally, positive implicative GE and pre-Hilbert algebras are considered, their connections with some algebras of logic are presented. In addition, the hierarchies existing between the classes of algebras studied here are shown.

DOI: 10.22034/as.2023.20655.1678

MSC(2010): Primary: 03G25; Secondary: 06A06, 06F35.

Keywords: BE algebra, GE algebra, Hilbert algebra, Pre-Hilbert algebra.

Received: 22 September 2023, Accepted: 14 November 2023.

*Corresponding author

1. INTRODUCTION

L. Henkin [8] introduced the notion of “implicative model”, as a model of positive implicative propositional calculus. In 1960, Monteiro [15] has given the name “Hilbert algebras” to the dual algebras of Henkin’s implicative models. In 1966, Iséki [11] introduced a new notion called a BCK algebra. It is an algebraic formulation of the BCK-propositional calculus system of Meredith [14], and generalize the concept of implicative algebras (see [1]). In 2021, Cintula and Noguera [6] presented of the most important logics that one can find in the literature. In particular, they considered the *BCK logic* and its many extensions. In [13], as a generalization of BCK algebras, Kim and Kim introduced BE algebras. Rezaei et al. [16] investigated connections between Hilbert algebras and BE algebras. In 2008, Walendziak [20] defined commutative BE algebras and proved that they are BCK algebras. Later on, in 2010, Buşneag and Rudeanu [5] introduced the notion of pre-BCK algebra. A BCK algebra is just a pre-BCK algebra satisfying also the antisymmetry. In 2016, Iorgulescu [9] introduced new generalizations of BCK and Hilbert algebras (RML, aBE, pi-BE, piml-RML algebras and many others). Bandaru et al. [2] introduced the concepts of GE algebra (generalized exchange algebra) and transitive GE algebra (tGE algebra for short). They studied the various properties and filter theory of GE algebras. Some types of filters (imploring, prominent, belligerent, interior, voluntary GE-filters) have been widely investigated and many important results are obtained (see [18, 17, 3, 19, 4]). Recently, as a natural generalization of Hilbert algebras, Walendziak [22] introduced pre-Hilbert algebras (the definition of a pre-Hilbert algebra is inspired by Henkin’s Positive Implicative Logic [8]) and studied their properties.

In the paper, we consider connections between GE algebras, BE algebras and pre-Hilbert algebras. In particular, we prove that pi-BE algebras are equivalent with GE algebras satisfying the exchange property. We give some characterizations of transitive GE algebras and show that the class of these algebras is the intersection of classes of GE algebras and pre-Hilbert algebras. We introduce and study exchange pre-Hilbert algebras. Moreover, we consider GE, BE and pre-Hilbert algebras with the antisymmetry property and give some examples. We also prove that tGE algebras verifying the property of antisymmetry are equivalent with Hilbert algebras. Finally, we present the interrelationships between the classes of algebras considered here.

The motivation of this study consists algebraic and logical arguments. Namely, generalized exchange algebras and pre-Hilbert algebras are related to Henkin’s Positive Implicative Logic, they belong to a wide class of algebras of logic. An additional motivation is the fact that the present paper is a continuation of previous papers: [2, 21] on GE algebras and [22] on pre-Hilbert algebras. Moreover, the results of the paper may have applications for future studies of the relationships between some generalizations of Hilbert algebras.

2. PRELIMINARIES

Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type $(2, 0)$. We consider the following list of properties ([9]) that can be satisfied by \mathcal{A} (the properties in the list are the most important properties satisfied by a Hilbert algebra):

- (An) (Antisymmetry) $x \rightarrow y = 1 = y \rightarrow x \implies x = y$,
- (B) $(y \rightarrow z) \rightarrow [(x \rightarrow y) \rightarrow (x \rightarrow z)] = 1$,
- (C) $[x \rightarrow (y \rightarrow z)] \rightarrow [y \rightarrow (x \rightarrow z)] = 1$,
- (Ex) (Exchange) $x \rightarrow (y \rightarrow z) = y \rightarrow (x \rightarrow z)$,
- (GE) (Generalized exchange) $x \rightarrow (y \rightarrow z) = x \rightarrow [y \rightarrow (x \rightarrow z)]$,
- (K) $x \rightarrow (y \rightarrow x) = 1$,
- (L) (Last element) $x \rightarrow 1 = 1$,
- (M) $1 \rightarrow x = x$,
- (Re) (Reflexivity) $x \rightarrow x = 1$,
- (Tr) (Transitivity) $x \rightarrow y = 1 = y \rightarrow z \implies x \rightarrow z = 1$,
- (pi) $x \rightarrow y = x \rightarrow (x \rightarrow y)$,
- (p-1) $x \rightarrow (y \rightarrow z) \leq (x \rightarrow y) \rightarrow (x \rightarrow z)$,
- (p-2) $(x \rightarrow y) \rightarrow (x \rightarrow z) \leq x \rightarrow (y \rightarrow z)$,
- (pimpl) $x \rightarrow (y \rightarrow z) = (x \rightarrow y) \rightarrow (x \rightarrow z)$.

Lemma 2.1. *Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type $(2, 0)$. Then the following hold:*

- (i) $(M) + (B)$ imply (Re) , (Tr) ;
- (ii) $(An) + (C)$ imply (Ex) ;
- (iii) $(Re) + (L) + (Ex)$ imply (K) ;
- (iv) $(M) + (L) + (B)$ imply (K) ;
- (v) $(Re) + (M) + (pimpl)$ imply (pi) ;
- (vi) $(Ex) + (pi)$ imply (GE) ;
- (vii) $(B) + (Ex) + (pi)$ imply $(p-1)$.

Proof. (i) – (v) follow from Propositions 2.1 and 6.4 of [9].

(vi) follows from Proposition 2.7 of [21].

(vii). Let $x, y, z \in A$. By (Ex), $x \rightarrow (y \rightarrow z) = y \rightarrow (x \rightarrow z)$. Applying (B) and (pi), we get $y \rightarrow (x \rightarrow z) \leq (x \rightarrow y) \rightarrow [x \rightarrow (x \rightarrow z)] = (x \rightarrow y) \rightarrow (x \rightarrow z)$. Then $x \rightarrow (y \rightarrow z) \leq (x \rightarrow y) \rightarrow (x \rightarrow z)$, that is, (p-1) holds. \square

Following Iorgulescu [9], we say that $(A, \rightarrow, 1)$ is an *RML algebra* if it verifies the axioms (Re), (M), (L). We recall now the following definition.

Definition 2.2. ([9]) Let $\mathcal{A} = (A, \rightarrow, 1)$ be an RML algebra. The algebra \mathcal{A} is said to be:

1. a *BE algebra* if it verifies (Ex),
2. an *aRML algebra* if it verifies (An),
3. an *aBE algebra* if it verifies (Ex) and (An), that is, it is a BE algebra with (An),
4. a *pre-BCK algebra* if it verifies (B) and (Ex), that is, it is a BE algebra with (B),
5. a *BCK algebra* if it is a pre-BCK algebra verifying (An).

Denote by **RML**, **BE**, **aRML**, **aBE**, **pre-BCK** and **BCK** the classes of RML, BE, aRML, aBE, pre-BCK and BCK algebras, respectively.

By definition, **BE** = **RML** + (Ex), **aRML** = **RML** + (An), **aBE** = **BE** + (An), **pre-BCK** = **BE** + (B), **BCK** = **pre-BCK** + (An).

Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type $(2, 0)$. We define the binary relation \leq by: for all $x, y \in A$,

$$x \leq y \iff x \rightarrow y = 1.$$

It is known that \leq is an order relation in BCK algebras. By definition, in RML and BE algebras, \leq is a reflexive relation; in aRML and aBE algebras, \leq is reflexive and antisymmetric. Since (M) + (B) imply (Tr), see Lemma 2.1 (i), in pre-BCK algebras, \leq is reflexive and transitive (i.e., it is a pre-order relation).

Definition 2.3. ([2]) Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type $(2, 0)$. We say that \mathcal{A} is:

6. a *GE algebra* (generalized exchange algebra) if it verifies (Re), (M), (GE);
7. a *transitive GE algebra* (tGE algebra, for short) if it is a GE algebra verifying (B);
8. an *antisymmetric GE algebra* (aGE algebra, for short) if it is a GE algebra verifying (An).

Denote by **GE**, **tGE** and **aGE** the classes of all GE algebras, transitive GE algebras and aGE algebras, respectively.

Proposition 2.4. (Corollary 3.2 of [21]) Any GE algebra satisfies (Re), (M), (L), (C), (K), (GE), (pi).

Remark 2.5. Since GE algebras satisfy (L), we get **GE** = **RML** + (GE). By definition, **tGE** = **GE** + (B) and **aGE** = **GE** + (An).

Remark 2.6. GE algebras do not have to satisfy (An), (B), (Tr), (Ex), (p-1), (pimpl); see example below.

Example 2.7. ([21]) Consider the set $A = \{a, b, c, d, e, 1\}$ and the operation \rightarrow given by the following table:

| | | | | | | |
|---------------|-----|-----|-----|-----|-----|-----|
| \rightarrow | a | b | c | d | e | 1 |
| a | 1 | 1 | c | c | 1 | 1 |
| b | a | 1 | d | d | 1 | 1 |
| c | a | 1 | 1 | 1 | 1 | 1 |
| d | a | 1 | 1 | 1 | 1 | 1 |
| e | a | 1 | 1 | 1 | 1 | 1 |
| 1 | a | b | c | d | e | 1 |

We can observe that the properties (Re), (M), (L), (GE) (hence (pi)) are satisfied. Therefore, $(A, \rightarrow, 1)$ is a GE algebra. It does not satisfy (An) for $(x, y) = (c, d)$; (Ex) for $(x, y, z) = (a, b, c)$; (Tr), (B), (p-1), (pimpl) for $(x, y, z) = (a, e, c)$.

Proposition 2.8. (Corollary 3.12 of [21]) Any transitive GE algebra verifies (Re), (M), (L), (B), (Tr), (GE), (pi), (p-1), (p-2).

Remark 2.9. Transitive GE algebras do not have to verify (An), (Ex), (pimpl); see example below.

Example 2.10. Let $A = \{a, b, c, 1\}$ and \rightarrow be defined as follows:

| | | | | | |
|---------------|-----|-----|-----|-----|-----|
| \rightarrow | a | b | c | d | 1 |
| a | 1 | 1 | c | c | 1 |
| b | a | 1 | d | d | 1 |
| c | a | 1 | 1 | 1 | 1 |
| d | a | 1 | 1 | 1 | 1 |
| 1 | a | b | c | d | 1 |

It is easy to see that $\mathcal{A} = (A, \rightarrow, 1)$ is a transitive GE algebras. It does not verify (An) for $x = c, y = d$; (Ex) and (pimpl) for $x = b, y = a, z = c$.

In [22], we introduced the following notion:

Definition 2.11. A pre-Hilbert algebra is an algebra $(A, \rightarrow, 1)$ of type $(2, 0)$ satisfying (M), (K) and (p-1).

Recall that an algebra $(A, \rightarrow, 1)$ is a *Hilbert algebra* ([7]) if it verifies the axioms (An), (K), (p-1).

Remark 2.12. In [7], A. Diego proved that Hilbert algebras satisfy (Re), (M), (L), (B), (Ex), (pi), (p-2), (pimpl). Moreover, he showed that the class of all Hilbert algebras is a variety.

Let us denote by **pre-H** and **H** the classes of pre-Hilbert and Hilbert algebras, respectively.

Remark 2.13. Since $(An) + (K) + (p-1)$ imply (M) (see [7]), a Hilbert algebra is in fact a pre-Hilbert algebra verifying (An) , that is, $\mathbf{H} = \mathbf{pre-H} + (An)$.

Proposition 2.14. (Theorem 3.9 of [22]) *Pre-Hilbert algebras satisfy (Re) , (M) , (L) , (K) , (Tr) , (B) , (C) , $(p-1)$, $(p-2)$.*

Proposition 2.15. *Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type $(2, 0)$. The following are equivalent:*

- (i) \mathcal{A} is a pre-Hilbert algebra;
- (ii) \mathcal{A} satisfies (M) , (L) , (B) and $(p-1)$.

Proof. (i) \implies (ii). Follows from Proposition 2.14.

(ii) \implies (i). By Lemma 2.2 (iv), $(M) + (L) + (B)$ imply (K) . Then \mathcal{A} satisfies (M) , (K) and $(p-1)$. Thus \mathcal{A} is a pre-Hilbert algebra. \square

Remark 2.16. From Proposition 2.15 we see that $\mathbf{pre-H} = \mathbf{RML} + (B) + (p-1)$.

Remark 2.17. Pre-Hilbert algebras do not have to satisfy (An) , (Ex) , (GE) , (pi) , $(pimpl)$; see example below.

Example 2.18. Consider the set $A = \{a, b, c, d, 1\}$ and the operation \rightarrow given by the following table:

| | | | | | |
|---------------|-----|-----|-----|-----|-----|
| \rightarrow | a | b | c | d | 1 |
| a | 1 | c | b | d | 1 |
| b | a | 1 | 1 | d | 1 |
| c | a | 1 | 1 | d | 1 |
| d | a | c | c | 1 | 1 |
| 1 | a | b | c | d | 1 |

We can observe that the properties (M) , (K) , $(p-1)$ are verified. Then, $(A, \rightarrow, 1)$ is a pre-Hilbert algebra. It does not verify (An) for $(x, y) = (b, c)$; (Ex) and $(pimpl)$ for $(x, y, z) = (a, d, b)$; (pi) for $(x, y) = (a, b)$; (GE) for $(x, y, z) = (a, 1, b)$.

Remark 2.19. It is easy to see that in GE algebras, \leq is a reflexive relation; in tGE and pre-Hilbert algebras, \leq is reflexive and transitive (i.e., it is a pre-order relation), in aGE algebras, \leq is reflexive and antisymmetric. In Hilbert algebras, \leq is an order relation.

3. CONNECTIONS BETWEEN GE , BE ALGEBRAS AND PRE-HILBERT ALGEBRAS

We recall the following definitions:

Definition 3.1. ([9]) **1.** A pi-RML algebra is an RML algebra verifying property (pi).
2. A positive implicative RML algebra, or a pimpl-RML algebra for short, is an RML algebra verifying property (pimpl).

Denote by **pi-RML** and **pimpl-RML** the classes of pi-RML and pimpl-RML algebras, respectively; similarly for the subclasses of **RML**.

By definition, **pi-RML** = **RML** + (pi), **pi-BE** = **BE** + (pi), **pi-pre-BCK** = **pre-BCK** + (pi) = **pi-BE** + (B), **pimpl-RML** = **RML** + (pimpl).

Remark 3.2. From Lemma 2.1 (v) it follows that in RML algebras, (pimpl) implies (pi). For BCK algebras, (pimpl) and (pi) are equivalent (cf. Theorem 8 of [12]).

By definition and Proposition 2.4, we have

Proposition 3.3. Any GE algebra is a pi-RML algebra.

From Lemma 2.1 (vi) we get

Proposition 3.4. Any pi-BE algebra is a GE algebra.

Proposition 3.5. Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type (2, 0). The following statements are equivalent:

- (i) \mathcal{A} is a pi-BE algebra,
- (ii) \mathcal{A} is a GE algebra satisfying (Ex).

Proof. (i) \implies (ii). Let \mathcal{A} be a pi-BE algebra. It is sufficient to show that \mathcal{A} satisfies (GE). Let $x, y, z \in A$. We obtain

$$x \rightarrow (y \rightarrow z) \stackrel{(pi)}{=} x \rightarrow [x \rightarrow (y \rightarrow z)] \stackrel{(Ex)}{=} x \rightarrow [y \rightarrow (x \rightarrow z)].$$

Thus (GE) holds.

(ii) \implies (i). Since GE algebras satisfy (pi), we see that (ii) implies (i). \square

Now we give a characterization of transitive GE algebras.

Theorem 3.6. Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type (2, 0). The following are equivalent:

- (i) \mathcal{A} is a transitive GE algebra;
- (ii) \mathcal{A} satisfies (M), (GE), (B);
- (iii) \mathcal{A} satisfies (Re), (M), (GE) and (p-1);
- (iv) \mathcal{A} is a pre-Hilbert algebra with (GE).

Proof. (i) \implies (ii). Obviously, by definition.

(ii) \implies (iii). By Lemma 2.1 (i), (M) + (B) imply (Re). Consequently, \mathcal{A} is a tGE algebra. From Proposition 2.8 it follows that \mathcal{A} satisfies (p-1).

(iii) \implies (iv). Let \mathcal{A} satisfy (Re), (M), (GE) and (p-1). Then \mathcal{A} is a GE algebra. By Proposition 2.4, \mathcal{A} satisfies (K). Therefore, (M), (K), (p-1) and (GE) hold in \mathcal{A} . Thus \mathcal{A} is a pre-Hilbert algebra with (GE).

(iv) \implies (i). Since pre-Hilbert algebras satisfy (B), we conclude that (iv) implies (i). \square

From Theorem 3.6 we have

Corollary 3.7. $\mathbf{tGE} = \mathbf{GE} \cap \mathbf{pre-H}$.

We now introduce a new algebra. We say that an algebra $\mathcal{A} = (A, \rightarrow, 1)$ is an *exchange pre-Hilbert algebra* if it is a pre-Hilbert algebra verifying (Ex).

Denote by **Ex-pre-H** the class of exchange pre-Hilbert algebras. By definition, **Ex-pre-H** = **pre-H** + (Ex).

Example 3.8. (Example 8.20 of [10]) Consider the set $A = \{a, b, c, 1\}$ with the following table of \rightarrow :

| | | | | |
|---------------|-----|-----|-----|-----|
| \rightarrow | a | b | c | 1 |
| a | 1 | a | a | 1 |
| b | 1 | 1 | 1 | 1 |
| c | 1 | 1 | 1 | 1 |
| 1 | a | b | c | 1 |

The algebra $\mathcal{A} = (A, \rightarrow, 1)$ verifies (Re), (M), (L), (Ex), (B), (p-1). It does not verify (An) for $x = b, y = c$; (pi) for $x = a, y = b$. Hence, \mathcal{A} is an exchange pre-Hilbert algebra, without (pi).

Proposition 3.9. *Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type (2, 0). The following statements are equivalent:*

- (i) \mathcal{A} is an exchange pre-Hilbert algebra,
- (ii) \mathcal{A} is a pre-BCK algebra with (p-1),
- (iii) \mathcal{A} is a BE algebra satisfying (p-1).

Proof. (i) \implies (ii) and (ii) \implies (iii) are obvious.

(iii) \implies (i). By Lemma 2.1 (iii), (Re) + (L) + (Ex) imply (K). Then \mathcal{A} satisfies (M), (K), (p-1), (Ex). Thus \mathcal{A} is an exchange pre-Hilbert algebra. \square

Example 3.10. (Example 9.35 of [10]) Let $A = \{a, b, c, 1\}$ and \rightarrow be defined as follows:

| | | | | |
|---------------|-----|-----|-----|-----|
| \rightarrow | a | b | c | 1 |
| a | 1 | a | b | 1 |
| b | 1 | 1 | 1 | 1 |
| c | 1 | 1 | 1 | 1 |
| 1 | a | b | c | 1 |

Properties (Re), (M), (L), (Ex) hold in $\mathcal{A} = (A, \rightarrow, 1)$. Hence, \mathcal{A} is a BE algebra. It does not verify (An) for $x = b, y = c$; (B) and (p-1) for $x = a, y = b, z = c$; (pi) for $x = a, y = b$. Therefore $\mathbf{Ex - pre - H} \subset \mathbf{BE}$ and $\mathbf{pi - BE} \subset \mathbf{BE}$.

Remark 3.11. (1) By Propositions 3.5, 3.6 and 3.9, $\mathbf{pi - BE} = \mathbf{GE} + (Ex)$, $\mathbf{tGE} = \mathbf{pre - H} + (GE)$, $\mathbf{Ex - pre - H} = \mathbf{pre - BCK} + (p - 1) = \mathbf{BE} + (p - 1)$.

(2) Hence $\mathbf{Ex - pre - H} + (pi) = \mathbf{pre - BCK} + (p - 1) + (pi) = \mathbf{pi - pre - BCK}$, since (B) + (Ex) + (pi) imply (p-1), see Lemma 2.1 (vii).

(3) We have $\mathbf{tGE} + (Ex) = \mathbf{GE} + (B) + (Ex) = \mathbf{pi - BE} + (B) = \mathbf{pi - pre - BCK}$.

The interrelationships between the classes of algebras mentioned before are visualized in Figure 1 (see Remarks 2.5, 2.16, 3.11).




FIGURE 1.

We now consider GE, BE and pre-Hilbert algebras with the antisymmetry. First we give a characterization of antisymmetric GE algebras.

Proposition 3.12. *Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type $(2, 0)$. The following statements are equivalent:*

- (i) \mathcal{A} is an aGE algebra,
- (ii) \mathcal{A} is a pi-aBE algebra,
- (iii) \mathcal{A} is an aRML algebra with (GE).

Proof. (i) \implies (ii). Let \mathcal{A} be a GE algebra with (An). From Proposition 2.4 it follows that \mathcal{A} satisfies (C) and (pi). By Lemma 2.1 (ii), (An) + (C) imply (Ex). Therefore \mathcal{A} is a pi-aBE algebra.

(ii) \implies (iii). To prove (GE), let $x, y, z \in A$. We obtain

$$x \rightarrow (y \rightarrow z) \stackrel{\text{(pi)}}{=} x \rightarrow [x \rightarrow (y \rightarrow z)] \stackrel{\text{(Ex)}}{=} x \rightarrow [y \rightarrow (x \rightarrow z)].$$

Thus (GE) holds. Hence \mathcal{A} is an aRML algebra with (GE).

(iii) \implies (i). It is obvious. \square

Remark 3.13. By Proposition 3.12, $\mathbf{aGE} = \mathbf{pi} - \mathbf{aBE}$.

Proposition 3.14. *Let $\mathcal{A} = (A, \rightarrow, 1)$ be an algebra of type $(2, 0)$. The following statements are equivalent:*

- (i) \mathcal{A} is a pre-Hilbert algebra with (An),
- (ii) \mathcal{A} is a Hilbert algebra,
- (iii) \mathcal{A} is a pi-aBE algebra with (B),
- (iv) \mathcal{A} is an aGE algebra with (B),
- (v) \mathcal{A} is a tGE algebra with (An).

Proof. (i) \iff (ii), by Remark 2.13.

(ii) \implies (iii). Let \mathcal{A} be a Hilbert algebra. Then \mathcal{A} satisfies (An), (Re), (M), (L), (Ex), (B), (pi). Hence \mathcal{A} is a pi-aBE algebra with (B).

(iii) \implies (ii). By Lemma 2.1 (iii) and (vii), (Re) + (L) + (Ex) imply (K) and (B) + (Ex) + (pi) imply (p-1). Hence (iii) forces (ii).

(iii) \iff (iv), by Remark 3.13.

(iv) \iff (v), by definition. \square

Remark 3.15. From Proposition 3.14 it follows that $\mathbf{H} = \mathbf{pre-H} + (\text{An}) = \mathbf{pi-aBE} + (\text{B}) = \mathbf{aGE} + (\text{B}) = \mathbf{tGE} + (\text{An})$.




FIGURE 2.

By Definitions and Remarks 3.13 and 3.15, we can draw Figure 2.

Remark 3.16. (1) Positive implicative GE algebras were studied in [21]. By Remark 4.9 of [21], $\mathbf{pimpl - RML} = \mathbf{pimpl - GE} = \mathbf{pimpl - tGE}$.

(2) Positive implicative pre-Hilbert algebras were studied in [22]. By Proposition 4.7 of [22], $\mathbf{pimpl - pre - H} = \mathbf{pimpl - RML}$. Therefore $\mathbf{pimpl - pre - H} = \mathbf{pimpl - GE} = \mathbf{pimpl - tGE}$.

(3) From Corollary 6.19 of [9] we have $\mathbf{pimpl - BE} = \mathbf{pimpl - pre - BCK} = \mathbf{pimpl - RML} + (Ex)$ and $\mathbf{H} = \mathbf{pimpl - BCK} = \mathbf{pimpl - aRML} = \mathbf{pimpl - aBE}$.

(4) From the above we see that $\mathbf{H} \subset \mathbf{pimpl - BE} \subset \mathbf{pimpl - GE} = \mathbf{pimpl - pre - H}$.

4. SUMMARY AND FUTURE WORK

In this paper, we studied connections between BE algebras, GE algebras and pre-Hilbert algebras. We proved that GE algebras satisfying the exchange property are equivalent with pi-BE algebras and showed that the class of transitive GE algebras is the intersection of classes of GE algebras and pre-Hilbert algebras. We introduced exchange pre-Hilbert algebras and gave some examples and characterizations of these algebras. Moreover, we considered GE, BE and pre-Hilbert algebras verifying the antisymmetry property. In particular, we proved that tGE algebras verifying (An) are equivalent with Hilbert algebras. Finally, we presented the interrelationships between the classes of algebras considered here.

The results obtained in the paper can be a starting point for future research. We suggest the following topics:

- (1) Studying GE and pre-Hilbert algebras with the implicative property, that is, verifying the identity $(x \rightarrow y) \rightarrow x = x$.
- (2) Describing the deductive systems, the congruences, the quotient algebras, etc. of pre-Hilbert algebras and GE algebras.
- (3) Investigating more deeply the exchange pre-Hilbert algebras.

5. ACKNOWLEDGMENTS

The author is indebted to the referees for their very careful reading and suggestions.

REFERENCES

- [1] J. C. Abbott, *Semi-boolean algebras*, Mat. Vesnik, **4** No. 19 (1967) 177-198.
- [2] R. K. Bandaru, A. Borumand Saeid and Y. B. Jun, *On GE-algebras*, Bull. Sect. Logic, **50** No. 1 (2021) 81-96.
- [3] R. K. Bandaru, A. Borumand Saeid and Y. B. Jun, *Belligerent GE-filters in GE-algebras*, J. Indones. Math. Soc., **28** (2021) 31-43.
- [4] A. Borumand Saeid, A. Rezaei, R. K. Bandaru and Y. B. Jun, *Voluntary GE-filters and further results of GE-filters in GE-algebras*, J. Algebr. Syst., **10** No. 1 (2022) 31-47.
- [5] D. Busneag and S. Rudeanu, *A glimpse of deductive systems in algebra*, Cent. Eur. J. Math., **8** No. 4 (2010) 688-705.
- [6] P. Cintula and C. Noguera, *Logic and Implication: An Introduction to the General Algebraic Study of Non-Classical Logics*, Trends in Logic 57, Springer, Berlin, 2021.
- [7] A. Diego, *Sur les algèbres de Hilbert*, Collection de Logique Mathématique, Edition Hermann, Serie A, XXI, 1966.
- [8] L. Henkin, *An algebraic characterization of quantifiers*, Fundam. Math., **37** (1950) 63-74.
- [9] A. Iorgulescu, *New generalizations of BCI, BCK and Hilbert algebras - Part I*, J. Mult.-Valued Log. Soft Comput., **27** No. 4 (2016) 353-406.
- [10] A. Iorgulescu, *New generalizations of BCI, BCK and Hilbert algebras - Part II*, J. Mult.-Valued Log. Soft Comput., **27** No. 4 (2016) 407-456.
- [11] K. Iséki, *An algebra related with a propositional calculus*, Proc. Jpn. Acad., **42** (1966) 26-29.
- [12] K. Iséki and S. Tanaka, *An introduction to the theory of BCK-algebras*, Math. Japon., **23** (1978) 1-26.
- [13] H. S. Kim and Y. H. Kim, *On BE-algebras*, Scient. Math. Japon., **66** No. 1 (2007) 113-116.
- [14] C. A. Meredith, *Formal Logics*, 2nd ed., Oxford, 1962.
- [15] A. Monteiro, *Lectures on Hilbert and Tarski Algebras*, Insitituto de Matemática, Universidad Nacional del Sur, Bahía Blanca, Argentina, 1960.
- [16] A. Rezaei, A. Borumand Saeid and R. A. Borzooei, *Relation between Hilbert algebras and BE-algebras*, Appl. Math., **8** (2013) 573-584.
- [17] A. Rezaei, R. K. Bandaru, A. Borumand Saeid and Y. B. Jun, *Prominent GE-filters and GE-morphisms in GE-algebras*, Afr. Mat., **32** (2021) 1121-1136.
- [18] S. Z. Song, R. K. Bandaru and Y. B. Jun, *Imploring GE-filters of GE-algebras*, J. Math., 2021, 1-7.

- [19] S. Z. Song, R. K. Bandaru, D. A. Romano and Y. B. Jun, *Interior GE-filters of GE-algebras*, Diss. Math. Gen. Alg. App., **42** (2022) 217-235.
- [20] A. Walendziak, *On commutative BE-algebras*, Scient. Math. Japon., **69** No. 2 (2009) 281-284.
- [21] A. Walendziak, *On implicative and positive implicative GE algebras*, Bull. Sect. Logic, 1-19.
- [22] A. Walendziak, *On pre-Hilbert and positive implicative pre-Hilbert algebras*, Submitted.

Andrzej Walendziak

Institute of Mathematics,

Faculty of Exact and Natural Sciences,

University of Siedlce, PL-08110 Siedlce, Poland.

walent@interia.pl