



Research Paper

ϕ - (k, n) -ABSORBING (PRIMARY) HYPERIDEALS IN A KRASNER
 (m, n) -HYPERRING

MAHDI ANBARLOEI*

ABSTRACT. Various expansions of prime hyperideals have been studied in a Krasner (m, n) -hyperring R . For instance, a proper hyperideal Q of R is called weakly (k, n) -absorbing (primary) provided that for $r_1^{kn-k+1} \in R$, $g(r_1^{kn-k+1}) \in Q - \{0\}$ implies that there are $(k-1)n-k+2$ of the r_i 's whose g -product is in Q ($g(r_1^{(k-1)n-k+2}) \in Q$ or a g -product of $(k-1)n-k+2$ of r_i 's ,except $g(r_1^{(k-1)n-k+2})$, is in $\mathfrak{r}^{(m,n)}(Q)$). In this paper, we aim to extend the notions to the concepts of ϕ - (k, n) -absorbing and ϕ - (k, n) -absorbing primary hyperideals. Assume that ϕ is a function from $\mathcal{HI}(R)$ to $\mathcal{HI}(R) \cup \{\varphi\}$ such that $\mathcal{HI}(R)$ is the set of hyperideals of R and k is a positive integer. We call a proper hyperideal Q of R a ϕ - (k, n) -absorbing (primary) hyperideal if for $r_1^{kn-k+1} \in R$, $g(r_1^{kn-k+1}) \in Q - \phi(Q)$ implies that there are $(k-1)n-k+2$ of the r_i 's whose g -product is in Q ($g(r_1^{(k-1)n-k+2}) \in Q$ or a g -product of $(k-1)n-k+2$ of r_i 's ,except $g(r_1^{(k-1)n-k+2})$, is in $\mathfrak{r}^{(m,n)}(Q)$). Several properties and characterizations of them are presented.

DOI: 10.22034/as.2024.20052.1638

MSC(2010): Primary: 20N20, 16Y99, 20N15, 06E20.

Keywords: ϕ - (k, n) -absorbing hyperideal, ϕ - (k, n) -absorbing primary hyperideal.

Received: 02 May 2023, Accepted: 05 May 2024.

*Corresponding author

1. INTRODUCTION

Extensions of prime and primary ideals to the context of ϕ -prime and ϕ -primary ideals were studied in [7, 12]. Afterwards, Khaksari in [20] and Badawi et al. in [9] introduced ϕ -2-prime and ϕ -2-primary ideals, respectively. Let R be a commutative ring. Suppose that ϕ is a function from $\mathcal{I}(R)$ to $\mathcal{I}(R) \cup \{\varphi\}$ where $\mathcal{I}(R)$ is the set of ideals of R . A proper ideal I of R is said to be a ϕ -2-absorbing ideal if whenever $x, y, z \in R$, with $xyz \in I - \phi(I)$ implies that $xy \in I$ or $xz \in I$ or $yz \in I$. Also, A proper ideal I of R is called a ϕ -2-absorbing primary ideal if for every $x, y, z \in R$, $xyz \in I - \phi(I)$ implies that $xy \in I$ or $xz \in \mathfrak{r}(I)$ or $yz \in \mathfrak{r}(I)$.

Hyperstructures are algebraic structures equipped with at least one multi-valued operation, called a hyperoperation. A hyperoperation on a non-empty set is a mapping from the nonempty power set. Hundreds of papers and several books have been written on this topic (for more details see [2, 10, 11, 13, 17, 21, 26, 30, 32, 33, 34]). An n -ary extension of algebraic structures is the most natural method for deeper understanding of their fundamental properties. Mirvakili and Davvaz in [28] introduced (m, n) -hyperrings and gave several results in this respect. They defined and described a generalization of the notion of a hypergroup and a generalization of an n -ary group, which is called n -ary hypergroup [14]. Some review of the n -ary structures can be found in in [22, 23, 24, 25, 31]. One important class of hyperrings, where the addition is a hyperoperation, while the multiplication is an ordinary binary operation, is Krasner hyperring. An extension of the Krasner hyperrings, which is a subclass of (m, n) -hyperrings, was presented by Mirvakili and Davvaz [27], which is called Krasner (m, n) -hyperring. Some important hyperideals namely Jacobson radical, nilradical, n -ary prime and primary hyperideals and n -ary multiplicative subsets of Krasner (m, n) -hyperrings were defined by Ameri and Norouzi in [1]. Afterward, the concept of (k, n) -absorbing (primary) hyperideals was studied by Hila et al. [18]. Norouzi et al. gave a new definition for normal hyperideals in Krasner (m, n) -hyperrings, with respect to that one given in [27] and they showed that these hyperideals correspond to strongly regular relations [29]. Direct limit of a direct system was defined and analysed by Asadi and Ameri in the category of Krasner (m, n) -hyperrings [8]. The notion of δ -primary hyperideals in Krasner (m, n) -hyperrings, which unifies the prime and primary hyperideals under one frame, was presented in [4]. Recently, Davvaz et al. introduced new expansion classes, namely weakly (k, n) -absorbing (primary) hyperideals in a Krasner (m, n) -hyperring [16].

In this paper, we introduce and study the notions of ϕ - (k, n) -absorbing and ϕ - (k, n) -absorbing primary hyperideals in a commutative Krasner (m, n) -hyperring. A number of main results are given to explain the general framework of these structures. Among many results in this paper, it is shown (Theorem 3.6) that if Q is a ϕ - (k, n) -absorbing hyperideal of R , then Q is a ϕ - (s, n) -absorbing hyperideal for all $s \geq k$. Although every ϕ - (k, n) -absorbing of a Krasner

(m, n) -hyperring is ϕ - (k, n) -absorbing primary, Example 4.3 shows that the converse may not be always true. It is shown (Theorem 4.13) that Q is a ϕ - (k, n) -absorbing primary hyperideal of R if and only if $Q/\phi(Q)$ is a weakly (k, n) -absorbing primary hyperideal of $R/\phi(Q)$. In Theorem 4.16, we show that if Q is a ϕ - (k, n) -absorbing primary hyperideal of R but is not a (k, n) -absorbing primary, then $g(Q^{k(n-1)+1}) \subseteq \phi(Q)$. As a result of the theorem we conclude that if Q is a ϕ - (k, n) -absorbing primary hyperideal of R that is not a (k, n) -absorbing primary hyperideal of R , then $\mathbf{r}^{(m,n)}(Q) = \mathbf{r}^{(m,n)}(\phi(Q))$.

2. KRASNER (m, n) -HYPERRINGS

In this section, we summarize the preliminary definitions which are related to Krasner (m, n) -hyperrings.

Let A be a non-empty set and $P^*(A)$ the set of all the non-empty subsets of A . An n -ary hyperoperation on A is a map $f : A^n \rightarrow P^*(A)$ and the couple (A, f) is called an n -ary hypergroupoid. The notation a_i^j will denote the sequence a_i, a_{i+1}, \dots, a_j for $j \geq i$ and it is the empty symbol for $j < i$. If G_1, \dots, G_n are non-empty subsets of A , then we define $f(G_1^n) = f(G_1, \dots, G_n) = \bigcup \{f(a_1^n) \mid a_i \in G_i, 1 \leq i \leq n\}$. If $b_{i+1} = \dots = b_j = b$, we write $f(a_1^i, b_{i+1}^j, c_{j+1}^n) = f(a_1^i, b^{(j-i)}, c_{j+1}^n)$. If f is an n -ary hyperoperation, then t -ary hyperoperation $f_{(l)}$ is given by

$$f_{(l)}(a_1^{l(n-1)+1}) = f\left(f(\dots, f(f(a_1^n), a_{n+1}^{2n-1}), \dots), a_{(l-1)(n-1)+1}^{l(n-1)+1}\right),$$

where $t = l(n - 1) + 1$.

Definition 2.1. [27] (R, f, g) , or simply R , is defined as a Krasner (m, n) -hyperring if the following statements hold:

- (1) (R, f) is a canonical m -ary hypergroup;
- (2) (R, g) is a n -ary semigroup;
- (3) The n -ary operation g is distributive with respect to the m -ary hyperoperation f , i.e., for every $a_1^{i-1}, a_{i+1}^n, x_1^m \in R$, and $1 \leq i \leq n$,

$$g\left(a_1^{i-1}, f(x_1^m), a_{i+1}^n\right) = f\left(g(a_1^{i-1}, x_1, a_{i+1}^n), \dots, g(a_1^{i-1}, x_m, a_{i+1}^n)\right);$$

- (4) 0 is a zero element of the n -ary operation g , i.e., for each $a_1^n \in R$, $g(a_1^{i-1}, 0, a_{i+1}^n) = 0$.

Throughout this paper, R denotes a commutative Krasner (m, n) -hyperring with the scalar identity 1 .

A non-empty subset T of R is called a subhyperring of R if (T, f, g) is a Krasner (m, n) -hyperring. The non-empty subset I of R is a hyperideal if (I, f) is an m -ary subhypergroup of (R, f) and $g(x_1^{i-1}, I, x_{i+1}^n) \subseteq I$, for each $x_1^n \in R$ and $1 \leq i \leq n$.

Definition 2.2. [1] Let I be a proper hyperideal of R . I refers to a prime hyperideal if for hyperideals I_1^n of R , $g(I_1^n) \subseteq P$ implies $I_i \subseteq I$ for some $1 \leq i \leq n$.

Lemma 4.5 in [1] shows that the proper hyperideal I of R is prime if for all $a_1^n \in R$, $g(a_1^n) \in I$ implies $a_i \in I$ for some $1 \leq i \leq n$.

Definition 2.3. [1] The radical of a proper hyperideal I of R , denoted by $\mathfrak{r}^{(m,n)}(I)$ is the intersection of all prime hyperideals of R containing I . If the set of all prime hyperideals which contain I is empty, then $\mathfrak{r}^{(m,n)}(I) = R$.

It was shown (Theorem 4.23 in [1]) that if $a \in \mathfrak{r}^{(m,n)}(I)$ then there exists $s \in \mathbb{N}$ with $g(a^{(s)}, 1_R^{(n-s)}) \in I$ for $s \leq n$, or $g_{(l)}(a^{(s)}) \in I$ for $s = l(n-1) + 1$.

Definition 2.4. [1] A proper hyperideal I of R is primary if $g(a_1^n) \in I$ for $a_1^n \in R$ implies $a_i \in I$ or $g(a_1^{i-1}, 1_R, a_{i+1}^n) \in \mathfrak{r}^{(m,n)}(I)$ for some $1 \leq i \leq n$.

Theorem 4.28 in [1] shows that the radical of a primary hyperideal of R is prime.

Definition 2.5. [18] Let I be a proper hyperideal of R . I refers to an

- (1) (k, n) -absorbing hyperideal if for $r_1^{kn-k+1} \in R$, $g(r_1^{kn-k+1}) \in I$ implies that there exist $(k-1)n - k + 2$ of the r_i 's whose g -product is in I . In this case, if $k = 1$, then I is an n -ary prime hyperideal of R . If $n = 2$ and $k = 1$, then I is a classic prime hyperideal of R .
- (2) (k, n) -absorbing primary hyperideal if for $r_1^{kn-k+1} \in R$, $g(r_1^{kn-k+1}) \in I$ implies that $g(r_1^{(k-1)n-k+2}) \in I$ or a g -product of $(k-1)n - k + 2$ of the r_i 's, except $g(r_1^{(k-1)n-k+2})$, is in $\mathfrak{r}^{(m,n)}(I)$.

3. ϕ -(k, n)-ABSORBING HYPERIDEALS

In his paper [16], Davvaz et al. introduced a generalization of the n -ary prime hyperideals in a Krasner (k, n) -hyperring, which they defined as weakly (k, n) -absorbing hyperideals. In this section, we generalize this notion to the context of ϕ -(k, n)-absorbing hyperideals.

Definition 3.1. Assume that $\mathcal{HI}(R)$ is the set of hyperideals of R and $\phi : \mathcal{HI}(R) \longrightarrow \mathcal{HI}(R) \cup \{\varphi\}$ is a function. Let k be a positive integer. A proper hyperideal Q of R is said to be ϕ -(k, n)-absorbing provided that for $r_1^{kn-k+1} \in R$, $g(r_1^{kn-k+1}) \in Q - \phi(Q)$ implies that there are $(k-1)n - k + 2$ of the r_i 's whose g -product is in Q .

Example 3.2. Consider the Krasner $(2, 2)$ -hyperring $R = \{0, 1, x\}$ with the hyperaddition and multiplication defined by

+	0	1	x
0	0	1	x
1	1	R	1
x	x	1	$\{0, x\}$

\cdot	0	1	x
0	0	0	0
1	0	1	x
x	0	x	0

Assume that ϕ is a function from $\mathcal{HI}(R)$ to $\mathcal{HI}(R) \cup \{\varphi\}$ defined $\phi(I) = g(I^{(2)})$ for $I \in \mathcal{HI}(R)$. Then the hyperideal $Q = \{0, x\}$ is a ϕ -(2, 2)-absorbing hyperideal of R .

Example 3.3. Let $t > 4$. Consider Krasner $(4, 3)$ -hyperring $(\mathbb{Z}_{5^{5t}}, +, \cdot)$ where $+$ and \cdot are usual addition and multiplication. Defined $\phi(I) = I^5$ for $I \in \mathcal{HI}(\mathbb{Z}_{5^{5t}})$. Then $I = \langle 5^t \rangle$ is not a $(2, 3)$ -absorbing hyperideal of $\mathbb{Z}_{5^{5t}}$ since $5.5.5.5.5^{t-4} \in I - \phi(I)$ but $5.5.5, 5.5.5^{t-4} \notin I$.

Let $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. Clearly, every (k, n) -absorbing hyperideal in a Krasner (m, n) -hyperring is a ϕ -(k, n)-absorbing hyperideal. But, the following example shows that the converse does not necessarily hold.

Example 3.4. Assume that R is the Krasner $(2, 4)$ -hyperring given in Example 4.7 in [1]. In [16], it was shown that $\langle 0 \rangle$ is not a $(1, 4)$ -absorbing hyperideal of R . Now, defined $\phi(I) = g(I^{(4)})$ for $I \in \mathcal{HI}(R)$. In this hyperring, $\langle 0 \rangle$ is a ϕ -(1, 4)-absorbing hyperideal of R .

Theorem 3.5. Let $\phi_1, \phi_2 : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be two functions such that for all $I \in \mathcal{HI}(R)$, $\phi_1(I) \subseteq \phi_2(I)$. If Q is a ϕ_1 -(k, n)-absorbing hyperideal of R , then Q is a ϕ_2 -(k, n)-absorbing hyperideal.

Proof. Suppose that $g(r_1^{kn-k+1}) \in Q - \phi_2(Q)$ for $r_1^{kn-k+1} \in R$. From $\phi_1(Q) \subseteq \phi_2(Q)$, it follows that $g(r_1^{kn-k+1}) \in Q - \phi_1(Q)$. Since Q is a ϕ_1 -(k, n)-absorbing hyperideal of R , we conclude that there are $(k - 1)n - k + 2$ of the r_i 's whose g -product is in Q , as needed. \square

Theorem 3.6. Let $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. If Q is a ϕ -(k, n)-absorbing hyperideal of R , then Q is a ϕ -(s, n)-absorbing hyperideal for all $s \geq k$.

Proof. Let us use the induction on k that if Q is ϕ -(k, n)-absorbing hyperideal of R , then Q is ϕ -($k+1, n$)-absorbing. Assume that Q is ϕ -(2, n)-absorbing and $g(r_1^{2n-2}, g(r_{2n-1}^{3n-2})) \in Q - \phi(Q)$ for some $r_1^{3n-2} \in R$. Since Q is ϕ -(2, n)-absorbing, then there are n of the r_i 's except $g(r_{2n-1}^{3n-2})$ whose g -product is in Q and so there are $2n - 1$ of the r_i 's whose g -product is in Q . This shows that Q is ϕ -(3, n)-absorbing. Assume that Q is ϕ -(k, n)-absorbing and $g(g(r_1^{2n-2}), r_{2n-1}^{(k+1)n-(k+1)+1}) \in Q - \phi(Q)$ for some $r_1^{(k+1)n-(k+1)+1} \in R$. Since Q is ϕ -(k, n)-absorbing, we conclude that $g(g(r_1^{2(n-1)}), r_{2n-1}, \dots, \widehat{r_i}, \dots, r_{(k+1)n-(k+1)+1}) \in Q$ for some $2(n-1) \leq i \leq (k+1)n-(k+1)+1$ or $g(r_{2n-1}^{(k+1)n-(k+1)+1}) \in Q$. The former case shows that Q is ϕ -($k + 1, n$)-absorbing. In the

latter case, we obtain $g(r_1^{n-1}, r_{2n-1}^{(k+1)n-(k+1)+1}) \in Q$ since $g(r_1^{2(n-1)}) \in Q$. Thus Q is ϕ - $(k+1, n)$ -absorbing. \square

Recall from [15] that if (R_1, f_1, g_1) and (R_2, f_2, g_2) are two Krasner (m, n) -hyperrings such that 1_{R_1} and 1_{R_2} are scalar identities of R_1 and R_2 , respectively, then $(R_1 \times R_2, f_1 \times f_2, g_1 \times g_2)$ is a Krasner (m, n) -hyperring where

$$\begin{aligned} f &= f_1 \times f_2((a_1, b_1), \dots, (a_m, b_m)) = \{(a, b) \mid a \in f_1(a_1^m), b \in f_2(b_1^m)\}, \\ g &= g_1 \times g_2((x_1, y_1), \dots, (x_n, y_n)) = (g_1(x_1^n), g_2(y_1^n)), \end{aligned}$$

for all $a_1^m, x_1^n \in R_1$ and $b_1^m, y_1^n \in R_2$.

Theorem 3.7. *Let (R_i, f_i, g_i) be a commutative Krasner (m, n) -hyperring for each $1 \leq i \leq kn - k + 1$ and $\phi_i : \mathcal{HI}(R_i) \rightarrow \mathcal{HI}(R_i) \cup \{\varphi\}$ be a function. Let Q_i be a hyperideal of R_i for each $1 \leq i \leq kn - k + 1$ and $\phi = \phi_1 \times \dots \times \phi_{kn-k+1}$. If $Q = Q_1 \times \dots \times Q_{kn-k+1}$ is a ϕ - $(k+1, n)$ -absorbing hyperideal of $R = R_1 \times \dots \times R_{kn-k+1}$, then Q_i is a ϕ_i - (k, n) -absorbing hyperideal of R_i and $Q_i \neq R_i$ for all $1 \leq i \leq kn - k + 1$.*

Proof. Let $r_1^{kn-k+1} \in R_i$ such that $g(r_1^{kn-k+1}) \in Q_i - \phi_i(Q_i)$. Suppose by contradiction that Q_i is not a ϕ_i - (k, n) -absorbing hyperideal of R_i . Define

$$\begin{aligned} a_1 &= (1_{R_1}, \dots, 1_{R_{i-1}}, r_1, 1_{R_{i+1}}, \dots, 1_{R_{kn-k+1}}), \\ a_2 &= (1_{R_1}, \dots, 1_{R_{i-1}}, r_2, 1_{R_{i+1}}, \dots, 1_{R_{kn-k+1}}), \\ &\vdots \\ a_{kn-k+1} &= (1_{R_1}, \dots, 1_{R_{i-1}}, r_{kn-k+1}, 1_{R_{i+1}}, \dots, 1_{R_{kn-k+1}}), \\ a_{kn-k} &= (1_{R_1}, \dots, 1_{R_{i-1}}, 1_{R_i}, 1_{R_{i+1}}, \dots, 1_{R_{kn-k+1}}), \\ a_{(k+1)n-(k+1)+1} &= (0, \dots, 0, 1_{R_i}, 0, \dots, 0). \end{aligned}$$

Hence $g(a_1^{(k+1)n-(k+1)+1}) \in Q - \phi(Q)$ but $g(a_1^{kn-k+1}) \notin Q$. Since Q is a ϕ - $(k+1, n)$ -absorbing hyperideal of R , we conclude that one of g -products of $kn - k + 1$ of a_i 's except $g(a_1^{(k+1)n-(k+1)+1})$ is in Q . This implies that there exist $(k-1)n - k + 2$ of r_i 's whose g -product is in Q_i which is a contradiction. Consequently, Q_i is a ϕ_i - (k, n) -absorbing hyperideal of R_i . \square

Assume that (R_1, f_1, g_1) and (R_2, f_2, g_2) are two Krasner (m, n) -hyperrings. Recall from [27] that a mapping $h : R_1 \rightarrow R_2$ is called a homomorphism if for all $a_1^m \in R_1$ and $b_1^n \in R_1$ we have (1) $h(f_1(a_1, \dots, a_m)) = f_2(h(a_1), \dots, h(a_m))$, (2) $h(g_1(b_1, \dots, b_n)) = g_2(h(b_1), \dots, h(b_n))$. Moreover, recall from [19] that a function $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ is called a reduction function of $\mathcal{HI}(R)$ if $\phi(P) \subseteq P$ and $P \subseteq Q$ implies that $\phi(P) \subseteq \phi(Q)$ for all $P, Q \in \mathcal{HI}(R)$.

Now, assume that R_1 and R_2 are two Krasner (m, n) -hyperring such that $h : R_1 \rightarrow R_2$ is a homomorphism. Suppose that ϕ_1 and ϕ_2 are two reduction functions of $\mathcal{HI}(R_1)$ and $\mathcal{HI}(R_2)$, respectively. If $\phi_1(h^{-1}(I_2)) = h^{-1}(\phi_2(I_2))$ for all $I_2 \in \mathcal{HI}(R_2)$, then we say h is a ϕ_1 - ϕ_2 -homomorphism. Let h be a ϕ_1 - ϕ_2 -epimorphism from R_1 to R_2 and let I_1 be a hyperideal of R_1 with $\text{Ker}(h) \subseteq I_1$. It is easy to see that $\phi_2(h(I_1)) = h(\phi_1(I_1))$.

Example 3.8. Let R_1 and R_2 be two Krasner (m, n) -hyperrings and ϕ_1 and ϕ_2 be two empty reduction functions of $\mathcal{HI}(R_1)$ and $\mathcal{HI}(R_2)$, respectively. Then every homomorphism h from R_1 to R_2 is a ϕ_1 - ϕ_2 -homomorphism.

Theorem 3.9. Let $h : R_1 \rightarrow R_2$ be a ϕ_1 - ϕ_2 -homomorphism, where ϕ_1 and ϕ_2 are two reduction functions of $\mathcal{HI}(R_1)$ and $\mathcal{HI}(R_2)$, respectively. Then

- (1) If Q_2 is a ϕ_2 - (k, n) -absorbing hyperideal of R_2 , then $h^{-1}(Q_2)$ is a ϕ_1 - (k, n) -absorbing of R_1 .
- (2) If h is surjective and Q_1 is a ϕ_1 - (k, n) -absorbing hyperideal of R_1 with $\text{Ker}(h) \subseteq Q_1$, then $h(Q_1)$ is a ϕ_2 - (k, n) -absorbing hyperideal of R_2 .

Proof. (1) Let Q_2 be a ϕ_2 - (k, n) -absorbing hyperideal of R_2 and $g(r_1^{kn-k+1}) \in h^{-1}(Q_2) - \phi_1(h^{-1}(Q_2))$ for some $r_1^{kn-k+1} \in R_1$. Then we get $h(g(r_1^{kn-k+1})) = g(h(r_1), \dots, h(r_{kn-k+1})) \in Q_2 - \phi_2(Q_2)$. Since Q_2 is a ϕ_2 - (k, n) -absorbing hyperideal of R_2 , we conclude that the image of h of $(k-1)n-k+2$ of r_i 's whose g -product is in Q_2 . Then there exist $(k-1)n-k+2$ of r_i 's whose g -product is in $h^{-1}(Q_2)$. Thus $h^{-1}(Q_2)$ is a ϕ_1 - (k, n) -absorbing of R_1 .

(2) Suppose that Q_1 is a ϕ_1 - (k, n) -absorbing hyperideal of R_1 with $\text{Ker}(h) \subseteq Q_1$ and h is surjective. Let $g(s_1^{kn-k+1}) \in h(Q_1) - \phi_2(h(Q_1))$ for some $s_1^{kn-k+1} \in R_2$. Then there exists $r_i \in R_1$ for every $1 \leq i \leq kn-k+1$ such that $h(r_i) = s_i$. Hence we get $h(g(r_1^{kn-k+1})) = g(h(r_1), \dots, h(r_{kn-k+1})) = g(s_1^{kn-k+1}) \in h(Q_1)$. Since $\text{Ker}(h) \subseteq Q_1$ and h is a ϕ_1 - ϕ_2 -epimorphism, we have $g(r_1^{kn-k+1}) \in Q_1 - \phi_1(Q_1)$. Since Q_1 is a ϕ_1 - (k, n) -absorbing hyperideal of R_1 , there are $(k-1)n-k+2$ of r_i 's whose g -product is in Q_1 . Now, since h is a homomorphism, we are done. \square

Let P be a hyperideal of R . Then the set $R/P = \{f(a_1^{i-1}, P, a_{i+1}^m) \mid a_1^{i-1}, a_{i+1}^m \in R\}$ with m -ary hyperoperation f and n -operation g is the quotient Krasner (m, n) -hyperring of R by P . Theorem 3.2 in [1] shows that the projection map π from R to R/P , defined by $\pi(r) = f(r, P, 0^{(m-2)})$, is homomorphism. Let P be a hyperideal of R and ϕ be a reduction function of $\mathcal{HI}(R)$. Then the function ϕ_q from $\mathcal{HI}(R/P)$ to $\mathcal{HI}(R/P) \cup \{\varphi\}$ defined by $\phi_q(I/P) = \phi(I)/P$ is a reduction function. Now, we have the following theorem as a result of Theorem 3.9 that is easily verified, and hence we omit the proof.

Theorem 3.10. *Let Q and P be two hyperideals of R and ϕ be a reduction function of $\mathcal{HI}(R)$ such that $P \subseteq \phi(Q) \subseteq Q$. If Q is a ϕ - (k, n) -absorbing hyperideal of R , then Q/P is a ϕ_Q - (k, n) -absorbing hyperideal of R/P .*

4. ϕ - (k, n) -ABSORBING PRIMARY HYPERIDEALS

Definition 4.1. Suppose that $\mathcal{HI}(R)$ is the set of hyperideals of R and $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ is a function. Let k be a positive integer. A proper hyperideal Q of R is called ϕ - (k, n) -absorbing primary if $g(r_1^{kn-k+1}) \in Q - \phi(Q)$ for $r_1^{kn-k+1} \in R$ implies that $g(r_1^{(k-1)n-k+2}) \in Q$ or a g -product of $(k-1)n - k + 2$ of r_i 's, except $g(r_1^{(k-1)n-k+2})$, is in $\mathbf{r}^{(m,n)}(Q)$.

Example 4.2. Every ϕ - (k, n) -absorbing of a Krasner (m, n) -hyperring is ϕ - (k, n) -absorbing primary.

The converse may not be always true as it is shown in the following example.

Example 4.3. Consider the Krasner $(2, 2)$ -hyperring $R = [0, 1]$ with the 2-ary hyperoperation defined by

$$a \oplus b = \begin{cases} \{\max\{a, b\}\}, & \text{if } a \neq b, \\ [0, a], & \text{if } a = b, \end{cases}$$

and multiplication is the usual multiplication on real numbers. Suppose that ϕ is a function from $\mathcal{HI}(R)$ to $\mathcal{HI}(R) \cup \{\varphi\}$ defined $\phi(I) = \bigcap_{i=1}^{\infty} g(I^{(i)})$ for $I \in \mathcal{HI}(R)$. Then the hyperideal $Q = [0, 0.5]$ is a ϕ - $(2, 2)$ -absorbing primary hyperideal of R but it is not ϕ - $(2, 2)$ -absorbing.

The next theorem provides us how to determine ϕ - (k, n) -absorbing primary hyperideal to be (k, n) -absorbing primary.

Theorem 4.4. *Assume that Q is a hyperideal of R and $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ is a reduction function such that $\phi(Q)$ is a (k, n) -absorbing primary hyperideal of R . If Q is a ϕ - (k, n) -absorbing primary hyperideal of R , then Q is a (k, n) -absorbing primary hyperideal of R .*

Proof. Let $r_1^{kn-k+1} \in R$ such that $g(r_1^{kn-k+1}) \in Q$ and $g(r_1^{(k-1)n-k+2}) \notin Q$. Assume that $g(r_1^{kn-k+1}) \in \phi(Q)$. Since $\phi(Q)$ is a (k, n) -absorbing primary hyperideal of R and $g(r_1^{(k-1)n-k+2}) \notin \phi(Q)$, we conclude that a g -product of $(k-1)n - k + 2$ of the r_i 's, except $g(r_1^{(k-1)n-k+2})$ is in $\mathbf{r}^{(m,n)}(\phi(Q)) \subseteq \mathbf{r}^{(m,n)}(Q)$, as needed. Suppose that $g(r_1^{kn-k+1}) \notin \phi(Q)$. Since Q is a ϕ - (k, n) -absorbing primary hyperideal of R , we are done. \square

In the following, the relationship between a ϕ - (k, n) -absorbing primary hyperideal of R and its radical is considered.

Theorem 4.5. *Let Q be a hyperideal of R and $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function such that $\mathbf{r}^{(m,n)}(\phi(Q)) = \phi(\mathbf{r}^{(m,n)}(Q))$. If Q is a ϕ - (k, n) -absorbing primary hyperideal of R , then $\mathbf{r}^{(m,n)}(Q)$ is a ϕ - (k, n) -absorbing hyperideal of R .*

Proof. Let $r_1^{kn-k+1} \in R$ such that $g(r_1^{kn-k+1}) \in \mathbf{r}^{(m,n)}(Q) - \phi(\mathbf{r}^{(m,n)}(Q))$. Assume that all products of $(k-1)n-k+2$ of the r_i 's except $g(r_1^{(k-1)n-k+2})$ are not in $\mathbf{r}^{(m,n)}(Q)$. Since $g(r_1^{kn-k+1}) \in \mathbf{r}^{(m,n)}(Q)$, then there exists $s \in \mathbb{Z}^+$ with $g(g(r_1^{kn-k+1})^{(s)}, 1^{(n-s)}) \in Q$, for $s \leq n$ or $g_{(l)}(g(r_1^{kn-k+1})^{(s)}) \in Q$, for $s > n$, $s = l(n-1) + 1$. In the former case, we get $g(g(r_1)^{(s)}, g(r_2)^{(s)}, \dots, g(r_{kn-k+1})^{(s)}, 1^{(n-s)}) \in Q$. If $g(g(r_1)^{(s)}, g(r_2)^{(s)}, \dots, g(r_{kn-k+1})^{(s)}, 1^{(n-s)}) \in \phi(Q)$, we obtain $g(r_1^{kn-k+1}) \in \mathbf{r}^{(m,n)}(\phi(Q)) = \phi(\mathbf{r}^{(m,n)}(Q))$, a contradiction. Since Q is a ϕ - (k, n) -absorbing primary hyperideal of R , then $g(g(r_1)^{(s)}, g(r_2)^{(s)}, \dots, g(r_{(k-1)n-k+2})^{(s)}, 1^{(n-s)}) = g(g(r_1^{(k-1)n-k+2})^{(s)}, 1^{(n-s)}) \in Q$ which means $g(r_1^{(k-1)n-k+2}) \in \mathbf{r}^{(m,n)}(Q)$. For the other case, we have a similar argument. Consequently, $\mathbf{r}^{(m,n)}(Q)$ is a ϕ - (k, n) -absorbing hyperideal of R . \square

Example 4.6. Assume that $H = \mathbb{Z}_3[X, Y, Z]$ and $Q = \langle X^3Y^3Z^3 \rangle$. Then $R = H/Q$ is a Krasner (m, n) -hyperring with ordinary addition and ordinary multiplication. Defined $\phi(I/Q) = 0R$ for $I/Q \in \mathcal{HI}(R)$. In the hyperring, Q/Q is a ϕ - $(1, 3)$ -absorbing primary hyperideal of R and $\mathbf{r}^{(m,n)}(\phi(Q/Q)) \neq \phi(\mathbf{r}^{(m,n)}(Q/Q))$. Note that $\mathbf{r}^{(m,n)}(Q/Q)$ is not a ϕ - $(1, 3)$ -absorbing hyperideal of R because $2XYZ + Q = (2X + Q)(Y + Q)(Z + Q) \in \mathbf{r}^{(m,n)}(Q/Q) - \phi(\mathbf{r}^{(m,n)}(Q/Q))$ but none of the elements $(2X + Q)$, $(Y + Q)$ and $(Z + Q)$ are not in $\mathbf{r}^{(m,n)}(Q/Q)$.

Theorem 4.7. *Assume that $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ is a function. If Q is a ϕ - (k, n) -absorbing primary hyperideal of R , then Q is a ϕ - (s, n) -absorbing primary hyperideal for all $s \geq k$.*

Proof. Let Q be a ϕ - (k, n) -absorbing primary hyperideal of R . Suppose that $g(g(r_1^{n+2}), r_{n+3}^{(k+1)n-(k+1)+1}) \in Q - \phi(Q)$ for some $r_1^{(k+1)n-(k+1)+1} \in R$. Put $g(r_1^{n+2}) = a_1$. Then we conclude that $g(a_1, \dots, r_{(k+1)n-(k+1)+1}) \in Q$ or a g -product of $kn-k+1$ of the r_i 's, except $g(a_1, \dots, r_{(k+1)n-(k+1)+1})$ is in $\mathbf{r}^{(m,n)}(Q)$ as Q is a ϕ - (k, n) -absorbing primary hyperideal of R . Since $\mathbf{r}^{(m,n)}(Q)$ is a hyperideal of R and $r_1^{n+2} \in R$, we conclude that $g(r_1, r_{n+3}, \dots, r_{(k+1)n-(k+1)+1}) \in \mathbf{r}^{(m,n)}(Q)$ or \dots or $g(r_{n+2}, r_{n+3}, \dots, r_{(k+1)n-(k+1)+1}) \in \mathbf{r}^{(m,n)}(Q)$ and so Q is $(k+1, n)$ -absorbing primary. \square

Theorem 4.8. *Let $\phi_1, \phi_2 : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be two functions such that for all $I \in \mathcal{HI}(R)$, $\phi_1(I) \subseteq \phi_2(I)$. If Q is a ϕ_1 - (k, n) -absorbing primary hyperideal of R , then Q is a ϕ_2 - (k, n) -absorbing primary hyperideal.*

Proof. It is proved in a similar way to Theorem 3.5. \square

Theorem 4.9. *Let $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. If Q is a ϕ - $(1, n)$ -absorbing primary hyperideal of R , then Q is a ϕ - $(2, n)$ -absorbing primary hyperideal.*

Proof. Let Q be a ϕ - $(1, n)$ -absorbing primary hyperideal and $g(r_1^n, \dots, r_{2n-1}) \in Q - \phi(Q)$ for some $r_1^{2n-1} \in R$. Then we get $g(r_1^n) \in Q$ or $g(r_{n+1}^{2n-1}) \in \mathbf{r}^{(m,n)}(Q)$. By definition of hyperideal, we conclude that $g(r_1, r_{n+1}, \dots, r_{2n-1}) \in \mathbf{r}^{(m,n)}(Q)$ or \dots or $g(r_1, r_{n+1}, \dots, r_{2n-1}) \in \mathbf{r}^{(m,n)}(Q)$ since $r_1^n \in R$. Consequently, Q is a ϕ - $(2, n)$ -absorbing primary hyperideal of R . \square

Let Q be a proper hyperideal of R and $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. Q refers to a strongly ϕ - (k, n) -absorbing primary hyperideal of R if $g(Q_1^{kn-k+1}) \subseteq Q - \phi(Q)$ for some hyperideals Q_1^{kn-k+1} of R implies that $g(Q_1^{(k-1)n-k+2}) \subseteq Q$ or a g -product of $(k-1)n-k+2$ of Q_i 's, except $g(Q_1^{(k-1)n-k+2})$, is a subset of $\mathbf{r}^{(m,n)}(Q)$. In the sequel, we assume that all ϕ - (k, n) -absorbing primary hyperideals of R are strongly ϕ - (k, n) -absorbing primary hyperideal. Recall from [16] that a proper hyperideal Q of R is called weakly (k, n) -absorbing primary if $0 \neq g(r_1^{kn-k+1}) \in Q$ for $r_1^{kn-k+1} \in R$ implies that $g(r_1^{(k-1)n-k+2}) \in Q$ or a g -product of $(k-1)n-k+2$ of r_i 's, except $g(r_1^{(k-1)n-k+2})$, is in $\mathbf{r}^{(m,n)}(Q)$.

Theorem 4.10. *Suppose that Q is a proper hyperideal of a commutative Krasner $(m, 2)$ -hyperring R and $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ is a function. Then the followings are equivalent:*

- (1) Q is a ϕ - $(2, 2)$ -absorbing primary hyperideal of R .
- (2) $Q/\phi(Q)$ is a weakly $(2, 2)$ -absorbing primary hyperideal of $R/\phi(Q)$.

Proof. (1) \implies (2) Let Q be ϕ - $(2, 2)$ -absorbing primary and for $a_{11}^{1m}, a_{21}^{2m}, a_{31}^{3m} \in R$,

$$\begin{aligned} \phi(Q) &\neq g(f(a_{11}^{1(i-1)}, \phi(Q), a_{1(i+1)}^{1m}), f(a_{21}^{2(i-1)}, \phi(Q), a_{2(i+1)}^{2m}), f(a_{31}^{3(i-1)}, \phi(Q), a_{3(i+1)}^{3m})) \\ &= f(g(a_{11}^{31}), \dots, g(a_{1(i-1)}^{3(i-1)}), \phi(Q), g(a_{1(i+1)}^{3(i+1)}), \dots, g(a_{1m}^{3m})) \\ &\in Q/\phi(Q). \end{aligned}$$

Then

$$\begin{aligned} &f(g(a_{11}^{31}), \dots, g(a_{1(i-1)}^{3(i-1)}), 0, g(a_{1(i+1)}^{3(i+1)}), \dots, g(a_{1m}^{3m})) \\ &= g(f(a_{11}^{1(i-1)}, 0, a_{1(i+1)}^{1m}), f(a_{21}^{2(i-1)}, 0, a_{2(i+1)}^{2m}), f(a_{31}^{3(i-1)}, 0, a_{3(i+1)}^{3m})) \\ &\in Q - \phi(Q). \end{aligned}$$

Since Q is a ϕ -(2, 2)-absorbing primary hyperideal of R , we get

$$\begin{aligned} &g(f(a_{11}^{1(i-1)}, 0, a_{1(i+1)}^{1m}), f(a_{21}^{2(i-1)}, 0, a_{2(i+1)}^{2m})) \\ &= f(g(a_{11}^{21}), \dots, g(a_{1(i-1)}^{2(i-1)}), 0, g(a_{1(i+1)}^{2(i+1)}), \dots, g(a_{1m}^{2m})) \subseteq Q, \\ &g(f(a_{21}^{2(i-1)}, 0, a_{2(i+1)}^{2m}), f(a_{31}^{3(i-1)}, 0, a_{3(i+1)}^{3m})) \\ &= f(g(a_{21}^{31}), \dots, g(a_{2(i-1)}^{3(i-1)}), 0, g(a_{2(i+1)}^{3(i+1)}), \dots, g(a_{2m}^{3m})) \subseteq \mathbf{r}^{(m,n)}(Q), \end{aligned}$$

or

$$\begin{aligned} &g(f(a_{11}^{1(i-1)}, 0, a_{1(i+1)}^{1m}), f(a_{31}^{3(i-1)}, 0, a_{3(i+1)}^{3m})) \\ &= f(g(a_{11}^{31}), \dots, g(a_{1(i-1)}^{3(i-1)}), 0, g(a_{1(i+1)}^{3(i+1)}), \dots, g(a_{1m}^{3m})) \subseteq \mathbf{r}^{(m,n)}(Q). \end{aligned}$$

It implies that

$$\begin{aligned} &f(g(a_{11}^{21}), \dots, g(a_{1(i-1)}^{2(i-1)}), \phi(Q), g(a_{1(i+1)}^{2(i+1)}), \dots, g(a_{1m}^{2m})) \\ &= g(f(a_{11}^{1(i-1)}, \phi(Q), a_{1(i+1)}^{1m}), f(a_{21}^{2(i-1)}, \phi(Q), a_{2(i+1)}^{2m})) \in Q/\phi(Q), \end{aligned}$$

or

$$\begin{aligned} &f(g(a_{21}^{31}), \dots, g(a_{2(i-1)}^{3(i-1)}), \phi(Q), g(a_{2(i+1)}^{3(i+1)}), \dots, g(a_{2m}^{3m})) \\ &= g(f(a_{21}^{2(i-1)}, \phi(Q), a_{2(i+1)}^{2m}), f(a_{31}^{3(i-1)}, \phi(Q), a_{3(i+1)}^{3m})) \\ &\in \mathbf{r}^{(m,n)}(Q)/\phi(Q) = \mathbf{r}^{(m,n)}(Q/\phi(Q)), \end{aligned}$$

or

$$\begin{aligned} &f(g(a_{11}^{31}), \dots, g(a_{1(i-1)}^{3(i-1)}), \phi(Q), g(a_{1(i+1)}^{3(i+1)}), \dots, g(a_{1m}^{3m})) \\ &= g(f(a_{11}^{1(i-1)}, \phi(Q), a_{1(i+1)}^{1m}), f(a_{31}^{3(i-1)}, \phi(Q), a_{3(i+1)}^{3m})) \\ &\in \mathbf{r}^{(m,n)}(Q)/\phi(Q) = \mathbf{r}^{(m,n)}(Q/\phi(Q)). \end{aligned}$$

(2) \implies (1) Let $g(r_1^3) \in Q - \phi(Q)$ for some $r_1^3 \in R$. Therefore we obtain $f(g(r_1^3), \phi(Q), 0^{(m-2)}) \neq \phi(Q)$. It follows that

$$\phi(Q) \neq g(f(r_1, \phi(Q), 0^{(m-2)}), f(r_2, \phi(Q), 0^{(m-2)}), f(r_3, \phi(Q), 0^{(m-2)})) \in Q/\phi(Q).$$

By the hypothesis, we get

$$g(f(r_1, \phi(Q), 0^{(m-2)}), f(r_2, \phi(Q), 0^{(m-2)})) = f(g(r_1^2), \phi(Q), 0^{(m-2)}) \in Q/\phi(Q).$$

or

$$g(f(r_2, \phi(Q), 0^{(m-2)}), f(r_3, \phi(Q), 0^{(m-2)})) = f(g(r_2^3), \phi(Q), 0^{(m-2)}) \in \mathbf{r}^{(m,n)}(Q)/\phi(Q).$$

or

$$g(f(r_1, \phi(Q), 0^{(m-2)}), f(r_3, \phi(Q), 0^{(m-2)})) = f(g(r_1^3), \phi(Q), 0^{(m-2)}) \in \mathbf{r}^{(m,n)}(Q)/\phi(Q).$$

This shows that $g(r_1^2) \in Q$ or $g(r_2^3) \in \mathbf{r}^{(m,n)}(Q)$ or $g(r_1^3) \in \mathbf{r}^{(m,n)}(Q)$. Consequently, Q is a ϕ -(2, 2)-absorbing primary hyperideal of R . \square

Suppose that I is a weakly (2, 2)-absorbing primary hyperideal of a commutative Krasner $(m, 2)$ -hyperring R . Recall from [16] that (x, y, z) is said to be (2, 2)-zero primary of I for $x, y, z \in R$, if $g(x, y, z) = 0$, $g(x, y) \notin I$, $g(y, z) \notin \mathbf{r}^{(m,n)}(I)$ and $g(x, z) \notin \mathbf{r}^{(m,n)}(I)$. Now, assume that Q is a ϕ -(2, 2)-absorbing primary hyperideal of a commutative Krasner $(m, 2)$ -hyperring R . Then we say (x, y, z) is a ϕ -(2, 2) primary of Q for some $x, y, z \in R$ if $g(x, y, z) \in \phi(Q)$, $g(x, y) \notin Q$, $g(y, z) \notin \mathbf{r}^{(m,n)}(Q)$ and $g(x, z) \notin \mathbf{r}^{(m,n)}(Q)$. It is easy to see that a proper hyperideal Q of R is ϕ -(2, 2)-absorbing primary that is not (2, 2)-absorbing primary if and only if Q has a ϕ -(2, 2) primary (x, y, z) for some $x, y, z \in R$.

Theorem 4.11. *Let R be a commutative Krasner $(m, 2)$ -hyperring and let $\phi : \mathcal{HI}(R) \longrightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. Let Q be a ϕ -(2, 2)-absorbing primary hyperideal of R and $x, y, z \in R$. Then the followings are equivalent:*

- (1) (x, y, z) is a ϕ -(2, 2) primary of Q .
- (2) $(f(x, \phi(Q), 0^{(m-2)}), f(y, \phi(Q), 0^{(m-2)}), f(z, \phi(Q), 0^{(m-2)}))$ is a (2, 2)-zero primary of $Q/\phi(Q)$.

Proof. (1) \implies (2) Let (x, y, z) be a ϕ -(2, 2) primary of Q . This means that $g(x, y, z) \in \phi(Q)$, $g(x, y) \notin Q$, $g(y, z) \notin \mathbf{r}^{(m,n)}(Q)$ and $g(x, z) \notin \mathbf{r}^{(m,n)}(Q)$. This implies that $f(g(x, y), Q, 0^{(m-2)}) \notin Q/\phi(Q)$, $f(g(y, z), \phi(Q), 0^{(m-2)}) \notin \mathbf{r}^{(m,n)}(Q)/\phi(Q)$ and $f(g(x, z), \phi(Q), 0^{(m-2)}) \notin \mathbf{r}^{(m,n)}(Q)/\phi(Q)$. By Theorem 4.10, we conclude that $(f(x, \phi(Q), 0^{(m-2)}), f(y, \phi(Q), 0^{(m-2)}), f(z, \phi(Q), 0^{(m-2)}))$ is a (2, 2)-zero primary of $Q/\phi(Q)$.

(2) \implies (1) Assume that $(f(x, \phi(Q), 0^{(m-2)}), f(y, \phi(Q), 0^{(m-2)}), f(z, \phi(Q), 0^{(m-2)}))$ is a (2, 2)-zero primary of $Q/\phi(Q)$. Thus $g(x, y, z) \in \phi(Q)$ but $f(g(x, y), Q, 0^{(m-2)}) \notin Q/\phi(Q)$, $f(g(y, z), \phi(Q), 0^{(m-2)}) \notin \mathbf{r}^{(m,n)}(Q)/\phi(Q)$ and $f(g(x, z), \phi(Q), 0^{(m-2)}) \notin \mathbf{r}^{(m,n)}(Q)/\phi(Q)$. Hence $g(x, y, z) \in \phi(Q)$, $g(x, y) \notin Q$, $g(y, z) \notin \mathbf{r}^{(m,n)}(Q)$ and $g(x, z) \notin \mathbf{r}^{(m,n)}(Q)$. It implies that (x, y, z) is a ϕ -(2, 2) primary of Q . \square

Theorem 4.12. *Let R be a commutative Krasner $(m, 2)$ -hyperring and let $\phi : \mathcal{HI}(R) \longrightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. Let Q be a ϕ -(2, 2)-absorbing primary hyperideal of R . If (x, y, z) is a ϕ -(2, 2) primary of Q for some $x, y, z \in R$, then*

- (1) $g(x, y, Q), g(y, z, Q), g(x, z, Q) \subseteq \phi(Q)$.

- (2) $g(x, Q^{(2)}), g(y, Q^{(2)}), g(z, Q^{(2)}) \subseteq \phi(Q)$.
- (3) $g(Q^{(3)}) \subseteq \phi(Q)$.

Proof. (1) Let (x, y, z) be a ϕ -(2, 2) primary of a ϕ -(2, 2)-absorbing primary hyperideal Q . By Theorem 4.11, $(f(x, \phi(Q), 0^{(m-2)}), f(y, \phi(Q), 0^{(m-2)}), f(z, \phi(Q), 0^{(m-2)}))$ is a (2, 2)-zero primary of $Q/\phi(Q)$ since (x, y, z) is a ϕ -(2, 2) primary of Q . Thus

$$f(g(x, y, Q), \phi(Q), 0^{(m-2)}) = f(g(y, z, Q), \phi(Q), 0^{(m-2)}) = f(g(x, z, Q), \phi(Q), 0^{(m-2)}) = \phi(Q),$$

by Theorem 4.9 in [16], which implies $g(x, y, Q)$, $g(y, z, Q)$ and $g(x, z, Q)$ are subsets of $\phi(Q)$.

(2) Theorem 4.11 shows that $(f(x, \phi(Q), 0^{(m-2)}), f(y, \phi(Q), 0^{(m-2)}), f(z, \phi(Q), 0^{(m-2)}))$ is a (2, 2)-zero primary of $Q/\phi(Q)$. Moreover, Theorem 4.10 shows that $Q/\phi(Q)$ is a weakly (2, 2)-absorbing primary of $R/\phi(Q)$. Then $f(g(x, Q^{(2)}), \phi(Q), 0^{(m-2)}) = f(g(y, Q^{(2)}), \phi(Q), 0^{(m-2)}) = f(g(z, Q^{(2)}), \phi(Q), 0^{(m-2)}) = \phi(Q)$, by Theorem 4.9 of [16]. Consequently, $g(x, Q^{(2)}), g(y, Q^{(2)}), g(z, Q^{(2)})$ are subsets of $\phi(Q)$.

(3) Again, $(f(x, \phi(Q), 0^{(m-2)}), f(y, \phi(Q), 0^{(m-2)}), f(z, \phi(Q), 0^{(m-2)}))$ is a (2, 2)-zero primary of $Q/\phi(Q)$ and $Q/\phi(Q)$ is a weakly (2, 2)-absorbing primary of $R/\phi(Q)$ by Theorem 4.11 and Theorem 4.10, respectively, then $f(g(Q^{(3)}), \phi(Q), 0^{(m-2)}) = \phi(Q)$ by Theorem 4.10 in [16]. Thus $g(Q^{(3)})$ is a subset of $\phi(Q)$. \square

Theorem 4.13. *Suppose that Q is a proper hyperideal of a commutative Krasner (m, n) -hyperring R and $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ is a function. Then the followings are equivalent:*

- (1) Q is a ϕ -(k, n)-absorbing primary hyperideal of R .
- (2) $Q/\phi(Q)$ is a weakly (k, n)-absorbing primary hyperideal of $R/\phi(Q)$.

Proof. It can be easily proved in a similar manner to the proof of Theorem 4.10. \square

Suppose that Q is a ϕ -(k, n)-absorbing primary hyperideal of R . Then we say $(r_1^{k(n-1)+1})$ is a ϕ -(k, n) primary of Q for some $r_1^{k(n-1)+1} \in R$ if $g(r_1^{k(n-1)+1}) \in \phi(Q)$, $g(r_1^{(k-1)n-k+2}) \notin Q$ and a g -product of $(k-1)n-k+2$ of r_i 's, except $g(r_1^{(k-1)n-k+2})$, is not in $\mathfrak{r}^{(m,n)}(Q)$.

Theorem 4.14. *Let R be a commutative Krasner $(m, 2)$ -hyperring and let $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. Let Q be a ϕ -(k, n)-absorbing primary hyperideal of R and $r_1^{k(n-1)+1} \in R$. Then the followings are equivalent:*

- (1) $(r_1^{k(n-1)+1})$ is a ϕ -(k, n) primary of Q .
- (2) $(f(r_1, \phi(Q), 0^{(m-2)}), \dots, f(r_{k(n-1)+1}, \phi(Q), 0^{(m-2)}))$ is a (k, n)-zero primary of $Q/\phi(Q)$.

Proof. It is seen to be true in a similar manner to Theorem 4.11. \square

Theorem 4.15. *Let R be a commutative Krasner (m, n) -hyperring and let $\phi : \mathcal{HI}(R) \longrightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. Let Q be a ϕ - (k, n) -absorbing primary hyperideal of R . If $(r_1^{k(n-1)+1})$ is a ϕ - (k, n) primary of Q for some $r_1^{k(n-1)+1} \in R$, then $g(r_1, \dots, \widehat{r_{i_1}}, \dots, \widehat{r_{i_2}}, \dots, \widehat{r_{i_s}}, \dots, r_{k(n-1)+1}, Q^{(s)}) \subseteq \phi(Q)$ for every $i_1, \dots, i_s \in \{1, \dots, k(n-1) + 1\}$ and $1 \leq s \leq (k-1)n - k + 2$.*

Proof. $(f(r_1, \phi(Q), 0^{(m-2)}), \dots, f(r_{k(n-1)+1}, \phi(Q), 0^{(m-2)}))$ is a (k, n) -zero primary of $Q/\phi(Q)$ by Theorem 4.14 and $Q/\phi(Q)$ is a weakly (k, n) -absorbing primary of $R/\phi(Q)$ by Theorem 4.13. Then we conclude that

$$f(g(f(r_1, \phi(Q), 0^{(m-2)}), \dots, f(\widehat{r_{i_1}}, \phi(Q), 0^{(m-2)}), \dots, f(\widehat{r_{i_2}}, \phi(Q), 0^{(m-2)}), \dots, f(\widehat{r_{i_s}}, \phi(Q), 0^{(m-2)}), \dots, f(r_{k(n-1)+1}, \phi(Q), 0^{(m-2)}), Q^{(s)}, \phi(Q), 0^{(m-2)}) = \phi(Q).$$

for every $i_1, \dots, i_s \in \{1, \dots, k(n-1) + 1\}$ and $1 \leq s \leq (k-1)n - k + 2$, by Theorem 4.9 of [16]. Thus, $g(r_1, \dots, \widehat{r_{i_1}}, \dots, \widehat{r_{i_2}}, \dots, \widehat{r_{i_s}}, \dots, r_{k(n-1)+1}, Q^{(s)}) \subseteq \phi(Q)$. \square

Theorem 4.16. *Let R be a commutative Krasner (m, n) -hyperring and let $\phi : \mathcal{HI}(R) \longrightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. Let Q be a ϕ - (k, n) -absorbing primary hyperideal of R but is not a (k, n) -absorbing primary. Then $g(Q^{k(n-1)+1}) \subseteq \phi(Q)$.*

Proof. This can be proved, by using Theorem 4.15, in a very similar manner to the way in which 4.12 was proved. \square

Now, let give some related corollaries.

Corollary 4.17. *Let $\phi : \mathcal{HI}(R) \longrightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function. If Q is a ϕ - (k, n) -absorbing primary hyperideal of R such that $g(Q^{k(n-1)+1}) \not\subseteq \phi(Q)$, then Q is a (k, n) -absorbing primary hyperideal of R .*

Corollary 4.18. *Let $\phi : \mathcal{HI}(R) \longrightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function and let Q be a ϕ - (k, n) -absorbing primary hyperideal of R that is not a (k, n) -absorbing primary hyperideal of R . Then $\mathbf{r}^{(m, n)}(Q) = \mathbf{r}^{(m, n)}(\phi(Q))$.*

Proof. By Theorem 4.16, we have $g(Q^{k(n-1)+1}) \subseteq \phi(Q)$ as Q is not a (k, n) -absorbing primary. This means $\mathbf{r}^{(m, n)}(Q) \subseteq \mathbf{r}^{(m, n)}(\phi(Q))$. On the other hand, from $\phi(Q) \subseteq Q$, it follows that $\mathbf{r}^{(m, n)}(\phi(Q)) \subseteq \mathbf{r}^{(m, n)}(Q)$. Hence $\mathbf{r}^{(m, n)}(Q) = \mathbf{r}^{(m, n)}(\phi(Q))$. \square

Corollary 4.19. *Let $\phi : \mathcal{HI}(R) \rightarrow \mathcal{HI}(R) \cup \{\varphi\}$ be a function and let Q be a proper hyperideal of R such that $\mathbf{r}^{(m,n)}(\phi(Q))$ is a (k, n) -absorbing hyperideal of R . Then Q is a ϕ - $(k+1, n)$ -absorbing primary hyperideal of R if and only if Q is a $(k+1, n)$ -absorbing primary hyperideal of R .*

Proof. (\implies) Let Q be a ϕ - $(k+1, n)$ -absorbing primary hyperideal of R . If Q is not a $(k+1, n)$ -absorbing primary hyperideal of R . Hence $\mathbf{r}^{(m,n)}(Q) = \mathbf{r}^{(m,n)}(\phi(Q))$ by Corollary 4.18. Then $\mathbf{r}^{(m,n)}(Q)$ is a (k, n) -absorbing hyperideal of R which implies that Q is a $(k+1, n)$ -absorbing primary hyperideal of R by Theorem 4.9 in [18].

(\impliedby) It is clear. \square

Theorem 4.20. *Let $h : R_1 \rightarrow R_2$ be a ϕ_1 - ϕ_2 -homomorphism, where ϕ_1 and ϕ_2 are two reduction functions of $\mathcal{HI}(R_1)$ and $\mathcal{HI}(R_2)$, respectively. Then*

- (1) *If Q_2 is a ϕ_2 - (k, n) -absorbing primary hyperideal of R_2 , then $h^{-1}(Q_2)$ is a ϕ_1 - (k, n) -absorbing primary hyperideal of R_1 .*
- (2) *If h is surjective and Q_1 is a ϕ_1 - (k, n) -absorbing primary hyperideal of R_1 with $\text{Ker}(h) \subseteq Q_1$, then $h(Q_1)$ is a ϕ_2 - (k, n) -absorbing primary hyperideal of R_2 .*

Proof. (1) Let Q_2 be a ϕ_2 - (k, n) -absorbing primary hyperideal of R_2 . Assume that $r_1^{kn-k+1} \in R_1$ such that $g(r_1^{kn-k+1}) \in h^{-1}(Q_2) - \phi_1(h^{-1}(Q_2))$. Then we get $h(g(r_1^{kn-k+1})) = g(h(r_1), \dots, h(r_{kn-k+1})) \in Q_2 - \phi_2(Q_2)$. Since Q_2 is a ϕ_2 - (k, n) -absorbing primary hyperideal of R_2 , we obtain either $g(h(r_1), \dots, h(r_{(k-1)n-k+2})) = h(g(r_1^{(k-1)n-k+2})) \in Q_2$ which means $g(r_1^{(k-1)n-k+2}) \in h^{-1}(Q_2)$, or $g(h(r_1), \dots, \widehat{h(r_i)}, \dots, h(r_{kn-k+1})) = h(g(r_1, \dots, \widehat{r_i}, \dots, r_{kn-k+1})) \in \mathbf{r}^{(m,n)}(Q_2)$ which means $g(r_1, \dots, \widehat{r_i}, \dots, r_{kn-k+1}) \in h^{-1}(\mathbf{r}^{(m,n)}(Q_2)) = \mathbf{r}^{(m,n)}(h^{-1}(Q_2))$ for some $1 \leq i \leq n$. Hence $h^{-1}(Q_2)$ is a ϕ_1 - (k, n) -absorbing primary hyperideal of R_1 .

(2) Assume that h is surjective and Q_1 is a ϕ_1 - (k, n) -absorbing primary hyperideal of R_1 with $\text{Ker}(h) \subseteq Q_1$. Let $s_1^{kn-k+1} \in R_2$ such that $g(s_1^{kn-k+1}) \in h(Q_1) - \phi_2(h(Q_1))$. Therefore there exist $r_1^{kn-k+1} \in R_1$ with $h(r_1) = s_1, \dots, h(r_{kn-k+1}) = s_{kn-k+1}$. Hence we get $h(g(r_1^{kn-k+1})) = g(h(r_1), \dots, h(r_{kn-k+1})) = g(s_1^{kn-k+1}) \in h(Q_1)$. Since h is a ϕ_1 - ϕ_2 -epimorphism and $\text{Ker}(h) \subseteq Q_1$, we have $g(r_1^{kn-k+1}) \in Q_1 - \phi_1(Q_1)$. Since Q_1 is a ϕ_1 - (k, n) -absorbing primary hyperideal of R_1 , we conclude that $g(r_1^{(k-1)n-k+2}) \in Q_1$ which implies

$$h(g(r_1^{(k-1)n-k+2})) = g(h(r_1), \dots, h(r_{(k-1)n-k+2})) = g(s_1^{(k-1)n-k+2}) \in h(Q_1),$$

or $g(r_1, \dots, \widehat{r_i}, \dots, r_{kn-k+1}) \in \mathbf{r}^{(m,n)}(Q_1)$ implies $h(g(r_1, \dots, \widehat{r_i}, \dots, r_{kn-k+1})) = g(h(r_1), \dots, \widehat{h(r_i)}, \dots, h(r_{kn-k+1})) = g(s_1, \dots, \widehat{s_i}, \dots, s_{kn-k+1}) \in h(\mathbf{r}^{(m,n)}(Q_1)) \subseteq \mathbf{r}^{(m,n)}(h(Q_1))$ for some $1 \leq i \leq (k-1)n - k + 2$. Consequently, $h(Q_1)$ is a ϕ_2 - (k, n) -absorbing primary hyperideal of R_2 . \square

As an instant consequence of the previous theorem, we get the following explicit result.

Theorem 4.21. *Let Q and P be two hyperideals of R and ϕ be a reduction function of $\mathcal{HI}(R)$ such that $P \subseteq \phi(Q) \subseteq Q$. If Q is a ϕ - (k, n) -absorbing primary hyperideal of R , then Q/P is a ϕ_q - (k, n) -absorbing primary hyperideal of R/P .*

Theorem 4.22. *Let (R_i, f_i, g_i) be a commutative Krasner (m, n) -hyperring for each $1 \leq i \leq kn - k + 1$ and $\phi_i : \mathcal{HI}(R_i) \rightarrow \mathcal{HI}(R_i) \cup \{\varphi\}$ be a function. Let Q_i be a hyperideal of R_i for each $1 \leq i \leq kn - k + 1$ and $\phi = \phi_1 \times \cdots \times \phi_{kn - k + 1}$. If $Q = Q_1 \times \cdots \times Q_{kn - k + 1}$ is a ϕ - $(k + 1, n)$ -absorbing primary hyperideal of $R = R_1 \times \cdots \times R_{kn - k + 1}$, then Q_i is a ϕ_i - (k, n) -absorbing primary hyperideal of R_i and $Q_i \neq R_i$ for all $1 \leq i \leq kn - k + 1$.*

Proof. By using an argument similar to that in the proof of Theorem 3.7, one can easily complete the proof. \square

5. CONCLUSION

In this paper, motivated by the research works on ϕ -2-absorbing (primary) ideals of commutative rings, we proposed and investigated the notions of ϕ - (k, n) -absorbing and ϕ - (k, n) -absorbing primary hyperideals in a Krasner (m, n) -hyperring. Some of their essential characteristics were analysed. Moreover, the stability of the notions were examined in some hyperring-theoretic constructions. As a new research subject, we suggest the concept of ϕ - (k, n) -absorbing δ -primary hyperideals, where δ is an expansion function of $\mathcal{HI}(R)$.

6. ACKNOWLEDGMENTS

The author wish to sincerely thank the referees for several useful comments.

REFERENCES

- [1] R. Ameri and M. Norouzi, *Prime and primary hyperideals in Krasner (m, n) -hyperrings*, European J. Combin., **34** (2013) 379-390.
- [2] R. Ameri, A. Kordi and S. Sarka-Mayerova, *Multiplicative hyperring of fractions and coprime hyperideals*, An. Stiint. Univ. Ovidius Constanta Ser. Mat., **25** No. 1 (2017) 5-23.
- [3] M. Anbarloei, *n -ary 2-absorbing and 2-absorbing primary hyperideals in Krasner (m, n) -hyperrings*, Mat. Vesn., **71** No. 3 (2019) 250-262.
- [4] M. Anbarloei, *Unifying the prime and primary hyperideals under one frame in a Krasner (m, n) -hyperring*, Comm. Algebra, **49** (2021) 3432-3446.
- [5] M. Anbarloei, *A study on a generalization of the n -ary prime hyperideals in Krasner (m, n) -hyperrings*, Afr. Mat., **33** (2021) 1021-1032.
- [6] M. Anbarloei, *Krasner (m, n) -hyperring of fractions*, Jordan J. Math. Stat., **16** No. 1 (2023) 165-185.
- [7] D. D. Anderson and M. Bataineh, *Generalizations of prime ideals*, Comm. Algebra, **36** (2008) 686-696.

- [8] A. Asadi and R. Ameri, *Direct limit of Krasner (m,n) -hyperrings*, J. Sci., **31** No. 1 (2020) 75-83.
- [9] A. Badawi, U. Tekir, E. A. Ugurlu, G. Ulucak and E. Y. Celikel, *Generalizations of 2-absorbing primary ideals of commutative rings*, Turkish J. Math., **40** No. 3 (2016) 703-717.
- [10] S. Corsini, *Prolegomena of Hypergroup Theory*, Second edition, Aviani editor, Italy, 1993.
- [11] S. Corsini and V. Leoreanu, *Applications of Hyperstructure Theory, Advances in Mathematics*, Vol. 5, Kluwer Academic Publishers, 2003.
- [12] A. Y. Darani, *Generalizations of primary ideals in commutative rings*, Novi Sad J. Math., **42** No.1 (2012) 27-35.
- [13] B. Davvaz and V. Leoreanu-Fotea, *Hyperring Theory and Applications*, International Academic Press, Palm Harbor, USA, 2007.
- [14] B. Davvaz and T. Vougiouklis, *n -ary hypergroups*, Iran. J. Sci. Technol., **30** No. A2 (2006) 165-174.
- [15] B. Davvaz, *Fuzzy Krasner (m,n) -hyperrings*, Comput. Math. Appl., **59** (2010) 3879-3891.
- [16] B. Davvaz, G. Ulucak and U. Tekir, *Weakly (k,n) -absorbing (Primary) hyperideals of a Krasner (m,n) -hyperring*, Hacet. J. Math. Stat., **52** No. 5 (2023) 1229-1238
- [17] W. Dorente, *Untersuchungen über einen verallgemeinerten Gruppenbegriff*, Math. Z., **29** (1928) 1-19.
- [18] K. Hila, K. Naka and B. Davvaz, *On (k,n) -absorbing hyperideals in Krasner (m,n) -hyperrings*, Q. J. Math., **69** (2018) 1035-1046.
- [19] A. Jaber, *Properties of ϕ - δ -primary and 2-absorbing δ -primary ideals of commutative rings*, Asian-Eur. J. Math., **13** No. 1 (2020), 2050026.
- [20] A. Khaksari, *ϕ -2-prime ideals*, Int. J. Pure Appl. Math., **99** No. 1 (2015) 1-10.
- [21] E. Kasner, *An extension of the group concept (reported by L.G. Weld)*, Bull. Amer. Math. Soc., **10** (1904) 290-291.
- [22] V. Leoreanu-Fotea, *Canonical n -ary hypergroups*, Ital. J. Pure Appl. Math., **24** (2008) 247-254.
- [23] V. Leoreanu-Fotea and B. Davvaz, *n -hypergroups and binary relations*, European J. Combin., **29** (2008) 1027-1218.
- [24] V. Leoreanu-Fotea and B. Davvaz, *Roughness in n -ary hypergroups*, Inform. Sci., **178** (2008) 4114-4124.
- [25] X. Ma, J. Zhan and B. Davvaz, *Applications of rough soft sets to Krasner (m,n) -hyperrings and corresponding decision making methods*, Filomat, **32** (2018) 6599-6614.
- [26] F. Marty, *Sur une generalization de la notion de groupe*, In 8th Congress Math. Scandenaves, Stockholm, 1934.
- [27] S. Mirvakili and B. Davvaz, *Relations on Krasner (m,n) -hyperrings*, European J. Combin., **31** (2010) 790-802.
- [28] S. Mirvakili and B. Davvaz, *Constructions of (m,n) -hyperrings*, Mat. Vesn., **67** No. 1 (2015) 1-16.
- [29] M. Norouzi, R. Ameri and V. Leoreanu-Fotea, *Normal hyperideals in Krasner (m,n) -hyperrings*, An. St. Univ. Ovidius Constanta, **26** No. 3 (2018) 197-211.
- [30] S. Omidi and B. Davvaz, *Contribution to study special kinds of hyperideals in ordered semihyperrings*, J. Taibah Univ. Sci., **11** (2017) 1083-1094.
- [31] S. Ostadhadi-Dehkordi and B. Davvaz, *A Note on isomorphism theorems of Krasner (m,n) -hyperrings*, Arab. J. Math., **5** (2016) 103-115.
- [32] T. Vougiouklis, *Hyperstructures and their Representations*, Hadronic Press Inc., Florida, 1994.

- [33] M. M. Zahedi and R. Ameri, *On the prime, primary and maximal subhypermultiples*, Ital. J. Pure Appl. Math., **5** (1999) 61-80.
- [34] J. Zhan, B. Davvaz and K. P. Shum, *Generalized fuzzy hyperideals of hyperrings*, Comput. Math. Appl., **56** (2008) 1732-1740.

Mahdi Anbarloei

Department of Mathematics, Faculty of Sciences

Imam Khomeini International University

Qazvin, Iran.

`m.anbarloei@sci.ikiu.ac.ir`