



Research Paper

THE STRONGLY IRREDUCIBLE DIMENSION OF RINGS VS. THE DERIVED DIMENSION OF THE SPACE OF STRONGLY IRREDUCIBLE IDEALS WITH V-TOPOLOGY

JAMAL HASHEMI* AND FATEMEH HASSANZADEH

ABSTRACT. An ideal I of a ring R is called strongly irreducible ideal (SI-ideal, for short), whenever the inclusion $J \cap K \subseteq I$, implies that $J \subseteq I$ or $K \subseteq I$. Let $X = \text{SSpec}(R)$ be the set of all strongly irreducible ideals of a ring R . Then X with certain topology has derived dimension if and only if R has strongly irreducible dimension. Moreover, the two dimensions differ by at most 1.

1. INTRODUCTION

Karamzadeh in [4] has proved a useful result connecting two different concepts from topological spaces and general rings which is toward the unity in mathematics. This in particular shows that given any non-limit ordinal $\alpha = \beta + 1$ we may use the set of $X = \text{Spec}(R)$ with a certain quasi-compact topology, where R is a Noetherian domain with Krull-dimension β

DOI: 10.22034/as.2024.21459.1710

MSC(2010): Primary: 13C15, 16P70.

Keywords: Arithmetical ring, Derived dimension, Strongly irreducible ideal, Strongly regular ring, V-topology.

Received: 12 April 2024, Accepted: 17 September 2024.

*Corresponding author

(note, such a domain always exists, see [4, Note added in proof]), to provide a quasi-compact space with given a non-limit ordinal as its derived dimension which is important in topology (note, derived dimension of a topological space is also called Cantor-Bendixon dimension, and it is well-known that this dimension for quasi-compact spaces is always non-limit ordinal, see [6, P. 176]. Motivated by this fact in [4], in what follows we are going to set the stage for proving a similar connecting result. A proper ideal I of a ring R is said to be strongly irreducible (briefly, SI-ideal) if for each pair of ideals J and K of R , $J \cap K \subseteq I$ implies that either $J \subseteq I$ or $K \subseteq I$. A strongly irreducible ring R (briefly, SI-ring) is a ring in which 0 is strongly irreducible. A minimal strongly irreducible ideal in a ring R is any strongly irreducible ideal of R such that it does not properly contain any other strongly irreducible ideal. It is easy to see that every SI-ideal contains a minimal strongly irreducible ideal, for more details see [1]. If I is an ideal in a ring R , the set of all minimal strongly irreducible ideals Q containing I is denoted by $\text{siMin}(I)$. In what follows we recall the classical Krull-dimension of a ring R . Let $X = \text{Spec}(R)$ be the set of all prime ideals in R and $\text{Spec}_0(R)$ denotes the set of all maximal ideals of R . For an ordinal $\alpha > 0$ denote $\text{Spec}_\alpha(R)$ to be the set of all prime ideals P in R such that whenever a prime ideal Q properly contains P , then Q belongs to $\text{Spec}_\beta(R)$ for some $\beta < \alpha$. The smallest ordinal α such that $\text{Spec}_\alpha(R) = X$ is called classical Krull-dimension of R , denoted by $\text{cl.K.dim}(R)$. It is well-known that $\text{cl.K.dim}(R)$ exists if and only if R has acc on prime ideals, for more details, see [2, 4]. Motivated by the above definition we introduce a new dimension for a ring R using SI-ideals instead of prime ideals and denote by $\text{si.dim}(R)$. In this article, we extend the results of [4], in the same vein. For example we show that the existence of $\text{si.dim}(R)$ is equivalent having acc on SI-ideals. Also similarly we study the derived dimension of a topological space defined on $X = \text{SSpec}(R)$ (The set of all SI-ideals). In particular we show that this derived dimension exists if and only if $\text{si.dim}(R)$ exists and they differ by at most one. In this article all rings are associative with $1 \neq 0$.

2. STRONGLY IRREDUCIBLE DIMENSION

Definition 2.1. Let $\text{SSpec}(R)$ be the set of all SI-ideals of a ring R and $\text{SSpec}_0(R)$ denote the set of all maximal ideals of R . Let $\alpha > 0$ be an ordinal and define $\text{SSpec}_\alpha(R)$ to be the set of all ideals Q of R such that for any SI-ideal Q' where $Q \subset Q'$, then Q' belongs to $\text{SSpec}_\beta(R)$ for some $\beta < \alpha$. Then the smallest ordinal α for which $\text{SSpec}(R) = \text{SSpec}_\alpha(R)$ is called *strongly irreducible dimension of R* , denoted by $\text{si.dim}(R)$.

The following result provides an equivalent condition for the existence of $\text{si.dim}(R)$ in a ring R .

Theorem 2.2. *A ring R has acc on SI-ideals if and only if $\text{si.dim}(R)$ exists.*

Proof. First suppose that R has acc on SI-ideals and define the sets $\text{SSpec}_\alpha(R)$ of SI-ideals as in the previous definition. Since for each α , $\text{card}(\text{SSpec}_\alpha(R)) \leq 2^{\text{card}(R)}$, the chain $\text{SSpec}_0(R) \subseteq \text{SSpec}_1(R) \subseteq \dots$ cannot be properly increasing forever. Hence, there exists an ordinal γ such that $\text{SSpec}_\gamma(R) = \text{SSpec}_{\gamma+1}(R)$. If $\text{si.dim}(R)$ does not exist, then using acc on SI-ideals, there is an SI-ideal Q which is maximal with respect to the property $Q \notin \text{SSpec}_\alpha(R)$. Hence all SI-ideals properly containing Q lie in $\text{SSpec}_\alpha(R)$. Therefore, we infer that $Q \in \text{SSpec}_{\alpha+1}(R) = \text{SSpec}_\alpha(R)$, which is a contradiction.

For the converse, we show that every nonempty set of $\text{SSpec}(R)$ has a maximal element. So, let $\text{si.dim}(R)$ exists and S be a nonempty set of $\text{SSpec}(R)$. Therefore, there is an ordinal $\alpha \geq 0$ such that $\text{SSpec}_\alpha(R) = \text{SSpec}(R)$, and we may assume that β is the smallest ordinal such that $S \cap \text{SSpec}_\beta(R)$ is not empty. Let $Q \in S \cap \text{SSpec}_\beta(R)$ and assume there is an element Q' in S which contains Q properly. Then $Q' \in \text{SSpec}_\gamma(R)$ for some $\gamma < \beta$ and hence $Q' \in \text{SSpec}_\gamma(R) \cap S$ for some $\gamma < \beta$, and this contradicts the fact that β is minimal. Thus Q is maximal in S , i.e., R has acc on SI-ideals. \square

Remark 2.3. Clearly that every prime ideal of a ring R is an SI-ideal. Thus, if R is a ring with no acc on prime ideals (i.e., $\text{cl.K.dim}(R)$ does not exist, see [2, Ex. 14A(b)] and [2, Proposition 14.1]), then from theorem 2.2, we conclude that $\text{si.dim}(R)$ also does not exist.

Example 2.4. Let \mathbb{Z} be the ring of integers. It is clear that every nonzero strongly irreducible ideal in \mathbb{Z} is of the form $p^i\mathbb{Z}$, where p is a prime number and i is a positive integer. Therefore for any nonnegative integer n , $\text{SSpec}_n(\mathbb{Z})$ consists of all ideals of the form $p^i\mathbb{Z}$ where $1 \leq i \leq n + 1$. So $\text{si.dim}(\mathbb{Z}) = \omega$, where ω is the first infinite ordinal. The same result holds for any commutative principal ideal domain that is not a field.

Example 2.5. Let R be a *strongly regular ring* (i.e., $a = a^2b$ for any $a \in R$ and some $b \in R$). Then $\text{cl.K.dim}(R) = \text{si.dim}(R) = 0$

Example 2.6. If R is a polynomial ring in an infinite number of indeterminates over a field k , then R does not have acc for prime ideals. Therefore according to Remark 2.3, $\text{si.dim}(R)$ is not exist.

Remark 2.7. We observe that if Q is an SI-ideal in R containing an ideal I , then Q/I is an SI-ideal of R/I . For the converse we also observe that if R is an arithmetical ring (note, we recall that a ring R is called arithmetical ring whenever the lattice of all ideals of R is distributive, i.e., for any three ideals I, J and K in R , $I + (J \cap K) = (I + J) \cap (I + K)$, or equivalently, $I \cap (J + K) = (I \cap J) + (I \cap K)$) and Q/I is an SI-ideal of R/I , then Q is an SI-ideal in R . To this end, let $J \cap K \subseteq Q$ then $(J \cap K) + I = (J + I) \cap (K + I) \subseteq Q$ and consequently

$(J + I)/I \cap (K + I)/I \subseteq Q/I$. Since Q/I is SI-ideal, we infer that $(J + I)/I \subseteq Q/I$ or $(K + I)/I \subseteq Q/I$. Hence $J \subseteq Q$ or $K \subseteq Q$, i.e., Q is an SI-ideal.

Lemma 2.8. *Let R be an arithmetical ring and $\alpha \geq 0$ be an ordinal. Then:*

(1) *If I is an ideal contained in the SI-ideal Q , then $Q \in \text{SSpec}_\alpha(R)$ if and only if $Q/I \in \text{SSpec}_\alpha(R/I)$.*

(2) *$\text{si.dim}(R) = \alpha$ implies $\text{si.dim}(R/I) \leq \alpha$ for every ideal I of R .*

(3) *If R is an SI-ring with $\text{si.dim}(R) = \alpha$ and $Q \neq 0$ is an SI-ideal, then $\text{si.dim}(R/Q) < \alpha$.*

Proof. (1) The statement is obvious for $\alpha = 0$. Let $\alpha \geq 0$ and assume (1) holds for all $\beta < \alpha$. From the definition of $\text{SSpec}_\alpha(R)$ and the induction hypothesis we get: $Q \in \text{SSpec}_\alpha(R)$ if and only if $Q \subset Q' \in \text{SSpec}(R)$ implies $Q' \in \text{SSpec}_\beta(R)$ for some $\beta < \alpha$. And we also note that $Q/I \subset Q'/I \in \text{SSpec}(R/I)$ implies $Q'/I \in \text{SSpec}_\beta(R/I)$ for some $\beta < \alpha$, if and only if $Q/I \in \text{SSpec}_\alpha(R/I)$.

(2) If $Q/I \in \text{SSpec}(R/I)$, then $I \subseteq Q \in \text{SSpec}(R) = \text{SSpec}_\alpha(R)$. By (1), $Q/I \in \text{SSpec}_\alpha(R/I)$ and hence $\text{si.dim}(R/I) \leq \alpha$.

(3) Since R is an SI-ring and $0 \subset Q$, clearly $Q \in \text{SSpec}_\beta(R)$ for some $\beta < \alpha$. Now, if $Q'/Q \in \text{SSpec}(R/Q)$ then $Q' \in \text{SSpec}_\gamma(R)$ for some $\gamma < \beta$ and hence $Q'/Q \in \text{SSpec}_\gamma(R/Q)$ by (1). Thus $\text{SSpec}(R/Q) \subseteq \text{SSpec}_\gamma(R/Q)$ which implies that $\text{si.dim}(R/Q) < \alpha$. \square

The following corollary is now immediate.

Corollary 2.9. *Let R be an arithmetical ring and $\text{si.dim}(R)$ exists. If P and Q are SI-ideals with $P \subset Q$, then $\text{si.dim}(R/Q) < \text{si.dim}(R/P)$.*

In what follows we present several more results concerning SI-ideal in arithmetical rings.

Lemma 2.10. *Let R be an arithmetical ring with $\text{si.dim}(R) = \alpha$. If $\beta \geq 0$ is any ordinal strictly less than α , then there is an SI-ideal Q such that $\text{si.dim}(R/Q) = \beta$. If R is a right or left Neotherain, then there is a minimal SI-ideal Q such that $\text{si.dim}(R/Q) = \alpha$.*

Proof. First, we observe that for an SI-ideal Q , we have $\text{si.dim}(R/Q) = \beta$ if and only if β is the smallest ordinal such that $Q \in \text{SSpec}_\beta(R)$. If there is no SI-ideal Q such that $\text{si.dim}(R/Q) = \beta$, then we must have $\text{SSpec}_\beta(R) = \text{SSpec}_{\beta+1}(R)$, which implies that $\text{SSpec}_\beta(R) = \text{SSpec}_\gamma(R)$ for any $\beta < \gamma$. Therefore, we infer that $\text{si.dim}(R) \leq \beta$, which is a contradiction. Here, we observe that this argument especially shows that $\text{si.dim}(R)$ is the supremum of the ordinals $\text{si.dim}(R/Q)$, where Q ranges over the set of SI-ideals, and it is clear that we may restrict the

set of SI-ideals to the set of minimal SI-ideals, for, every SI-ideals contains a minimal SI-ideal. If R is right or left Noetherian, then there are only finitely many minimal SI-ideals, therefore there is a minimal SI-ideal, Q say, such that $\text{si.dim}(R/Q) = \alpha$. \square

Lemma 2.11. *Let R be an arithmetical ring with $\text{si.dim}(R) \geq \alpha \geq 0$. If $\text{si.dim}(R/I) < \alpha$ for every ideal $I \neq 0$, then R is an SI-ring with $\text{si.dim}(R) = \alpha$.*

Proof. We show that $\text{SSpec}_\alpha(R) = \text{SSpec}(R)$. Let P and Q be two SI-ideals with $P \subset Q$. Since $\text{si.dim}(R/P) = \beta < \alpha$ and Q/P is an SI-ideal of R/P , by Lemma 2.8(1), we get $Q \in \text{SSpec}_\beta(R)$. Thus $P \in \text{SSpec}_\alpha(R)$ for all $P \in \text{SSpec}(R)$, and this yields $\text{si.dim}(R) \leq \alpha$, i.e., $\text{si.dim}(R) = \alpha$.

Now suppose that R is not an SI-ring, therefore there are nonzero SI-ideals I and J such that $I \cap J = 0$. Let β be the maximum of $\text{si.dim}(R/I)$ and $\text{si.dim}(R/J)$. It is sufficient to prove that $\text{SSpec}_\beta(R) = \text{SSpec}(R)$, which is a contradiction because we show that $\text{si.dim}(R) = \alpha$. To this end, suppose that P is an SI-ideal of R . Since $I \cap J = 0 \subseteq P$, we may assume that $I \subseteq P$. Hence (P/I) is an SI-ideal of (R/I) (note that R is an arithmetical ring). Now by Lemma 2.8 (2), $\text{si.dim}(R/I)/(P/I) \leq \text{si.dim}(R/I)$, so $\text{si.dim}(R/P) \leq \beta < \alpha$, i.e., $(P/I) \in \text{SSpec}_\beta(R/I)$ and consequently $P \in \text{SSpec}_\beta(R)$. \square

Lemma 2.12. *Let R be an arithmetical ring with acc on two-sided ideals. The following are equivalent:*

- (1) R is an SI-ring.
- (2) For every SI-ideal $Q \neq 0$, $\text{si.dim}(R/Q) < \text{si.dim}(R)$.
- (3) For every ideal $I \neq 0$, $\text{si.dim}(R/I) < \text{si.dim}(R)$.

Proof. (1) implies (2) by Lemma 2.8 (3). (3) implies (1) by the previous lemma. Finally, in what follows we prove (2) implies (3). Suppose that (2) holds and let I be an ideal which is maximal with respect to $\text{si.dim}(R/I) = \text{si.dim}(R)$. Now, suppose that K/I is a nonzero ideal of R/I , then by the maximality of I , we have $\text{si.dim}(R/I)/(K/I) = \text{si.dim}(R/K) < \text{si.dim}(R)$. Thus by the previous lemma, (R/I) is an SI-ring. So I is an SI-ideal and by (2), we get $I = 0$.

\square

The following corollary is evident.

Corollary 2.13. *Let R be a right or left Noetherian arithmetical ring and $\text{si.dim}(R)$ exists. If Q is an SI-ideal of R and I is an ideal of R with $Q \subset I$, then $\text{si.dim}(R/I) < \text{si.dim}(R/Q)$.*

Proposition 2.14. *Let R be a ring with acc on SI-ideals. Then R has acc on ideals I of the form $I_k = \bigcap_{Q \in F_k} Q$, where F_k is a finite set of noncomparable SI-ideals.*

Proof. Let $I_1 \subseteq I_2 \subseteq \cdots \subseteq I_n \subseteq \cdots$ be an infinite ascending chain of ideals, each of which is of the form $I_k = \bigcap_{P \in F_k} P$, where F_k is a finite set of noncomparable SI-ideals. If it happens that $F_{r_1} = F_{r_2} = \cdots = F_{r_n} = \cdots$, where $r_1 < r_2, \dots < r_n < \cdots$ is an infinite sequence, then $I_{r_1} = I_{r_2} = \cdots = I_{r_n} = \cdots$ and we are through. Now we may assume $F_{n+1} - F_n \neq \emptyset$, for all n and complete the proof by obtaining a contradiction. We note that $F_i \cap F_r \subseteq F_{i-1} \cap F_r$ for all r and $r \leq i-1$, for if not, then there exists $Q_i \in F_i \cap F_r$ such that $Q_i \notin F_{i-1}$. Hence there exists $Q_{i-1} \in F_{i-1}$ such that $Q_{i-1} \subset Q_i$ and since $r \leq i-1$, there exists $Q_r \in F_r$ such that $Q_r \subseteq Q_{i-1} \subset Q_i$. But Q_r and Q_i are both in F_r and can not be comparable. This shows that without loss of generality we can assume that $F_{i-1} \cap F_r = F_i \cap F_r$, for all r and $r \leq i-1$. Now let $Q_m \in F_m - F_{m-1}$, for any integer $m > 0$. Then $Q_m \notin \bigcup_{i=1}^{m-1} F_i$, for otherwise $Q_m \in F_r$, for some $r \leq m-1$ and $F_m \cap F_r = F_{m-1} \cap F_r$ implies that $Q_m \in F_{m-1}$, which is impossible. Hence there exists $Q_{m-1} \in F_{m-1}$ such that $Q_{m-1} \subset Q_m$ and $Q_{m-1} \notin \bigcup_{i=1}^{m-2} F_i$, for otherwise $Q_{m-1} \in F_{m-1} \cap F_r$ for some $r \leq m-2$ implies that $Q_{m-1} \in F_m$, which is impossible. Repeating this process we get a chain $Q_1 \subset Q_2 \subset \cdots \subset Q_n$ of SI-ideals such that each Q_i belongs to F_i . Now put $F_1^n = \{Q_1 \in F_1 : \text{there exists a chain } Q_1 \subset Q_2 \subset \cdots \subset Q_n, \text{ where } Q_i \in F_i, i = 1, \dots, n\}$. We have already shown that $F_1^n \neq \emptyset$ for all n . Moreover, F_1^n is finite and $F_1^n \subseteq F_1^m$, for $m \leq n$. Therefore the chain $F_1^1 \supseteq F_1^2 \supseteq \cdots \supseteq F_1^n \supseteq \cdots$ stationary and we can choose $Q'_1 \in \bigcap_{n=1}^{\infty} F_1^n$. Now for each $n \geq 2$, let $F_2^n = \{Q_2 \in F_2 : \text{there exists a chain } Q'_1 \subset Q_2 \subset \cdots \subset Q_n \text{ where } Q_i \in F_i, i = 2, \dots, n\}$. It is clear that $F_2^n \neq \emptyset$ for all n . Now we can choose $Q'_2 \in \bigcap_{n=2}^{\infty} F_2^n$. Hence proceeding inductively we get a chain $Q'_1 \subset Q'_2 \subset \cdots \subset Q'_n \subset \cdots$ of SI-ideals, which is the desired contradiction. \square

First let us recall that the Z-topology on the set of prime ideals in noncommutative rings as in [4]. If A is an ideal of R we let $V(A)$ denote the subset of $\text{Spec}(R)$ consisting of those prime ideals that contain A . Now just by replacing the prime ideal in the previous definition by SI-ideals we get a topology on the SI-ideals, see also [1].

It is shown that in [4] that the set of $X - V(A)$ satisfy the axioms for open sets in this topological space and we call it the *SIZ-topology* on $X = \text{SSpec}(R)$. Put $B = \{V(A) : A \text{ is an ideal of } R\}$, then clearly B can be take as a base for open sets on $\text{SSpec}(R)$. This topology is also called V-topology. The name V-topology is first introduced by Karamzadeh in [4].

The following result is in [4], without proof. Next we give a proof for the sake of the reader.

Proposition 2.15. *Let $X = \text{SSpec}(R)$ be with the SIZ-topology, then the following statements are equivalent:*

- (1) X has acc on open subsets.
- (2) Every subset of X is quasi-compact.
- (3) X has acc on intersections of SI-ideals.

Proof. (1) \Rightarrow (2) Let $A \subseteq \bigcup_{\lambda \in \Lambda} O_\lambda$, where each O_λ is an open subset of X and A is a subset of X . Since X has acc on open subsets, then $T = \{\bigcup_{i=1}^n O_{\lambda_i} : \lambda_i \in \Lambda, n \in \mathbb{N}\}$ has maximal element, say $\bigcup_{i=1}^n O_{\lambda_i}$. It is clear that $A \subseteq \bigcup_{i=1}^n O_{\lambda_i}$.

(2) \Rightarrow (3) Let $I_1 \subseteq I_2 \subseteq \dots$ be an infinite ascending chain of ideals, where each I_k is an intersection of a family of SI-ideals. Then

$$X - V(I_1) \subseteq X - V(I_2) \subseteq \dots.$$

Let $A = \bigcup_{k \in \mathbb{N}} (X - V(I_k))$, then By assumption, there exists $n \in \mathbb{N}$ such that $A = X - V(I_n)$. It follows that $V(I_n) = V(I_{n+k})$, and consequently $I_n = I_{n+k}$ for all $k \in \mathbb{N}$.

(3) \Rightarrow (1) Since every open set of X is of the form $X - V(I)$, where I is an ideal of R , so to prove 3, it suffices to prove that X has the dcc on closed sets. Hence let $V(I_1) \supseteq V(I_2) \supseteq \dots$ be an infinite descending chain of closed subset of X . It is clear that $\bigcap V(I_1) \subseteq \bigcap V(I_2) \subseteq \dots$ and by hypothesis, there exists $n \in \mathbb{N}$ such that $\bigcap V(I_n) = \bigcap V(I_{n+k})$, for all $k \in \mathbb{N}$. Since for every ideal I of R , $V(\bigcap V(I)) = V(I)$, we infer that $V(I_n) = V(I_{n+k})$ for all $k \in \mathbb{N}$, and the proof is complete. \square

In what follows we prove some needed results which are counterparts of similar results in [4]

Corollary 2.16. *If R is a ring with $\text{si.dim}(R)$ and has only finitely many minimal SI-ideals over any ideal, then every SI-ideal is minimal over some finitely generated subideal.*

Proof. If I is an ideal in R , let $P(I)$ denote the intersection of SI-ideal containing I . It is sufficient to show that $P(I) = P(\langle x_1, x_2, \dots, x_n \rangle)$, where $\langle x_1, x_2, \dots, x_n \rangle$ is the ideal generated by $x_1, x_2, \dots, x_n \in I$. It is clear that $V(I) = \bigcap_{x \in I} V(\langle x \rangle)$ and $X - V(I) = \bigcup_{x \in I} (X - V(\langle x \rangle))$. Now by Proposition 2.14, R has acc on intersections of SI-ideals. Therefore Proposition 2.15 shows that every subset and in particular $X - V(I)$ is quasi-compact. Thus there are some element $x_1, x_2, \dots, x_n \in I$ such that $X - V(I) = \bigcup_{i=1}^n (X - V(\langle x_i \rangle))$, hence $V(I) = \bigcap_{i=1}^n V(\langle x_i \rangle)$ implies that $P(I) = P(\langle x_1, x_2, \dots, x_n \rangle)$. \square

We cite the following proposition from [3], which is counterpart of the prime avoidance lemma for SI-ideals. We give the proof for the sake of the reader.

Proposition 2.17. *Let $I, Q_1, Q_2, \dots, Q_n, n \geq 2$, be ideals of a ring R and $I \subseteq \bigcup_{i=1}^n Q_i$. If at most two of the Q_i 's are not SI-ideal, then $I \subseteq Q_i$ for some Q_i .*

Proof. For $n = 2$, the assertion holds, even if Q_1 and Q_2 are not SI-ideal, which is a classical result in ring theory. Now assume $n \geq 3$. In this case, without loss of generality we may assume that Q_1 is an SI-ideal and $Q_i \not\subseteq Q_j$ for $i \neq j$. Also by induction we may assume that

$I \not\subseteq \bigcup_{i=2}^n Q_i$. Hence there is $x \in I$ such that $x \notin \bigcup_{i=2}^n Q_i$. We show that $I \subseteq Q_1$ and we are done. Let us put $J = \bigcap_{i=2}^n Q_i$ and note that for each $y \in I \cap J$ we have $x + y \notin Q_i$ for all $i \geq 2$. Therefore $x + y \in Q_1$ which means $y \in Q_1$ and so $I \cap J \subseteq Q_1$. Since Q_1 is an SI-ideal and $J \not\subseteq Q_1$, we infer that $I \subseteq Q_1$. \square

The next proposition is the counterpart of Proposition 2 in [4]. But before that, we express the concept of the rank of an SI-ideal. Let Q, Q_1, \dots, Q_n be distinct SI-ideals and $Q = Q_0 \supset Q_1 \supset \dots \supset Q_n$, then we say that this chain is of length n . Now, we say that Q has rank n which is abbreviated by $\text{rank}(Q) = n$, if there exists a chain of length n descending from Q , but no longer chain. If for any positive integer n , there exists a descending chain from Q of length n , then we say that Q has rank ∞ . We note that a minimal SI-ideal has rank 0.

Proposition 2.18. *Let R be an arithmetical ring with $\text{si.dim}(R) = n$ and have only finitely many minimal SI-ideals over any ideal, then every SI-ideal is minimal over a subideal generated by less than or equal to n elements.*

Proof. Let $\text{si.dim}(R) = n$, Q is an SI-ideal and Q_1, Q_2, \dots, Q_t be all minimal SI-ideals. By Corollary 2.9, it is clear that for any SI-ideal Q , $\text{rank}(Q) \leq \text{si.dim}(R)$. We may proceed by induction on $k = \text{rank}(Q)$ and show that Q is minimal over a subideal which is generated by $\leq k$ elements. For $\text{rank}(Q) = 0$ is clear. Now suppose that it is true for $\text{rank}(Q) \leq k - 1$. And let $\text{rank}(Q) = k$ where $k > 0$. By Proposition 2.17, we infer that $Q \not\subseteq \bigcup_{i=1}^t Q_i$ because $k > 0$, so there exists $a_1 \in Q$ such that $a_1 \notin \bigcup_{i=1}^t Q_i$. Now let $\pi : R \rightarrow R/\langle a_1 \rangle$ with $\pi(x) = \bar{x}$ be the canonical projection and we observe that $\text{rank}(Q/\langle a_1 \rangle) \leq k - 1$. Therefore $Q/\langle a_1 \rangle$ is minimal over $\langle \bar{a}_2, \bar{a}_3, \dots, \bar{a}_k \rangle$ by hypothesis induction, then it is clear that Q is minimal over $\langle a_1, a_2, \dots, a_k \rangle$. \square

3. DRIVED DIMENSION OF A TOPOLOGICAL SPACE

Recall that in a topological space X an element $x \in X$ is called a limit point of a subset A of X if each open set of X contains at least one point of A distinct from x . The set of all limit points of A is denoted by A' and is called the *drived* set of A . A point $a \in A$ is called *isoleted* whenever $a \in A - A'$.

Without further ado we begin with the definition of the above dimension.

Definition 3.1. The α -derivative of a topological space X is defined by transfinite induction: $X_0 = X$, $X_{\alpha+1} = X'_\alpha$ and $X_\alpha = \bigcap_{\beta < \alpha} X_\beta$, for a limit ordinal α . If for an ordinal α we have $X_\alpha = \emptyset$ then X is called *scattered*. If X is scattered and α is the smallest ordinal such that $X_\alpha = \emptyset$, then α is called *derived dimension* of X and is denoted by $d(X) = \alpha$, for more information see [4].

The following lemma is well known, see[4].

Lemma 3.2. *Let X be a topological space, then the following are equivalent.*

- (1) *Every non empty subset of X contains an isolated point.*
- (2) *There is an $\alpha > 0$ such that $X_\alpha = \emptyset$*

The following is the counterpart of [4, lemma 4].

Lemma 3.3. *Let $X = \text{SSpec}(R)$ be the space with the V-topology and $S \subseteq X$, then an element $Q \in S$ is an isolated point of S if and only if it is a maximal element of S .*

Proof. If $Q \in S$ is maximal element of S then $V(Q) \cap S = \{Q\}$ shows that P is an isolated point of S . Now suppose that Q is an isolated point of S then there exists an open subset G such that $P \in G$ and $G \cap S = \{Q\}$. But there exists $V(A)$ such that $Q \in V(A) \subseteq G$, then $V(A) \cap S = \{Q\}$. Now we claim that Q is a maximal in S . If $Q \subset Q'$ and $Q' \in S$ then $Q' \in V(A)$ which is impossible. \square

We need the next proposition which is also the counterpart of [4, Corollary 3].

Proposition 3.4. *Let $X = \text{SSpec}(R)$ be the space with the V-topology. Then $\text{SSpec}_\alpha(R) = \bigcup_{\beta \leq \alpha} S_\beta$, where S_β is the set of isolated points of X_β .*

Proof. We proceed by induction on α . For $\alpha = 0$ it is clear. Let us assume that $\text{SSpec}_\beta(R) = \bigcup_{\gamma \leq \beta} S_\gamma$ for all $\beta < \alpha$. Now let $Q \in \bigcup_{\beta \leq \alpha} S_\beta$. If $Q \in S_\alpha$, then Q is a maximal element of X_α and so $Q' \in X, Q \subset Q'$ implies that $Q' \notin X_\alpha = X - \bigcup_{\beta < \alpha} S_\beta$. Hence we have $Q' \in S_\beta$ for some $\beta < \alpha$. Thus $Q' \in \bigcup_{\gamma \leq \beta} S_\gamma = \text{SSpec}_\beta(R)$ which implies that $Q \in \text{SSpec}_\alpha(R)$, and if $Q \notin S_\alpha$, then $Q \in S_\beta$ for some $\beta < \alpha$ which implies that $Q \in \bigcup_{\gamma \leq \beta} S_\gamma = \text{SSpec}_\beta(R) \subseteq \text{SSpec}_\alpha(R)$. Therefore, we have $\bigcup_{\beta \leq \alpha} S_\beta \subseteq \text{SSpec}_\alpha(R)$.

Conversely, let $Q \in \text{SSpec}_\alpha(R)$, then if $Q \notin \bigcup_{\beta < \alpha} S_\beta$, we show that $Q \in S_\alpha$. To this end, let $Q' \in X, Q \subset Q'$, then $Q' \in \text{SSpec}_\beta(R) = \bigcup_{\gamma \leq \beta} S_\gamma$ implies that $Q' \notin X_\alpha = X - \bigcup_{\gamma < \alpha} S_\gamma$. But $Q \in X_\alpha$ shows that Q must be a maximal element of X_α , so by the previous lemma $Q \in S_\alpha$. Therefore we have $\text{SSpec}_\alpha(R) \subseteq \bigcup_{\beta \leq \alpha} S_\beta$. \square

Corollary 3.5. *Let $\text{si.dim}(R) = \alpha$, then $X = \text{SSpec}(R)$ with V-topology have derived dimension and $d(X) \leq \alpha + 1$*

Proof. Let S be a non empty subset of X , then by Theorem 2.2, there exists a maximal element Q in S . We note that $V(Q) \cap S = \{Q\}$. This shows that Q is an isolated points of S with respect to V-topology. Hence by Lemma 3.2, there is an $\alpha > 0$ such that $X_\alpha = \emptyset$. Hence $d(X)$

exists and since according to the previous proposition $X_{\alpha+1} = X - \bigcup_{\beta \leq \alpha} S_\beta = \emptyset$, therefore we have $d(X) \leq \alpha + 1$. \square

The next result is our main theorem.

Theorem 3.6. *Let $X = \text{SSpec}(R)$ be the space with the V -topology, then the derived dimension of X exists if and only if $\text{si.dim}(R)$ exists and $d(X) = \text{si.dim}(R)$ if $d(X)$ is a limit ordinal and $d(X) = \text{si.dim}(R) + 1$ if $d(X)$ is not a limit ordinal.*

Proof. Since $\text{SSpec}_\alpha(R) = \bigcup_{\beta \leq \alpha} S_\beta$ and $X_{\alpha+1} = X - \bigcup_{\beta \leq \alpha} S_\beta$, so the first part holds. For the last part, first consider $d(X) = \alpha$, where α is a limit ordinal. In this case we have $X_\alpha = X - \bigcup_{\beta < \alpha} S_\beta = \emptyset$, therefore $X = \bigcup_{\beta < \alpha} S_\beta = \bigcup_{\beta \leq \alpha} S_\beta = \text{SSpec}_\alpha(R)$. Hence $\text{si.dim}(R) \leq \alpha$, and since by Corollary 3.5, $d(X) \leq \text{si.dim}(R) + 1$, thus $\text{si.dim}(R) = \alpha$. Now let $d(X) = \alpha + 1$, then $X_{\alpha+1} = \emptyset$ which implies that $X = \bigcup_{\beta \leq \alpha} S_\beta = \text{SSpec}_\alpha(R)$. Therefore $\text{si.dim}(R) \leq \alpha$ and from $d(X) \leq \text{si.dim}(R) + 1$, we get $\text{si.dim}(R) = \alpha$. \square

4. ACKNOWLEDGMENTS

We are grateful to the referees for their insightful comments and suggestions on the paper. In addition, the first author grateful to the Research Council of Shahid Chamran University of Ahvaz for financial support (GN: SCU.MM1402.258).

REFERENCES

- [1] A. Azizi, *Strongly irreducible ideals*, J. Aust. Math. Soc., **84** No. 2 (2008) 145-154.
- [2] K. R. Goodearl and R. B. Warfield, *An Introduction to Noncommutative Noetherian Rings*, Cambridge University Press, 2004.
- [3] J. Hashemi and F. Hassanzadeh, *Some remarks on strongly irreducible ideals*, J. Math. Ext., To appear.
- [4] O. A. S. Karamzadeh, *On the classical Krull dimension of rings*, Fund. Math., **117** No. 2 (1983) 103-108.
- [5] G. Krause, *On fully left bounded left noetherian rings*, J. Algebra., **23** No. 1 (1972) 88-99.
- [6] M. Namdari, *The story of rings of continuous functions in Ahvaz: From $C(X)$ to $C_c(X)$* , J. Iran. Math. Soc., **4** No. 2 (2023) 149-177.

Jamal Hashemi

Department of Mathematics,

Faculty of Mathematical Sciences and Computer,

Shahid Chamran University of Ahvaz, Ahvaz, Iran.

j.hashemi@scu.ac.ir

Fatemeh Hassanzadeh

Department of Mathematics,

Faculty of Mathematical Sciences and Computer,

Shahid Chamran University of Ahvaz, Ahvaz, Iran.

`f.hassanzade@stu.scu.ac.ir`